

Listed below are fees for STI tests offered at our clinic. Please note that Japanese National Health Insurance **does not** cover the costs of screening tests. Test results can be reported in person, by mail, phone, or email. Consumption tax (currently 8%) will be added to the final fee.

Test	Recommended MINIMUM interval prior to testing:	Results available in:	Fee
STI screening panel (HIV**, syphilis RPR, gonorrhea, chlamydia)	---	4 days	¥12,000
STI screening panel (HIV**, syphilis rapid TP antibody, gonorrhea, chlamydia)	---	4 days	¥13,000
**Either the lab or in-house HIV antigen/antibody test can be chosen for the panels. Screening for gonorrhea and chlamydia is routinely done on urine samples, but testing of other sites (oral, anal, vaginal) may be recommended depending on risk factors. There will be additional fees for each site tested. Mycoplasma/Ureaplasma testing can be included for an additional ¥7,000			
HIV antigen/antibody [†] (sample tested at outside lab)	2 weeks	1 – 2 days	¥4,000
HIV antigen/antibody [†] (rapid in-house blood test)	2 weeks	20 minutes	¥4,000
†4 th generation HIV-1 p24 antigen, HIV-1,2 antibody			
Syphilis (RPR)	3 weeks	2 days	¥1,000
Syphilis (TP antibody, rapid in-house blood test)	3 weeks	30 minutes	¥2,000
Chlamydia (PCR test on urine, gargle, or swab sample)	1 week	4 days	¥4,000
Gonorrhea (PCR test on urine, gargle, or swab sample)	1 day	4 days	¥4,000
Mycoplasma / Ureaplasma (PCR test on urine or swab sample)	1 week	7 days	¥10,000
Hepatitis B surface antigen	1 month	3 days	¥1,000
Hepatitis B surface antibody	4 to 6 months	3 days	¥2,000
Hepatitis C antibody	2 to 6 months	3 days	¥2,400
Herpes simplex 1 and 2 antigen (can only be done if skin lesions present)		4 days	¥6,000
Herpes simplex 1 and 2 antibody (NT) ^{***}	3 to 4 months	7 – 10 days	¥5,000

- A brief consultation with explanation of the tests is included in the fees. If extensive discussion is required or other medical issues need to be addressed, a consultation fee will be charged (¥5,000 - ¥8,000).
- There is no reliable screening test for “genital warts”. Diagnosis can only be made when skin lesions are present.

*STI = Sexually Transmitted Infections.

***Antibody testing for herpes is not generally recommended as the test does not accurately distinguish between the virus types.