



ELSEVIER

# How microRNAs control cell division, differentiation and death

Eric A Miska

After the milestone discovery of the first microRNA in 1993, the past five years have seen a phenomenal surge of interest in these short, regulatory RNAs. Given that 2% of all known human genes encode microRNAs, one main goal is to uncover microRNA function. Although it has been more difficult to assign function to microRNAs in animals than it has been in plants, important roles are emerging: in invertebrates, microRNAs control developmental timing, neuronal differentiation, tissue growth and programmed cell death. Functional studies in zebrafish and mice point toward important roles for microRNAs during morphogenesis and organogenesis. Finally, microRNAs might regulate viral infection and human cancer.

## Addresses

Wellcome Trust/Cancer Research UK Gurdon Institute and Department of Biochemistry, University of Cambridge, The Henry Wellcome Building of Cancer and Developmental Biology, Tennis Court Road, Cambridge, CB2 1QN, UK

Corresponding author: Miska, Eric A (e.miska@gurdon.cam.ac.uk)

**Current Opinion in Genetics & Development** 2005, **15**:563–568

This review comes from a themed issue on  
Differentiation and gene regulation  
Edited by Tony Kouzarides and Andrew J Bannister

Available online 15th August 2005

0959-437X/\$ – see front matter  
© 2005 Elsevier Ltd. All rights reserved.

DOI 10.1016/j.gde.2005.08.005

## Introduction

Interest in the genes controlling developmental timing in the roundworm *Caenorhabditis elegans* [1–3] led to the cloning of the first microRNA, *lin-4*, [4] and the identification of the first microRNA target, *lin-14* [5]. The cloning of a second microRNA, *let-7*, which is involved in the same pathway in *C. elegans* [6], and the realization that *let-7* was conserved from worms to mammals [7] led to the identification of many microRNAs in worms, flies, mammals and flowering plants, using cloning and prediction strategies [8–12]. MicroRNAs are short, ~22 nucleotide, non-coding RNAs that are thought to regulate gene expression through sequence-specific base-pairing with their targets. The precise number of microRNA genes in the human genome is still unknown, but current estimates range from 500 to 1000 [13]. This review focuses only on animal microRNAs; the roles of microRNAs in plants have been reviewed recently elsewhere [14,15].

## MicroRNA biogenesis and mechanisms of action

MicroRNA genes are found in intergenic regions as well as in introns. When located within introns, they can be on either the coding or the non-coding strand. Co-regulation of microRNAs and their host genes has been observed in some cases [16]. The majority of microRNA genes are probably transcribed by RNA polymerase II. Primary microRNA (pri-miRNA) transcripts can be hundreds of bases to several kilobases in length and might contain a single (monocistronic) or several (polycistronic) miRNAs. Pri-miRNAs have a 5' 7-methylguanosine cap and a 3'-polyadenosine tail, and they might or might not contain introns that are spliced. For a detailed review of microRNA transcription, see [17].

Pri-miRNAs contain ~80-nucleotide stem-loop structures, named pre-miRNAs, with the mature miRNAs located on at least one of their arms. In the nucleus, pre-miRNAs are processed from pri-miRNAs by the RNase III enzyme Drosha and DGCR8/Pasha [18–22] and are exported to the cytoplasm by a mechanism that involves Exportin-5 [23]. In the cytoplasm, a second RNase III enzyme, Dicer, cuts the pre-miRNA to generate the mature microRNA as part of a short RNA duplex. This RNA is subsequently unwound by a helicase activity and incorporated into an RNA-induced silencing complex (RISC). Detailed reviews of microRNA biogenesis can be found elsewhere [17,24–27].

The mature microRNA is chemically identical to small interfering RNAs (siRNAs; see Glossary) that are generated during RNA interference (RNAi; see Glossary); for example, in experimental conditions and during viral infection (Figure 1). Although most microRNAs in animals are thought to function through the inhibition of effective mRNA-translation of target genes through imperfect base-pairing with the 3'UTR (3' untranslated region) of target mRNAs [25], the underlying mechanism is poorly understood. In addition to this 'classic' mechanism, at least one microRNA, miR-196, can cleave a target mRNA, HOXB8, in a similar fashion to cleavage by siRNA [28\*,29]. MicroRNAs might also play a role in AU-rich element-mediated mRNA degradation [30\*]. Finally, the involvement of microRNAs in transcriptional gene-silencing, which has been observed in plants, remains a possibility [14,31].

This review describes the recent advances in understanding microRNA function in animals. As this is very much a work in progress, I also point out new strategies to further our understanding of microRNA biology. All those animal microRNAs for which a function has been demonstrated *in vivo* are listed in Table 1. Table 2 summarizes all

**Glossary**

**Exocytosis:** Release of proteins and neurotransmitters, etc, from cells by the fusion of vesicles with the plasma membrane.

**RNA interference (RNAi):** Sequence-specific mRNA-degradation pathway. Requires small interfering RNAs to target specific mRNAs.

**Small interfering RNAs (siRNA):** Short, single-stranded RNAs of approximately 22 nucleotides that can cause the degradation of target mRNAs by perfect base-pairing with their target.

**Target prediction:** Bioinformatic approach to predict microRNA targets on the basis of sequence complementarity of microRNAs and microRNA binding sites in 3'UTRs of target mRNAs.

microRNAs for which a biological role has been proposed and substantiated by indirect means.

**Roles of microRNAs in invertebrates**


The developmental timing (heterochronic) pathway in *C. elegans* involves at least two microRNAs, *lin-4* and *let-7*. Both *lin-4* and *let-7* are required for endothelial cell fate decisions during larval development [1,2]. When *lin-4* or *let-7* is inactivated, specific endothelial cells undergo additional cell divisions instead of their normal differentiation. The two known target genes of *lin-4*, *lin-14* and *lin-28*, were identified through genetic interaction studies (see Table 1) [4,5,32]. This is also the case for the first two known *let-7* target genes, *lin-41* and *hbl-1* [33–35]. More recently, two additional *let-7* target genes, the transcription factor genes *daf-12* and *pha-4*, were identified with a combination of target prediction (see Glossary) using bioinformatics and RNAi [36\*\*]. A *let-7*-related microRNA, miR-84, was recently identified as a putative regulator of vulval development in *C. elegans* (see Table 2)

[37]. miR-84 is proposed to act through *let-60*, a *RAS* homologue [37]. MicroRNAs are also regulators of neuronal differentiation in *C. elegans*. The microRNA *lgy-6* is required for asymmetrical expression of a putative taste-receptor in a pair of sensory neurons, ASEL and ASER [38]. The *lgy-6* microRNA target is the Nkx-type homeobox gene *cog-1*. Both *lgy-6* and *cog-1* were identified in forward genetic screens for loss of asymmetry of ASEL and ASER. Interestingly, a second microRNA, miR-273, might act upstream of *lgy-6* and *cog-0031* as a direct regulator of *die-1*, which also encodes a transcription factor [39].

The first microRNA to be identified in the fruit fly, *Drosophila melanogaster*, is encoded by the *bantam* locus, which had previously been identified in a screen for deregulated tissue growth [40]. The *bantam* microRNA stimulates cell proliferation and reduces programmed cell death. It directly regulates the pro-apoptotic gene *hid/wrinkled*. A second *D. melanogaster* microRNA, miR-14, also suppressed programmed cell death [41]. Its target(s) remain to be determined. Additional roles for microRNAs in *D. melanogaster* can be inferred from conserved regulatory motifs in the 3'UTRs of Notch target genes, termed GY-, Brd- (Bearded) and K-boxes, and from microRNAs with complementarity to these motifs [42]. Candidate microRNAs regulating these motifs include miR-2, miR-4, miR-5, miR-6, miR-7, miR-11 and miR-79 [42–45].

**Roles of microRNAs in vertebrates**

Little direct evidence for *in vivo* roles for vertebrate microRNAs has been obtained to date. Not a single

**Figure 1**

Multiple roles for short RNAs. Primary microRNA gene transcripts contain stem-loop structures (pre-miRNAs) that are processed by the RNase III enzyme Dicer to give rise to the mature microRNA. Dicer is also required for the processing of double-stranded viral RNA and for generating siRNAs during RNAi. Mature microRNAs and siRNAs are chemically indistinguishable. Short RNAs are thought to have three modes of action: translational inhibition, mRNA degradation and transcriptional gene-silencing at the level of chromatin. In animals, most microRNAs are thought to inhibit effective target mRNA translation; however, microRNAs might use all three mechanisms to regulate target-gene expression.

**Table 1****Animal microRNAs for which a function has been demonstrated *in vivo*.**

MicroRNA	Known targets	Organism	Biological process	Cellular process	References
<i>lin-4</i> microRNA	<i>lin-14, lin-28</i>	<i>C. elegans</i>	Developmental timing	Differentiation/proliferation	[4,5,32]
<i>let-7</i> microRNA	<i>lin-41, hbl-1, daf-12, pha-4</i>	<i>C. elegans</i>	Developmental timing	Differentiation/proliferation	[6,33–35,36**]
<i>lisy-6</i> microRNA	<i>cog-1</i>	<i>C. elegans</i>	Left-right asymmetry	Differentiation	[38]
<i>bantam</i> microRNA	<i>hid</i>	<i>D. melanogaster</i>	Growth control	Proliferation/programmed cell death	[40]
miR-14	Unknown	<i>D. melanogaster</i>	Programmed cell death	Programmed cell death	[41]
miR-430 <sup>1</sup>	Unknown	<i>Drosophila rerio</i>	Neurogenesis	Unknown	[46**]

This table includes only those microRNAs that have been analyzed using loss-of-function studies. Functional roles for a number of other microRNAs have been inferred from overexpression and mis-expression studies; these data are summarized in Table 2. <sup>1</sup>miR-430 function was inferred from rescue experiments in a *Dicer*-mutant background.

microRNA knockout has been reported. However, in zebrafish (*Danio rerio*) experiments in which both maternal and zygotic Dicer product was removed, re-introduction of a single microRNA, miR-430, led to a dramatic improvement of severe brain morphogenesis defects [46\*\*]. This observation suggests an important function for miR-430, or a related microRNA, in brain morphogenesis in the zebrafish.

In the mouse, miR-181 has been implicated in haematopoietic lineage differentiation, and overexpression of miR-181 leads to an increase in B lymphoid cells *in vitro* and *in vivo* [47]. miR-375, which is specifically expressed in mouse pancreatic islet cells, might regulate insulin secretion at the level of exocytosis (see Glossary) [48]. *Myotrophin* was identified as one direct target for miR-375.

Finally, the *miR-17* microRNA cluster and *miR-32* have been linked to human disease and are discussed below [49,50\*\*].

### A role for microRNAs in human disease: cancer and viral infection

A potential involvement of microRNAs in human disease might be inferred from the biological roles of microRNAs reported to date. Indeed, the *lin-4* and *let-7* phenotypes observed in *C. elegans* can be interpreted as proliferative

defects [1,2], and the roles of *bantam* and *miR-14* in *D. melanogaster* also point to defects in proliferation [40,41]. Therefore, one might speculate that microRNAs might be de-regulated in proliferative diseases, such as cancer. Indeed, a large number of microRNAs appear to be de-regulated in primary human tumours [50\*\*,51,52,53\*], and many human microRNAs are located at fragile sites and genomic regions involved in cancer [54,55]. Of particular interest is the *miR-17* microRNA cluster, which is in a region on human chromosome 13 that is frequently amplified in B-cell lymphomas [50\*\*]. Over-expression of the *miR-17* cluster was found to co-operate with *c-Myc* to accelerate tumour development in a mouse B-cell lymphoma model. Although the direct targets of members of the *miR-17* cluster of microRNAs are unknown, their collective role might be to inhibit excessive programmed cell death [50\*\*]. In a separate study, the *miR-17* cluster was found to be up-regulated by over-expression of c-Myc, leading to the de-regulation of E2F1 expression [56\*].

Another link between microRNAs and human disease comes from the identification of microRNAs encoded by large DNA viruses of the herpesvirus family, including the Epstein-Barr virus [57,58\*]. These viral microRNAs have no apparent homologues in host genomes, and their function is currently not understood. Conversely, a cellular microRNA, miR-32, was shown to play an important

**Table 2****Proposed *in vivo* roles for additional microRNAs in animals.**

MicroRNA	Targets	Organism	Biological process	Cellular process	References
miR-273	<i>die-1</i>	<i>C. elegans</i>	Left-right asymmetry	Differentiation	[49]
miR-84	<i>let-60</i>	<i>C. elegans</i>	Vulval development	Differentiation/proliferation	[37]
miR-2a, -2b, -6, -7	<i>E(spl)/bHLH, Bearded families</i> <sup>1</sup>	<i>D. melanogaster</i>	Notch signalling	Unknown	[42–45]
miR-181	Unknown	<i>Mus musculus</i>	Hematopoiesis	Differentiation	[47]
miR-375	<i>Myotrophin (Mtpn)</i>	<i>M. musculus</i>	Insulin secretion	Exocytosis	[48]
miR-17, -18, -19a, -20, -19b-1, -92-1	Unknown	<i>M. musculus</i>	Tumourigenesis	Proliferation	[50**]
miR-32	<i>PFV-1</i>	<i>Homo sapiens</i>	Viral defense	Viral defense	[49]

This table lists microRNAs for which a probable function has been demonstrated using an indirect approach, for example mis-expression experiments. This table does not contain microRNAs for which a target mRNA has been predicted and validated using over-expression experiments but for which no further functional characterization at the cellular or organismal level has been carried out. Additional validated microRNA–target pairs can be found elsewhere [36\*\*,43–45,70\*]. <sup>1</sup>Family of Notch target transcription factors with GY-, Brd- and K-box motifs. Abbreviations: bHLH, basis helix–loop–helix; PFV-1, primate foamy virus type 1.

role in regulating the proliferation of the primate foamy virus type 1 (PFV-1) in cell culture [49].

### Approximating microRNA function through the analysis of *Dicer* mutants

The RNase III enzyme Dicer is essential for the processing of microRNAs and for RNAi. Therefore, the analysis of *Dicer*-knockout strains might point toward biological roles of microRNAs, with the caveat that RNAi and possibly other biological processes might be de-regulated. Dicer was first analyzed genetically in *C. elegans* (*dcr-1*), in which it was found to be essential for germline development [59]. It is likely that maternal contribution of *dcr-1* masks earlier phenotypes, and this notion is supported by additional phenotypes that are revealed by the inactivation of *dcr-1* using RNAi [60]. RNAi of *dcr-1* results in embryonic and developmental timing defects that are reminiscent of *lin-4* and *let-7* mutants [60], thereby suggesting additional roles for microRNAs in *C. elegans* development.

In the zebrafish, a knockout of the *Dicer1* gene leads to a developmental arrest 7 to 10 days post-fertilization [61]. An earlier role for Dicer is probably masked by maternal Dicer contribution. Indeed, removal of maternal Dicer by the generation of germ line clones leads to more severe defects: axis formation and early differentiation are normal, but gastrulation, brain formation, somitogenesis and heart development are abnormal [46\*\*].

*Dicer*-mutant mice have defects in axis formation and gastrulation, are depleted of stem cells and die at around 7.5 days of gestation [62]. A conditional knockout of *Dicer* early during T-cell development suggests a role for microRNAs in  $\alpha\beta$ -cell lineage [63]. Finally, a knockout of *Dicer* in the chicken-human hybrid DT40 cell line suggests a role for the enzyme in heterochromatin formation [64].

### Approximating microRNA function through mRNA target prediction

MicroRNA target prediction offers an attractive route to the discovery of microRNA function. Several algorithms have been applied to *C. elegans*, *D. melanogaster* and mammalian genomes to predict microRNA targets [43,44,65–69]. In a complementary effort, conserved sites in 3'UTRs of human genes have been identified [70\*]. For a subset of these predictions, the use of overexpression and green fluorescent protein- or luciferase-reporter assays has demonstrated that the microRNA can regulate the target candidate. It remains an open question as to how many microRNA targets that have been verified in these heterologous systems are important targets *in vivo*.

### Approximating microRNA function through expression analysis

Initially, microRNA expression was monitored using northern blotting [4]. Next, cloning of microRNAs from specific tissues was used to approximate expression levels

[71]. Over the past two years, microRNA microarray technologies have been developed for the high-throughput profiling of microRNA expression [15,52,53\*,72,73]. The most exciting discovery to date is that many microRNAs are tissue-specifically expressed. It is to be expected that microRNA expression profiles will help to guide functional studies, as exemplified by the analysis of the pancreatic islet-specific microRNA miR-340 [48]. Finally, the recent development of microRNA *in situ* technology has the potential to revolutionize microRNA expression analysis [74\*\*].

### Conclusions

We are just beginning to unravel the roles of microRNAs in animals, but the examples listed here already demonstrate the great importance of this class of short RNAs. Many of the functions elucidated to date point to important roles for microRNAs in cell fate determination, including the control of cell division, differentiation and death. Few loss-of-function studies have been reported to date, but the tool kit is expanding with devices for global de-regulation of microRNAs (*Dicer* knockouts), target predictions and expression patterns at our disposal. The most exciting time for microRNAs lies ahead.

### Acknowledgements

E Miska acknowledges support from a Cancer Research UK Programme Grant (C13474/A4613) and would like to thank Greg Hannon for sharing information before publication.

### Papers and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Ambros V, Horvitz HR: **Heterochronic mutants of the nematode *Caenorhabditis elegans***. *Science* 1984, **226**:409-416.
2. Chalfie M, Horvitz HR, Sulston JE: **Mutations that lead to reiterations in the cell lineages of *C. elegans***. *Cell* 1981, **24**:59-69.
3. Horvitz HR, Sulston JE: **Isolation and genetic characterization of cell-lineage mutants of the nematode *Caenorhabditis elegans***. *Genetics* 1980, **96**:435-454.
4. Lee RC, Feinbaum RL, Ambros V: **The *C. elegans* heterochronic gene *lin-4* encodes small RNAs with antisense complementarity to *lin-14***. *Cell* 1993, **75**:843-854.
5. Wightman B, Ha I, Ruvkun G: **Posttranscriptional regulation of the heterochronic gene *lin-14* by *lin-4* mediates temporal pattern formation in *C. elegans***. *Cell* 1993, **75**:855-862.
6. Reinhart BJ, Slack FJ, Basson M, Pasquinelli AE, Bettinger JC, Rougvie AE, Horvitz HR, Ruvkun G: **The 21-nucleotide *let-7* RNA regulates developmental timing in *Caenorhabditis elegans***. *Nature* 2000, **403**:901-906.
7. Pasquinelli AE, Reinhart BJ, Slack F, Martindale MQ, Kuroda MI, Maller B, Hayward DC, Ball EE, Degnan B, Muller P *et al.*: **Conservation of the sequence and temporal expression of *let-7* heterochronic regulatory RNA**. *Nature* 2000, **408**:86-89.
8. Lee RC, Ambros V: **An extensive class of small RNAs in *Caenorhabditis elegans***. *Science* 2001, **294**:862-864.

9. Lagos-Quintana M, Rauhut R, Lendeckel W, Tuschl T: **Identification of novel genes coding for small expressed RNAs.** *Science* 2001, **294**:853-858.
10. Lau NC, Lim LP, Weinstein EG, Bartel DP: **An abundant class of tiny RNAs with probable regulatory roles in *Caenorhabditis elegans*.** *Science* 2001, **294**:858-862.
11. Reinhart BJ, Weinstein EG, Rhoades MW, Bartel B, Bartel DP: **MicroRNAs in plants.** *Genes Dev* 2002, **16**:1616-1626.
12. Llave C, Kasschau KD, Rector MA, Carrington JC: **Endogenous and silencing-associated small RNAs in plants.** *Plant Cell* 2002, **14**:1605-1619.
13. Berezikov E, Guryev V, van de Belt J, Wienholds E, Plasterk RH, Cuppen E: **Phylogenetic shadowing and computational identification of human microRNA genes.** *Cell* 2005, **120**:21-24.
14. Kidner CA, Martienssen RA: **The developmental role of microRNA in plants.** *Curr Opin Plant Biol* 2005, **8**:38-44.
15. Baskerville S, Bartel DP: **Microarray profiling of microRNAs reveals frequent coexpression with neighboring miRNAs and host genes.** *RNA* 2005, **11**:241-247.
16. Zilberman D, Henikoff S: **Epigenetic inheritance in *Arabidopsis*: selective silencing.** *Curr Opin Genet Dev* 15: in press.
17. Cullen BR: **Transcription and processing of human microRNA precursors.** *Mol Cell* 2004, **16**:861-865.
18. Lee Y, Ahn C, Han J, Choi H, Kim J, Yim J, Lee J, Provost P, Radmark O, Kim S *et al.*: **The nuclear RNase III Droscha initiates microRNA processing.** *Nature* 2003, **425**:415-419.
19. Denli AM, Tops BB, Plasterk RH, Ketting RF, Hannon GJ: **Processing of primary microRNAs by the microprocessor complex.** *Nature* 2004, **432**:231-235.
20. Gregory RI, Yan KP, Amuthan G, Chendrimada T, Doratotaj B, Cooch N, Shiekhattar R: **The microprocessor complex mediates the genesis of microRNAs.** *Nature* 2004, **432**:235-240.
21. Han J, Lee Y, Yeom KH, Kim YK, Jin H, Kim VN: **The Droscha-DGCR8 complex in primary microRNA processing.** *Genes Dev* 2004, **18**:3016-3027.
22. Landthaler M, Yalcin A, Tuschl T: **The human DiGeorge syndrome critical region gene 8 and its *D. melanogaster* homolog are required for miRNA biogenesis.** *Curr Biol* 2004, **14**:2162-2167.
23. Lund E, Guttinger S, Calado A, Dahlberg JE, Kutay U: **Nuclear export of microRNA precursors.** *Science* 2003, **303**:95-98.
24. Pham JW, Sontheimer EJ: **The making of an siRNA.** *Mol Cell* 2004, **15**:163-164.
25. Bartel DP: **MicroRNAs: genomics, biogenesis, mechanism, and function.** *Cell* 2004, **116**:281-297.
26. Tomari Y, Zamore PD: **MicroRNA biogenesis: drosha can't cut it without a partner.** *Curr Biol* 2005, **15**:R61-R64.
27. Pasquinelli AE, Hunter S, Bracht J: **MicroRNAs: a developing story.** *Curr Opin Genet Dev* 15:200-205.
28. Yekta S, Shih IH, Bartel DP: **MicroRNA-directed cleavage of HOXB8 mRNA.** *Science* 2004, **304**:594-596.  
 The authors describe how a mammalian microRNA, miR-196, can direct the cleavage of the HOXB8 mRNA.
29. Mansfield JH, Harfe BD, Nissen R, Obenaus J, Srineel J, Chaudhuri A, Farzan-Kashani R, Zuker M, Pasquinelli AE, Ruvkun G *et al.*: **MicroRNA-responsive 'sensor' transgenes uncover Hox-like and other developmentally regulated patterns of vertebrate microRNA expression.** *Nat Genet* 2004, **36**:1079-1083.
30. Jing Q, Huang S, Guth S, Zarubin T, Motoyama A, Chen J, Di Padova F, Lin SC, Gram H, Han J: **Involvement of microRNA in AU-rich element-mediated mRNA instability.** *Cell* 2005, **120**:623-634.  
 Here, the authors describe a possible new role for microRNAs in the control of AU-rich element-mediated mRNA decay, implicating miR-16.
31. Baulcombe D: **RNA silencing in plants.** *Nature* 2004, **431**:356-363.
32. Moss EG, Lee RC, Ambros V: **The cold shock domain protein LIN-28 controls developmental timing in *C. elegans* and is regulated by the *lin-4* RNA.** *Cell* 1997, **88**:637-646.
33. Slack FJ, Bassom M, Liu Z, Ambros V, Horvitz HR, Ruvkun G: **The *lin-41* RBCC gene acts in the *C. elegans* heterochronic pathway between the *let-7* regulatory RNA and the LIN-29 transcription factor.** *Mol Cell* 2000, **5**:659-669.
34. Abrahamte JE, Daul AL, Li M, Volk ML, Tennessen JM, Miller EA, Rougvie AE: **The *Caenorhabditis elegans* hunchback-like gene *lin-57/hbl-1* controls developmental time and is regulated by microRNAs.** *Dev Cell* 2003, **4**:625-637.
35. Lin SY, Johnson SM, Abraham M, Vella MC, Pasquinelli A, Gamberi C, Gottlieb E, Slack FJ: **The *C. elegans* hunchback homolog, *hbl-1*, controls temporal patterning and is a probable microRNA target.** *Dev Cell* 2003, **4**:639-650.
36. Grosshans H, Johnson T, Reinert KL, Gerstein M, Slack FJ: **The temporal patterning microRNA *let-7* regulates several transcription factors at the larval to adult transition in *C. elegans*.** *Dev Cell* 2005, **8**:321-330.  
 The authors identify at least two new direct *let-7* target mRNAs, including the transcription factors genes *daf-12* and *pha-4*.
37. Johnson SM, Grosshans H, Shingara J, Byrom M, Jarvis R, Cheng A, Labourier E, Reinert KL, Brown D, Slack FJ: **RAS is regulated by the *let-7* microRNA family.** *Cell* 2005, **120**:635-647.
38. Johnston RJ, Hobert O: **A microRNA controlling left/right neuronal asymmetry in *Caenorhabditis elegans*.** *Nature* 2003, **426**:845-849.
39. Chang S, Johnston RJ Jr, Frokjaer-Jensen C, Lockery S, Hobert O: **MicroRNAs act sequentially and asymmetrically to control chemosensory laterality in the nematode.** *Nature* 2004, **430**:785-789.
40. Brennecke J, Hipfner DR, Stark A, Russell RB, Cohen SM: ***bantam* encodes a developmentally regulated microRNA that controls cell proliferation and regulates the proapoptotic gene *hid* in *Drosophila*.** *Cell* 2003, **113**:25-36.
41. Xu P, Vernoooy SY, Guo M, Hay BA: **The *Drosophila* microRNA mir-14 suppresses cell death and is required for normal fat metabolism.** *Curr Biol* 2003, **13**:790-795.
42. Lai EC: **Micro RNAs are complementary to 3' UTR sequence motifs that mediate negative post-transcriptional regulation.** *Nat Genet* 2002, **30**:363-364.
43. Stark A, Brennecke J, Russell RB, Cohen SM: **Identification of *Drosophila* microRNA targets.** *PLoS Biol* 2003, **1**:e60.
44. Brennecke J, Stark A, Russell RB, Cohen SM: **Principles of microRNA-target recognition.** *PLoS Biol* 2005, **3**:e85.
45. Lai EC, Tam B, Rubin GM: **Pervasive regulation of *Drosophila* Notch target genes by GY-box-, Brd-box-, and K-box-class microRNAs.** *Genes Dev* 2005, **19**:1067-1080.
46. Giraldez AJ, Cinalli RM, Glasner ME, Enright AJ, Thomson JM, Baskerville S, Hammond SM, Bartel DP, Schier AF: **MicroRNAs regulate brain morphogenesis in zebrafish.** *Science* 2005, **308**:833-838.  
 The authors' analysis of a maternal and zygotic *Dicer*-mutant zebrafish points to important roles for microRNAs in gastrulation and the development of brain, heart and somites. This study points toward a direct role for members of the miR-430 family of microRNAs in brain development.
47. Chen CZ, Li L, Lodish HF, Bartel DP: **MicroRNAs modulate hematopoietic lineage differentiation.** *Science* 2004, **303**:83-86.
48. Poy MN, Eliasson L, Krutzfeldt J, Kuwajima S, Ma X, Macdonald PE, Pfeffer S, Tuschl T, Rajewsky N, Rorsman P *et al.*: **A pancreatic islet-specific microRNA regulates insulin secretion.** *Nature* 2004, **432**:226-230.
49. Lecellier CH, Dunoyer P, Arar K, Lehmann-Che J, Eyquem S, Himber C, Saib A, Voinnet O: **A cellular microRNA mediates antiviral defense in human cells.** *Science* 2005, **308**:557-560.

50. He L, Thomson JM, Hemann MT, Hernando-Monge E, Mu D, Goodson S, Powers S, Cordon-Cardo C, Lowe SW, Hannon GJ *et al.*: **A microRNA polycistron as a potential human oncogene.** *Nature* 2005, **435**:828-833.
- Using a mouse *c-Myc* co-cooperativity model, the authors demonstrate the first functional link between a cluster of microRNAs and cancer.
51. Calin GA, Dumitru CD, Shimizu M, Bichi R, Zupo S, Noch E, Aldler H, Rattan S, Keating M, Rai K *et al.*: **Frequent deletions and down-regulation of micro-RNA genes *miR15* and *miR16* at 13q14 in chronic lymphocytic leukaemia.** *Proc Natl Acad Sci USA* 2002, **99**:15524-15529.
52. Calin GA, Liu CG, Sevignani C, Ferracin M, Felli N, Dumitru CD, Shimizu M, Cimmino A, Zupo S, Dono M *et al.*: **MicroRNA profiling reveals distinct signatures in B cell chronic lymphocytic leukemias.** *Proc Natl Acad Sci USA* 2004, **101**:11755-11760.
53. Lu J, Getz G, Miska EA, Alvarez-Saavedra E, Lamb J, Peck D, Sweet-Cordero A, Ebert BL, Mak RH, Ferrando AA *et al.*: **MicroRNA expression profiles classify human cancers.** *Nature* 2005, **435**:834-838.
- Many microRNAs may be regulated in primary human tumours, suggesting a role in human cancer.
54. Calin GA, Sevignani C, Dumitru CD, Hyslop T, Noch E, Yendamuri S, Shimizu M, Rattan S, Bullrich F, Negrini M *et al.*: **Human microRNA genes are frequently located at fragile sites and genomic regions involved in cancers.** *Proc Natl Acad Sci USA* 2004, **101**:2999-3004.
55. McManus MT: **MicroRNAs and cancer.** *Semin Cancer Biol* 2003, **13**:253-258.
56. O'Donnell KA, Wentzel EA, Zeller KI, Dang CV, Mendell JT: **c-Myc-regulated microRNAs modulate E2F1 expression.** *Nature* 2005, **435**:839-843.
- The authors provide the first demonstration of a transcription factor directly regulating a microRNA primary transcript.
57. Pfeffer S, Sewer A, Lagos-Quintana M, Sheridan R, Sander C, Grasser FA, van Dyk LF, Ho CK, Shuman S, Chien M *et al.*: **Identification of microRNAs of the herpesvirus family.** *Nat Methods* 2005, **2**:269-276.
58. Pfeffer S, Zavolan M, Grasser FA, Chien M, Russo JJ, Ju J, John B, Enright AJ, Marks D, Sander C *et al.*: **Identification of virus-encoded microRNAs.** *Science* 2004, **304**:734-736.
- The authors identify microRNAs that are encoded by the Epstein-Barr virus, a member of the herpesvirus family. This suggests a role for microRNA in viral infection.
59. Knight SW, Bass BL: **A role for the RNase III enzyme DCR-1 in RNA interference and germ line development in *Caenorhabditis elegans*.** *Science* 2001, **293**:2269-2271.
60. Grishok A, Pasquinelli AE, Conte D, Li N, Parrish S, Ha I, Baillie DL, Fire A, Ruvkun G, Mello CC: **Genes and mechanisms related to RNA interference regulate expression of the small temporal RNAs that control *C. elegans* developmental timing.** *Cell* 2001, **106**:23-34.
61. Wienholds E, Koudijs MJ, van Eeden FJ, Cuppen E, Plasterk RH: **The microRNA-producing enzyme Dicer1 is essential for zebrafish development.** *Nat Genet* 2003, **35**:217-218.
62. Bernstein E, Kim SY, Carmell MA, Murchison EP, Alcorn H, Li MZ, Mills AA, Elledge SJ, Anderson KV, Hannon GJ: **Dicer is essential for mouse development.** *Nat Genet* 2003, **35**:215-217.
63. Cobb BS, Nesterova TB, Thompson E, Hertweck A, O'Connor E, Godwin J, Wilson CB, Brockdorff N, Fisher AG, Smale ST *et al.*: **T cell lineage choice and differentiation in the absence of the RNase III enzyme Dicer.** *J Exp Med* 2005, **201**:1367-1373.
64. Fukagawa T, Nogami M, Yoshikawa M, Ikeno M, Okazaki T, Takami Y, Nakayama T, Oshimura M: **Dicer is essential for formation of the heterochromatin structure in vertebrate cells.** *Nat Cell Biol* 2004, **6**:784-791.
65. Kiriakidou M, Nelson PT, Kouranov A, Fitziev P, Bouyioukos C, Mourelatos Z, Hatzigeorgiou A: **A combined computational-experimental approach predicts human microRNA targets.** *Genes Dev* 2004, **18**:1165-1178.
66. Rajewsky N, Socci ND: **Computational identification of microRNA targets.** *Dev Biol* 2004, **267**:529-535.
67. John B, Enright AJ, Aravin A, Tuschl T, Sander C, Marks DS: **Human microRNA targets.** *PLoS Biol* 2004, **2**:e363.
68. Lewis BP, Burge CB, Bartel DP: **Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets.** *Cell* 2005, **120**:15-20.
69. Lewis BP, Shih IH, Jones-Rhoades MW, Bartel DP, Burge CB: **Prediction of mammalian microRNA targets.** *Cell* 2003, **115**:787-798.
70. Xie X, Lu J, Kulbokas EJ, Golub TR, Mootha V, Lindblad-Toh K, Lander ES, Kellis M: **Systematic discovery of regulatory motifs in human promoters and 3' UTRs by comparison of several mammals.** *Nature* 2005, **434**:338-345.
- The authors take a complementary approach to the search for microRNA targets. They perform a systematic screen for motifs in 3'UTRs of human genes that are conserved in mammals and identify many sequences with complementarity to microRNAs.
71. Lagos-Quintana M, Rauhut R, Yalcin A, Meyer J, Lendeckel W, Tuschl T: **Identification of tissue-specific microRNAs from mouse.** *Curr Biol* 2002, **12**:735-739.
72. Miska EA, Alvarez-Saavedra E, Townsend M, Yoshii A, Sestan N, Rakic P, Constantine-Paton M, Horvitz HR: **Microarray analysis of microRNA expression in the developing mammalian brain.** *Genome Biol* 2004, **5**:R68.
73. Thomson JM, Parker J, Perou CM, Hammond SM: **A custom microarray platform for analysis of microRNA gene expression.** *Nat Methods* 2004, **1**:47-53.
74. Wienholds E, Kloosterman WP, Miska EA, Alvarez-Saavedra E, Berezikov E, de Bruijn E, Horvitz HR, Kauppinen S, Plasterk RHA: **MicroRNA expression in zebrafish embryonic development.** *Science* 2005, **309**:310-311.
- Here, the first robust microRNA *in situ* technology is demonstrated on zebrafish embryos. This technique offers unprecedented spatial resolution of microRNA expression and has the potential to revolutionize the identification of microRNA function.

### Free journals for developing countries

The WHO and six medical journal publishers have launched the Access to Research Initiative, which enables nearly 70 of the world's poorest countries to gain free access to biomedical literature through the Internet.

Gro Harlem Brundtland, director-general for the WHO, said that this initiative was 'perhaps the biggest step ever taken towards reducing the health information gap between rich and poor countries'.

For more information, visit [www.healthinternetwork.net](http://www.healthinternetwork.net)