

Social Neuroscience

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/psns20>

Resting state glutamate predicts elevated pre-stimulus alpha during self-relatedness - A combined EEG-MRS study on 'rest-self overlap'

Bai Yu^a, Takashi Nakao^b, Jiameng Xu^{cegh}, Pengmin Qin^{cde}, Pedro Chaves^c, Alexander Heinzl^k, Niall Duncan^{ci}, Timothy Lane^{defg}, Nai-Shing Yen^{gh}, Shang-Yueh Tsai^{gj} & Georg Northhoff^{cdegi}

^a Department of Psychology, Nagoya University, Japan

^b Department of Psychology, Graduate School of Education, Hiroshima University, Higashi-Hiroshima, Japan

^c Institute of Mental Health Research, University of Ottawa, Ottawa, Canada

^d Taipei Medical University, Graduate Institute of Humanities in Medicine, Taipei, Taiwan

^e Taipei Medical University-Shuang Ho Hospital, Brain and Consciousness Research Center, Depts. of Psychiatry and Radiology, New Taipei City, Taiwan

^f Academia Sinica, Institute of European and American Studies, Taipei, Taiwan

^g National Chengchi University, Research Center for Mind, Brain and Learning, Taipei, Taiwan

^h National Chengchi University, Department of Psychology, Taipei, Taiwan

ⁱ Centre for Cognition and Brain Disorders (CBBB), Normal University Hangzhou, Hangzhou, China

^j National Chengchi University, Graduate Institute of Applied Physics, Taipei, Taiwan

^k Department of Nuclear Medicine, RWTH Aachen, Germany

Accepted author version posted online: 24 Jul 2015.



[Click for updates](#)

To cite this article: Bai Yu, Takashi Nakao, Jiameng Xu, Pengmin Qin, Pedro Chaves, Alexander Heinzl, Niall Duncan, Timothy Lane, Nai-Shing Yen, Shang-Yueh Tsai & Georg Northhoff (2015): Resting state glutamate predicts elevated pre-stimulus alpha during self-relatedness - A combined EEG-MRS study on 'rest-self overlap', *Social Neuroscience*, DOI: [10.1080/17470919.2015.1072582](https://doi.org/10.1080/17470919.2015.1072582)

To link to this article: <http://dx.doi.org/10.1080/17470919.2015.1072582>

Disclaimer: This is a version of an unedited manuscript that has been accepted for publication. As a service to authors and researchers we are providing this version of the accepted manuscript (AM). Copyediting, typesetting, and review of the resulting proof will be undertaken on this manuscript before final publication of the Version of Record (VoR). During production and pre-press, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal relate to this version also.

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for

any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Publisher: Taylor & Francis

Journal: *Social Neuroscience*

DOI: 10.1080/17470919.2015.1072582

Resting state glutamate predicts elevated pre-stimulus alpha during self-relatedness – A combined EEG-MRS study on ‘rest-self overlap’

Bai Yu (1) *, Takashi Nakao (2)*, Jiameng Xu (3, 5, 7, 8), Pengmin Qin (3, 4, 5), Pedro Chaves (3), Alexander Heinzl (11), Niall Duncan (3, 9), Timothy Lane (4, 5, 6, 7), Nai-Shing Yen (7, 8), Shang-Yueh Tsai (7,10), Georg Northoff (3, 4, 5, 7, 9)

* = Equal contributing authors

(1) Department of Psychology, Nagoya University, Japan

(2) Department of Psychology, Graduate School of Education, Hiroshima University, Higashi-Hiroshima, Japan

(3) Institute of Mental Health Research, University of Ottawa, Ottawa, Canada

(4) Taipei Medical University, Graduate Institute of Humanities in Medicine, Taipei, Taiwan

(5) Taipei Medical University-Shuang Ho Hospital, Brain and Consciousness Research Center, Depts. of Psychiatry and Radiology, New Taipei City, Taiwan

(6) Academia Sinica, Institute of European and American Studies, Taipei, Taiwan

(7) National Chengchi University, Research Center for Mind, Brain and Learning, Taipei, Taiwan

(8) National Chengchi University, Department of Psychology, Taipei, Taiwan

(9) Centre for Cognition and Brain Disorders (CBBB), Normal University Hangzhou, Hangzhou, China

(10) National Chengchi University, Graduate Institute of Applied Physics, Taipei, Taiwan

(11) Department of Nuclear Medicine, RWTH Aachen, Germany

Corresponding authors

Dr. Georg Northoff, MD, PhD, FRCP

Research Director

Mind, Brain Imaging and Neuroethics
Canada Research Chair
EJLB-Michael Smith Chair for Neuroscience and Mental Health
Royal Ottawa Healthcare Group
University of Ottawa Institute of Mental Health Research

1145 Carling Avenue, Room 6467

Ottawa, ON K1Z 7K4

Tel.: (613) 722-6521 Ext. 6801

Fax: (613) 798-2982

Email: georg.northoff@theroyal.ca

Website: www.georgnorthoff.com

Tim Lane, PhD

Taipei Medical University, Graduate Institute of Humanities in Medicine, Taipei, Taiwan

Taipei Medical University-Shuang Ho Hospital, Brain and Consciousness Research Center, New
Taipei City, Taiwan

timlane13@gmail.com

Acknowledgments

GN is grateful for financial support from CIHR, EJLB-CIHR, and ISAN/HDRF. This work was also supported by JSPS KAKENHI Grant Numbers 24390284, 25870467, and 26285168 to TN. TL and NY are grateful for financial support from Taiwan's National Science Council (Grant numbers: 100-2410-H-004-139-MY3 and 102-2420-H-038-002-MY2.). JX is grateful for financial support provided by National Chengchi University's Research Center for Mind, Brain, and Learning.

Abstract

Recent studies have demonstrated neural overlap between resting state activity and self-referential processing. This “rest-self” overlap occurs especially in anterior cortical midline structures like the perigenual anterior cingulate cortex (PACC). However, the exact neuro-temporal and biochemical mechanisms remain to be identified. Therefore, we conducted a combined EEG-MRS study. EEG focused on pre-stimulus (e.g., prior to stimulus presentation or perception) power changes to assess the degree to which those changes can predict subjects’ perception (and judgment) of subsequent stimuli as high or low self-related. Magnetic resonance spectroscopy (MRS) measured resting state concentration of glutamate, focusing on PACC. High pre-stimulus (e.g., prior to stimulus presentation or perception) alpha power significantly correlated with both perception of stimuli judged to be highly self-related, and with resting state glutamate concentrations in the PACC. In sum, our results show (i) pre-stimulus (e.g., prior to stimulus presentation or perception) alpha power and resting state glutamate concentration to mediate rest-self overlap which (ii) dispose or incline subjects to assign high degrees of self-relatedness to perceptual stimuli.

Keywords:

Self-referential processing, perigenual anterior cingulate cortex, MRS, EEG, glutamine

Introduction

The concept of “self” has been investigated by philosophers for millennia; recently, however, it has also become a major topic in neuroscience. Imaging studies that investigate the self-relatedness of stimuli reveal strong neural recruitment of subcortical and cortical midline structures (CMS) during presentation of stimuli specifically related to self (Northoff et al., 2006; Qin and Northoff, 2011). These regions include the ventromedial prefrontal cortex (VMPFC), the perigenual anterior cingulate (PACC), as well as subcortical regions like the ventral striatum (VS) and the ventral tegmental area (VTA). This set of midline regions also forms the core of the default-mode network (DMN), which has been associated with particularly high resting state activity (Raichle et al., 2001; Buckner et al., 2008).

Importantly, several studies (Schneider et al., 2008; d’Argembeau et al. 2005; Whitfield-Gabriel et al., 2011; Qin et al. 2011; Qin et al., 2013) discovered overlap between resting state activity and self-related activity in anterior CMS like the perigenual anterior cingulate cortex (PACC). In these regions, high self-related stimuli seem to correlate with less deviation from resting state activity, relative to low self-related activity. But the mechanisms of this neural overlap between resting state and self-related activity, what we have dubbed the *rest-self overlap*, remain unclear.

The brain’s resting state activity or intrinsic activity has come increasingly into focus recently (Northoff 2014a and b). The term ‘resting state activity’ is paradoxical though since it denotes exactly the opposite namely that the brain is never at rest. While a continuous level at rest can be found throughout the whole brain in any region, resting state activity in default-mode network and especially the midline regions seems to show some peculiarities in that it is particularly high and variable (see Raichle et al. 2001, Northoff 2014a and b). How though do these specific features of the midline

regions' resting state activity impact subsequent stimulus-related activity during the processing of especially self-related stimuli? We currently do not know. The observed rest-self overlap suggest special relationship between for instance pre-stimulus activity levels (that somehow in a yet unclear way must be related to the task-free the resting state) and stimulus-induced activity during high self-related stimuli.

Several studies have combined fMRI and EEG in demonstrating that pre-stimulus activity can indeed predict neural and behavioural effects of subsequent stimuli and tasks (for overviews see Northoff et al. 2010; Sadargahini et al., 2010). For example, the level of pre-stimulus (e.g., prior to stimulus presentation or perception) activity in the fusiform face area, as measured with fMRI, predicts both the degree of neural activity during subsequent stimulus presentation and whether a given stimulus is perceived as a face or a vase (for an excellent overview, see Sadargahini et al 2010; for similar results, albeit with different tasks, see Shibata et al., 2008; Park and Rugg, 2010; Hsieh et al., 2012). Several analogous investigations have been carried out using EEG: here power or phase patterns of the resting state, or pre-stimulus (e.g., prior to stimulus presentation or perception) activity, predicts neural and behavioural stimulus-related activity during decision making and perceptual tasks (Linkankaer-Hansen, 2004; Hanslmayr et al., 2007; van Dijk et al., 2008; Ai & Ro, 2014; Nakao et al. 2013; Weisz et al., 2014).

But whether pre-stimulus (e.g., prior to stimulus presentation or perception) or resting state activity enables prediction of how subjects will respond to *self-related stimuli* has not yet been investigated. Because of the observed rest-self overlap, and reasoning by analogy, we conjectured that pre-stimulus (e.g., prior to stimulus presentation or perception) activity can be employed to predict neural activity when subjects perceive or render judgments about self-related stimuli. In other words, the working hypothesis is—pre-stimulus (e.g., prior to stimulus presentation or perception) or resting state activity may enable us to predict whether the subject perceives or judges any given stimulus as

self-related, on a scale that ranges from high to low. Investigating high self-related stimuli, several EEG studies showed changes around 150-400 ms and especially lower (8-9Hz) alpha band power changes post-stimulus (Mu et al., 2008; Justen et al., 2014; Qin et al., 2008; and, for a recent review on EEG studies of self, Knyazev, 2013). These findings, however, leave untested the question as to whether pre-stimulus or resting state activity affects self-related stimulus-induced activity on either neural (viz., stimuli-induced electrophysiological activity) or behavioural levels (viz., judging stimuli for their degree of self-relatedness, high or low).

In addition to investigating pre-stimulus (e.g., prior to stimulus presentation or perception) electrophysiological activity, because this activity is mediated by the excitation-inhibition balance (EIB) and that balance has been successfully modelled as due to glutamate-GABA interaction, especially in the lower frequency bands upon which we focus (Logothetis, 2008), we sought to identify this activity's biochemical substrate. Although inhibitory GABA has recently been shown to mediate EEG task-evoked measures like gamma band oscillations (Muthukumaraswamy et al. 2009, 2013; Lally et al. 2014; Barr et al., 2013), studies on excitatory glutamate modulation of EEG measures have been reported less widely (for exceptions, see Lally et al. 2014; for animal studies, see Morales-Villagrán et al., 2008). Glutamate is an excitatory transmitter that fMRI and MRS analyses have shown to mediate resting state activity, including both intra-regional activity levels and trans-regional functional levels (Enzi et al. 2012; Duncan et al., 2011, 2013; also see Falkenberg et al., 2012; Scheidegger, 2012). Such glutamatergic modulation of the resting state suggests that glutamate might mediate the influence of pre-stimulus state activity on stimulus-related activity. But the hypothesis remains untested, both in general and in the case of self-related stimuli.

Accordingly, the general aim of our combined EEG-MRS study was to investigate both electrophysiological and biochemical correlates of the relationship between (i) resting state activity and (ii) stimulus-related activity while subjects perceive and render judgments concerning self-related

stimuli. In short, we investigated the electrophysiological and biochemical mechanisms of the rest-self overlap. More specifically, our first aim concerned whether pre-stimulus (e.g., prior to stimulus presentation or perception) activity levels can be marshalled so as to predict how stimuli are perceived and classified, as either high or low self-related. We hypothesized that pre-stimulus (e.g., prior to stimulus presentation or perception) activity in the lower alpha range (8-9Hz) would precede those stimuli that subjects perceived or judged to be high self-related. To test this hypothesis we conducted an EEG study wherein subjects were shown a standardized set of pictorial stimuli that they were required to assess as high or low self-related.

Our second aim comprised investigating glutamatergic modulation of pre-stimulus (e.g., prior to stimulus presentation or perception) activity changes. Toward that end, using magnetic resonance spectroscopy (MRS), we studied the same subjects who had participated in the EEG study with both EEG and MRS being measured in a random sequence. By so doing we were able to measure glutamate concentrations in several regions: perigenual anterior cingulate cortex (PACC), dorsolateral prefrontal cortex (DLPFC), occipital cortex (OCC), and thalamus (THA). Based upon the rest-self overlap considerations adumbrated above, we hypothesized that glutamate in the PACC—but not in the other regions—would mediate elevated pre-stimulus alpha power at frontal regions for stimuli that subjects assessed as high self-related. Our group has abundant experience in combining MRS with other imaging modalities (Duncan et al., 2011, 2013; Wiebking et al. 2013, 2014; Qin et al., 2013; Hayes et al., 2013; Enzi et al., 2013; Northoff et al., 2007; Walter et al. 2009); this experience enabled us to link pre-stimulus (e.g., prior to stimulus presentation or perception) activity levels (as measured in EEG) to biochemical modulation by resting state glutamate concentration (as measured in MRS).

(Figure 1)

Experimental Procedures

Subjects

Twenty-seven students were recruited from a local university to participate in this experiment and each was given a modest remuneration. Of the 27, data sets for 11 were excluded, both because of EEG or MRS artefact and because of the hypothesis-driven need to identify an equal distribution of high and low self-related responses. Thus, results reported here are for a total of 16 participants. All participants were right-handed, had normal or corrected-to-normal vision, and no history of neurological disorders; they were also 20 years of age, or older, as 20 is the age of the majority in Taiwan. Informed consent was obtained after the experiment was explained and after each participant was carefully screened. IRB approval for this project was granted by the Taipei Medical University-Joint Institutional Review Board (Approval No: 201209022).

Overall design

EEG was employed while participants viewed various emotional and non-emotional pictures. First, these pictures were just perceived, while participants exercised no judgment. Second, participants were required to evaluate or render judgment: to judge whether the objects or events in the pictures are personally relevant (viz., high or low self-related). While their EEG was being recorded, participants were able to choose between high and low by pressing a button available to them.

MRS measurement of Glx, the combined concentration ratio of glutamate/glutamine to creatine, was employed in order to separate Glutamate and Glutamine so as to identify differential correlation findings for both substances (e.g., Duncan et al. 2012). Target regions were the PACC, DLPFC, OCC, and the THA. After concluding both EEG and MRS, participants were asked to perform subjective

ratings of the pictures' distinct dimensions (emotional valence, arousal, meaningfulness, personal relevance, as well as time and space perception).

Visual Stimuli

Participants were presented with 114 emotional and neutral pictures from the International Affective Picture System (IAPS). The selection of the pictures was based upon previous ratings (Libkuman et al., 2007) for the dimensions personal relevance (viz., high and low self), valence and arousal, as well as ratings from Schneider et al. (2008), Northoff et al. (2009), Grimm et al. (2006, 2007, 2009), and Heinzl et al. (2008, 2009). Based upon these ratings, stimuli were balanced with regard to valence, arousal, and self-relatedness. This allowed us to vary single dimensions such as self-relatedness (personal relevance) while leaving the other dimensions (valence and arousal) maximally unaltered. The dimensions of self-relatedness, emotional valence, and arousal were all balanced in their high and low degrees; this means that there were no significant differences in the stimuli themselves concerning all three dimensions. Having balanced the dimensions in this way made it possible to attribute our subjects' assessments of stimuli during EEG to the subjects themselves (viz., their individual or subjective preferences) rather than to the stimuli or objective features of the stimuli.

All pictures were presented twice, in a randomized sequence. By pressing a button they could render a judgment of high or low self-relatedness/personal-relevance. After 10-14 s of fixation presentation (pre-stimulus), pictures were presented to participants for a 3 s period of "pure" perception, a period during which participants were not asked to do anything. This period was followed by a fixation cross with a varied duration of 3-5 s (See appendix, Figure 1). The fixation cross, in turn, was followed by a judgment period, wherein only the question requesting a judgment—but not the picture—was shown. Judgments concerned self-relatedness, as explained above. The judgment period lasted for 3 seconds and was followed by a period of 1-2 s to allow for a blink. The main focus for EEG analysis was on the period prior to the perception of the stimulus which we coin 'pre-stimulus'. In

contrast, the judgment period only served for behavioural purpose to obtain subject's ratings of the preceding picture. Hence, in the following the term 'pre-stimulus' refers to the period that immediately precedes the presentation of the stimulus, the perception period, rather than the judgment period.

EEG Data acquisition and analysis

EEG data were recorded using a Neuroscan amplifier (Compumedics Neuroscan, Charlotte, NC, USA) and Ag/AgCl electrodes through a 64-channel cap (according to the International Ten-Twenty System) referenced to the nose tip. Data were sampled at 1,000 Hz with DC recording. The impedance of each electrode was kept below 5 k Ω and 60 Hz was notched. The electrooculogram (EOG) was recorded by two pairs of electrodes, one pair above and below the right eye and the other on the outer canthus of each eye. A ground electrode was attached to the middle of the forehead.

EEG data analysis was performed using EEGLAB toolbox (Delorme and Makeig, 2004) running under Matlab 7.9.0 (The Mathworks Inc.). Data were filtered using a low-pass filter 120 Hz, and a high-pass filter of 1 Hz for the EEG data. Stimulus-locked data epochs starting from 3000 ms before and 3000 ms after the stimulus onset were extracted.

Artefact rejection

Epochs with irregular noise were identified and rejected using a computer algorithm based on abnormal statistical distribution, as well as by inferences from visual inspection (Delorme et al., 2007). Typical physiological artefacts (e.g., eye blinks, eye movement, and muscle potentials) were retained for the independent component analysis (ICA). The extended infomax ICA were performed to obtain

64 ICs from the stimulus locked epoch in each participant. For each IC, an equivalent current dipole was estimated (DIPFIT 2.2, EEGLAB plug-in using Fieldtrip toolbox functions; Robert Oostenveld). ICs representing the typical physiological artefacts and electrode artefacts were identified by visual inspection of their time course data, multi-trial ERP-image plots, the power spectrum, scalp topography, and dipole. On average, 3 ICs of stimulus-locked epoch were rejected from each participant's data. The remaining ICs were back-projected onto the scalp electrodes to obtain artefact-free EEG data. After completion of these artefact rejection analyses, each type of data epoch was divided into experimental conditions.

ERP analyses

For ERP extraction, data were filtered with a low-pass filter of 15 Hz, and were baseline corrected using data from -100 ms to 0 ms relative to stimulus onset, i.e., the presentation or perception of the stimulus (rather than the judgement). Data for high-self and low-self conditions were averaged across trials.

ERSP analyses

Event-related spectral perturbation (ERSP) is the degree to which spectral power differs from mean baseline power as a function of time and frequency. For ERSP calculation, the Morlet wavelet was used. 99 log-spaced frequencies ranging from 1 Hz to 100 Hz were calculated every 1 ms starting from -3000 ms prior to and ending up to 3000 ms following stimulus onset (baseline from -1900ms to -1700ms). Wavelet cycle was increasing from 2 Hz, at the lowest frequency measured, 0.08 Hz, to 10 Hz at the highest, 100 Hz (Delorme & Makeig, 2004). To compare the difference between the high self and low self-condition, we focused alpha band (8-9 Hz) in ERSP. Since several studies during self- vs

other-judgment self-reference showed the activity at Cz (Geng, Zhang, Li, Tao, & Xu, 2012; Mu & Han, 2010), we here also focused on ERSP at Cz.

We performed cluster-based permutation test (Cohen, 2014) to avoid the issue of multiple comparisons in the large time-frequency space. First, we calculated t -value for each pixel data of time-frequency window (-1000ms – 0 ms, 8-9 Hz) at Cz, and those were thresholded using uncorrected parametric p -value ($p < 0.05$). Then, `bwconncomp` Matlab function was applied to identify clusters in the thresholded map, and the sum of t -value in each cluster was calculated. To generate probability distribution of the sum of t -value under null hypothesis, the 2000 times iterations of the following three steps were conducted: First, the condition label (self vs. non-self) was randomly shuffled. Second, we calculated t -value for each pixel data, and those were thresholded using uncorrected parametric p -value ($p < 0.05$). Third, we collected the biggest sum of the absolute t -value in the cluster. The distribution generated by the iterations was used to calculate the critical value.

MRS data acquisition and analysis

Single-voxel 1H MR spectra were acquired in a separate session (in randomized sequence with EEG) during the resting state using a 3T whole body MRI system (Skyra, Siemens Medical Solutions, Erlangen, Germany) using an 32-channel head coil (PRESS, TR = 2 s; TE = 40 ms; Bandwidth=2000 Hz; sample points=2048; averages=256). Voxels were prescribed on a high-resolution T1-weighted 3D data set (MPRAGE, TR = 2 s; TI = 1.1 s; TE = 4.8 ms; flip angle = 7; FoV = 256 x256 x192 mm; spatial resolution = 1x1 x1 mm). One voxel of 20x10x20 mm was placed on the bilateral PACC, while a second was placed on the left DLPFC cortex (20x20x 20mm) in order to control for the regional specificity of the PACC effects. So as to account for visual effects, we also placed a voxel (20x20x20mm) on the occipital cortex (OCC). And, one additional voxel was placed on the THA (15x15x20mm).

Because of the crucial role played by the PACC during self-related processing in the cortical midline structure, we concentrated our correlation analysis on this region, while other regions were used as controls. For each region, additional non-water-suppression scan without pre-saturation of water signal was acquired using 4 averages for automatic phase correction.

Spectra were analyzed using fully automatic LCModel software version 6.1.0 (www.s-provencher.com/pages/lcmodel.shtml). Metabolite concentrations of N-acetyl aspartate (NAA), total creatine (tCr) including creatine and phosphocreatine, total choline (tCho), myo-inositol (mI), glutamate (Glu), glutamine (Gln), and the combination of glutamate and glutamine (Glx) were obtained. Evaluation of the spectral quality was based on the line width at full-width at half maximum (FWHM) and the SNR as provided by LCModel.

For individual metabolites, a standard deviation of the fitting error, Cramer-Rao lower bound (CRLB), was reported. Expressed in concentration percentage, CRLB can function as an indicator of the reliability of metabolic concentration quantification. The CRLB of each metabolite is commonly used to quantify the goodness-of-fit of the LCModel. Spectra with FWHM line widths larger than 8 Hz, and quantification results with a Cramer-Rao lower bound higher than 20%, were excluded from further analysis. The measurements for two subjects in the PACC and one subject in the LAI were discarded for these reasons (PACC: n= 16, DLPFC, OCC < Thalamus). Metabolite concentrations are given as their ratio to the measured tCr concentration. As a slight interdependence, due to a spectral overlap in their resonances, results for glutamate and glutamine these were quantified together. Recall that this combined concentration ratio of glutamate/glutamine to tCr is herein referred to as Glx. Because the LCModel quantifies glutamate and glutamine separately, individual glutamate and glutamine can be also subjected to further analysis associated with Glx.

Statistical analysis

ERP was averaged over a 1000-ms epoch separately for high-self condition and low-self condition. Baseline was corrected using the mean amplitude during the 200-ms pre-stimuli. A negativity was measured as the mean amplitude between 150 and 300 ms. A positivity was measured as the mean amplitude between 300 and 400 ms. ERP data were then compared using paired sample t-tests. Using Pearson correlation the relation was calculated between pre-stimulus (-600ms to -400 ms) ERSP/alpha power (8-9Hz) difference between high-self and low-self condition - and PACC Glutamate/Creatine. For each participant, differences in pre-stimulus power were calculated by subtracting the low-self condition from the high-self condition. Because we hypothesized that glutamate in the PACC would mediate elevated pre-stimulus alpha power at frontal regions, we have focused on 14 frontal channels (Fp1,Fpz,Fp2,AF3,AF4,F7,F5,F3,F1,Fz,F2,F4,F6,F8) and applied FDR p -value correction for the correlation analyses.

Results

Behavioral data

The mean number of pictures judged as high-self and low-self were, respectively, 21.04 pictures ($SD=7.35$) and 24.96 pictures ($SD=7.20$). A paired t-test revealed that no significant difference was found ($t(26)=-1.41$, $p=0.17$) between high-self and low-self. The mean RT for the high-self condition was 1101.77 ms ($SD=269.68$); for low-self, 1182.69 ms ($SD=314.31$). A paired t-test revealed that RTs for high-self were shorter than for low self ($t(26)=-2.77$, $p=0.01$). In accord with available ratings (Libkuman et al., 2007; Northoff et al., 2009; Schneider et al., 2008), we also compared high and low self-related stimuli as judged by our participants with regard to potential differences in emotional

valence, and arousal; comparisons yielded no significant differences or correlations with subject ratings. That there were no such differences or correlations suggests that differences in assessment between high and low self-related stimuli were dependent upon participants' subjective preferences rather than upon objective features of the stimuli themselves.

Comparison between high and low self-related stimuli in ERP and ERSP (EEG)

First we analysed the ERPs for the perception period. ERP's during this period for pictures that were subsequently rated as high self were compared with those pictures that were rated as low self. A comparison of high- to low- self-relatedness, as can be seen in Figure 2, yielded differential ERPs between 150-400ms—increased negativity between 150-300ms ($t(20)=-2.82$, $p<0.05$ in Fz); comparison also yielded delayed and reduced positivity at 300-400ms—($t(20)=-1.35$, $p<0.05$ in Cz). Since we controlled our stimulus set for valence, arousal, and dominance (see methods section), ERP and subsequent ERSP differences between high and low self-related stimuli must reflect the degree of self-relatedness assigned to the stimuli by subjects (see Figure 2a and b).

(Figure 2)

Based on these results, we calculated event-related spectral perturbation (ERSP) at Cz for the perception periods of high and low self-related stimuli. The ERSPs evidenced a significant difference between high and low-self-related stimuli in the lower alpha frequency range, 8-9 Hz and theta frequency range, 4-7 Hz (see Figure 3a). Most notably, alpha power changes were visible not only during the stimulus-related activity itself, e.g., during presentation/perception of the stimulus but also during the pre-stimulus period (preceding the presentation. e.g., perception of the stimulus) at around -

600 to -400ms prior to the onset of the stimulus presentation (see the box in Figure 3b). Cluster-based permutation test yielded significant differences (Cluster t -value sum=141.72, df =20, cluster count = 57, corrected p <0.05) in lower alpha (8-9hz) between high and low self in ERSP around -500 ms pre-stimulus at Cz (see Figure 3b). This finding was further confirmed by the topographic plot for that time period (see Figure 3c). We also excluded spill-over effects from preceding stimuli: we compared high versus low-self-relatedness judgment as sorted by the preceding judgment. We did not find any significant difference in ERSP alpha power between high self-relatedness (-0.20 dB) and low self-relatedness (0.18 dB) of the preceding judgment at -600 - -400 ms, ($t(20)$ = -0.09, uncorrected p >0.05 at Cz).

(Figure 3)

Collectively, these results demonstrate that the ERSP was significantly stronger in the lower alpha range (8-9Hz) prior to the onset of stimulus presentation, e.g., perception in those stimuli that were evaluated as high self, by comparison to those evaluated as low self. Thus it seems that the degree of pre-stimulus alpha power can predict subjects' perception (and subsequent assessment) of stimuli as high or low self-related. And, perception, in turn, is accompanied by higher or lower levels of post-stimulus alpha power.

Prediction of ERSP changes (EEG) by glutamate (MRS)

First, we obtained reliable spectra for glutamate in all four regions. We then investigated how the levels of glutamate (Glu/Cr as obtained in MRS) in the PACC (as controlled for by DLPFC, OCC,

and THA) were related to the pre-stimulus ERSP changes that predicted high and low self differentiation. In order to understand that relationship we entered the concentration of Glu/Cr in PACC as a regressor when calculating the topographic ERSP high and low self map at -600 to -400ms. This yielded a significant positive relationship between PACC Glu/Cr and low alpha (8-9hz) pre-stimulus (-600 - -400ms) high-low self difference at Cz and, in the frontal regions (see Figure 4: $r=0.72$, uncorrected $p=0.00227$, FDR corrected $p<0.05$): the higher the subject's resting state concentration of glutamate in the PACC, the larger its pre-stimulus (-600 - -400ms) low alpha (8-9 Hz) power/ERSP difference between high and low self (see Figure 4a).

Most importantly, the correlation with glutamate was specific for pre-stimulus (e.g., prior to the presentation or perception of the stimulus) alpha power, since glutamate did not correlate with the post-stimulus onset changes in alpha power. The correlation was also regionally specific, since neither did glutamate in the other regions (DLPFC, THA, and OCC) correlate with pre-stimulus alpha power (see Duncan et al. 2014).

To further illustrate the findings, we selected anterior frontal electrodes (F4, F5, F6, and F7) and correlated their respective pre-stimulus alpha power (-600 to -400ms, -400 to -200ms, -200 to 0ms, 0 to 200ms, and 200 to 400ms) with PACC glutamate concentration. This analysis yielded significant correlations between PACC glutamate and pre-stimulus alpha differences between high and low self for all of the pre-stimulus time windows at F7, F5 and F3 ($r = 0.38$ to 0.59 ; $p= 0.000$ to 0.03): the higher the subject's PACC glutamate concentration, the larger its pre-stimulus alpha ERSP difference between high and low self-related stimuli (see Figure 4b).

(Figure 4)

In sum, resting state concentration of glutamate in the PACC specifically predicts the pre-stimulus state alpha power (8-9 Hz) difference between stimuli that were subsequently perceived (and assessed) as high and low self-related. That is, a higher resting state concentration of glutamate leads to a stronger ERSP pre-stimulus alpha power difference between high and low self-related stimuli. This finding suggests that the level of resting state glutamate modulates pre-stimulus alpha power (prior to presentation or perception of stimuli) in a way relevant to subsequent information processing: viz., the assignment of different degrees of self-relatedness (e.g., high or low) to subsequently perceived stimuli.

Discussion

This is the first combined EEG-MRS investigation of the neural and biochemical correlates of the relationship between resting state and self-related activity—the rest-self overlap. We first focused on whether pre-stimulus activity levels (as accounted for by ERSP in EEG), i.e., preceding the presentation or perception of the stimulus, predicted the subjects' subsequent perception (and judgment) of the stimuli as high and low self-related. Second, we sought to determine whether pre-stimulus activity changes were modulated by glutamate resting state levels in the PACC. Our combined EEG-MRS study yielded three main results: (i) post-onset stimulus-related ERP differences and power (ERSP) differences between high and low self-related stimulus' perception at approximately 150-400ms; (ii) elevated pre-stimulus alpha power (ERSP) preceding the presentation, e.g., perception of those stimuli that were perceived to be highly self-related; and, (iii) glutamate resting state concentration in PACC that correlates with elevated pre-stimulus (i.e., prior to stimulus presentation or perception) alpha power for high self stimuli, in a regionally (not in other regions like the DLPFC, OCC or THA) and biochemically (only glutamate, no other substances) specific way.

ERP and alpha power (ERSP) differences between high and low self-related stimuli

The first main finding in EEG concerns the observation of post-onset stimulus-related differences during stimulus presentation, e.g., perception in both ERP and ERSP differences at around 150-400 ms between high and low self-related stimuli. The ERP showed increased negativity between 150-300ms, which is followed by delayed and reduced positivity between 300-400ms that characterizes high self-related stimuli as distinguished from those that exhibit low self-relatedness. ERSP yielded elevated power specifically in the lower alpha frequency range (8-9 Hz), during high self-related stimuli. Both ERP and ERSP findings are well in accord with previous EEG studies of self-relatedness (see Qin et al., 2008; Esslen et al., 2008; Knyazev, 2013).

Pictures were grouped according to how they were assessed by participants: high- or low self-related. Therefore the ERP and ERSP differences between high and low self-related stimuli must be related specifically to the participants' subjective assessment (rather than to objective features of the stimuli). Moreover, we controlled for other stimulus-related features (e.g., emotional) as well as for task-related effects (e.g., effects of the perception period). Accordingly, it appears to the case that observed post-stimulus ERP and ERSP differences specifically reflect the subjective component evoked during neural stimulus processing. This is a tentative hypothesis, however, one which requires future studies wherein objective and subjective components of stimulus processing are directly compared to one another.

Pre- stimulus alpha power during high self-related stimuli

In addition to the post-onset stimulus-related effects, we also observed pre-stimulus differences prior to the presentation, e.g., perception, between those stimuli assessed as high and those assessed as low self-related. More specifically, stimuli that were subsequently perceived and assessed to be high

self-related showed elevated power, specifically in the alpha range, beginning at -600ms. This finding concerning the impact on subsequent perception is compatible with other EEG studies, especially those that involve visual and auditory modalities or cognitive functions (for a review see, Sadaghiani et al., 2010).

This marks the first time that such pre-stimulus prediction has been found applicable to and extendable to participant attribution of self-relatedness. Boldly, elevated pre-stimulus alpha power prior to stimulus-onset, at approximately -600 ms to 0 ms, can be used to predict that subjects will perceive (and subsequently assess) the respective stimulus as high rather than low self-related. It follows then that subjective perception (and assessment) of stimuli as high or low self-related may depend not only upon the stimuli themselves and their objective features, but also on the subjects' prior state, e.g. pre-stimulus alpha power that precedes the presentation, e.g., perception of the stimulus.

Collectively, our data underscore the special role of power changes, specifically in alpha, during pre-stimulus processing of self-relatedness. Power increases in alpha during stimulus-induced activity when comparing high versus low self-related stimuli have been reported by others (Knyazev 2013; Mu et al., 2008; Justen et al., 2014). Our data, however, extend these findings: this is the first study that shows pre-stimulus alpha power can predict self-relatedness perception (and subsequent assessment). It seems to be the case that elevated, pre-stimulus alpha power disposes subjects' perception (and assessments) towards assigning a high degree of self-relatedness to the stimuli. In contrast, low level of pre-stimulus alpha power lets subjects perceive (and assess) the subsequent stimulus as low rather than high self-related.

But it remains unclear just what mechanisms enable alpha to play this distinctive role. Although it is well known that alpha is sensitive to changes in resting state activity level, as when shift from eyes open to eyes closed (Barry et al., 2007; Barry et al., 2009; Chen et al., 2008; Nakao et al., 2013), the functional implications of this role in the resting state have not yet been fully explored. One

such functional implication, however, may be indicated by our results—it seems to be a perfect candidate mechanism for neural mediation of the close relationship between resting state activity and self-relatedness, the rest-self overlap. Tentatively then, we propose the hypothesis that alpha power in the brain's resting state is a mechanism by means of which the resting state can select what is relevant for or important to any given person.

The hypothesis that resting state alpha is a mechanism for selecting high self-related stimuli accords well with what is known about the functional role of alpha in general. It has indeed been suggested that rhythmic alpha activity, through event-related synchronization and desynchronization, shapes the functional, timing and spatial structure available for incoming stimuli, through mediation of activation processes and selective gating by specific networks (Klimesch, 2007; Jensen, 2010). According to this framework, the fact that our participants subjectively perceive (and assess) standardized stimuli as more or less self-related could result from extrinsic stimuli encountering a distinctive spatiotemporal constellation shaped by pre-stimulus (e.g., prior to stimulus presentation or perception) state alpha power and timing. Future studies utilizing MEG or other technologies might enable more precise investigation of the pre-stimulus alpha's spatiotemporal coordinates, including the mechanisms by means of which those coordinate modulate the degree of self-relatedness attributed to stimuli.

Glutamate modulates pre- stimulus alpha power

In addition to EEG, the subjects also underwent MRS to measure the resting state concentration of glutamate in different regions. We detected that glutamate resting state concentration in the PACC (but not in the DLPFC, VCC, or THA) predicted the degree of pre-stimulus (e.g., prior to stimulus presentation or perception) state alpha power (8-9 Hz) difference between high and low self-related

stimuli: that is, the higher the glutamate resting state concentration in the PACC, the higher the pre-stimulus alpha power, and the more likely that the subsequent stimulus will be perceived and assessed as high (rather than low) self-related. This was observed for the whole brain, as shown by our topographical map correlation (Figure 4a) as well as for single anterior frontal electrodes in the pre-stimulus windows (Figure 4b). Most importantly, this correlation was biochemically, regionally, and neuronally specific: it applied to glutamate but not other metabolites; to the PACC, not other regions; and, to pre-stimulus (e.g., prior to presentation or perception of the stimulus), not post-onset (e.g., during stimulus presentation or perception) stimulus-related ERSP differences.

Glutamate has been demonstrated to modulate resting state measures like the BOLD amplitude in fMRI (Enzi et al. 2013), and the PACC functional connectivity to cortical and subcortical regions (Duncan et al.; 2011, 2013; Falkenberg et al. 2012; Jochum et al., 2012). These findings, however, left unresolved how glutamatergic modulation of resting state activity influences stimulus-induced activity, in general, and self-related stimuli, in particular. Our study is the first to demonstrate that glutamate specifically modulates pre-stimulus state alpha power as measured in EEG. This finding is not only in line with previous EEG investigations of glutamate (Lally et al., 2013), it also suggests that glutamate can influence indirectly, via pre-stimulus state, subsequent stimulus-induced activity: first glutamate mediates pre-stimulus state activity and, then resting state activity modulates stimulus-induced activity. This is unlike the case with GABA, which appears to influence directly stimulus-induced activity (Wiebking et al. 2013, 2014, Hayes et al. 2013). In sum, glutamate's influence on stimulus-related behaviour, as for instance whether a stimulus is judged to be high or low self-related, is indirect, not direct.

As one of the main players, along with GABA, of the excitation-inhibition balance (EIB) (Logothetis 2008), glutamate mediates predominantly the neural excitatory level. Our finding that glutamate mediates the level of pre-stimulus alpha power suggests that pre-stimulus alpha power is

related to the excitatory level and, ultimately, the shift of the EIB towards elevated excitation. This finding, in conjunction with our observation that increased alpha power mediates high self-related stimuli, suggests that high levels of glutamatergic mediated resting state excitation in the PACC incline or dispose subjects to assign high degrees of self-relatedness to subsequent stimuli. This is just a working hypothesis, however, one that requires more detailed investigation of intracranial data measuring firing rates and local field potentials, along with measurement and stimulation of glutamate.

Methodological limitations

We are not able to specify the neural substrate of early pre-stimulus differentiation. Although we are able to exclude spill-over effects from preceding judgments, the exact neural mechanisms of pre-stimulus alpha increase remain unclear. These mechanisms may be related to purely spontaneous activity changes in the resting state and also in lower frequency bands like infra-slow frequency bands, because spontaneous activity of this type predominates, especially in cortical midline structures like the PACC (see Raichle et al. 2001; Lloyd 2011; Northoff 2014a and b). In this context it is intriguing to note that cortical midline structures have been associated with both self (Northoff et al., 2006; van Meer et al., 2010) and the rest-self overlap (Qin and Northoff 2011; Schneider et al., 2008; d'Argembeau et al., 2005; Whitfield-Gabrieli et al., 2001). But in order to better understand how self and lower frequency bands might be related, future studies that combine the recording of slow cortical potentials in EEG and fMRI during self-relatedness will be necessary. Only by conducting such studies can we further specify the neural mechanisms of elevated pre-stimulus alpha and of processing high self-related stimuli. And, as mentioned above, the use of higher resolution methods such as intracranial recordings will likely be necessary in order to identify the mechanisms whereby the interplay between glutamate concentration and pre-stimulus alpha power is realized.

Another limitation of this study is that we only measured the resting state concentration of glutamate. We did not conduct functional MRS while we did not conduct functional MRS (cf., Lally et al., 2013). A functional MRS study would have enabled us to measure glutamate levels while high and low self-related stimuli were being processed. Functional MRS would have enabled us to match glutamate and alpha power changes temporally.

Finally, we did carefully select our stimuli so as to exclude differences between high and low self-related stimuli, with respect to dominance, valence, and arousal. Nevertheless, we remained unable to control for other possible factors like reward. High self-related stimuli may exhibit a higher degree of reward than those that exhibit low self-relatedness, because self appears to be related to distinctive activity levels in the reward system (deGreeck et al., 2008, 2009, and 2010). Hence, future EEG and MRS studies may be necessary in order to directly compare self-relatedness and reward, as well as to determine how these are modulated by glutamate.

Conclusion

This is the first direct investigation of the rest-self overlap in that we investigated the neural and biochemical mechanisms of the relationship between resting state and self-related processing. We demonstrated that the degree of pre-stimulus (e.g., prior to stimulus presentation or perception) changes in alpha power (ERSP) can be used to predict whether subsequent stimuli will be perceived (and assessed) as high or low self-related. Moreover, though conclusions are necessarily tentative, these pre-stimulus (e.g., prior to stimulus presentation or perception) state changes appear to be related to glutamate resting state concentration in the PACC. Taken together these findings are the first to reveal temporal-neuronal (e.g. alpha power) and biochemical (e.g., glutamate) mechanisms that might mediate the rest-self overlap. Results of this investigation enhance understanding the neuro-biochemical

mechanisms involved in the generation of self-relatedness. But the investigation's importance extends beyond just the issue of self-relatedness. These data also reveal some of the mechanisms—viz., alpha power and glutamatergic-mediated neural excitation—by means of which the pre-stimulus resting state (e.g., prior to stimulus presentation or perception) can influence subsequent stimulus processing. Of most importance, our study suggests that these mechanisms have significant psychological consequences, as is the case with the assignment of high or low self-relatedness to stimuli. In a word, these mechanisms partially explain what matters to self, and why.

Acknowledgments

GN is grateful for financial support from CIHR, EJLB-CIHR, and ISAN/HDRF. This work was also supported by JSPS KAKENHI Grant Numbers 24390284, 25870467, and 26285168 to TN. TL and NY are grateful for financial support from Taiwan's National Science Council (Grant numbers: 100-2410-H-004-139-MY3 and 102-2420-H-038-002-MY2.). JX is grateful for financial support provided by National Chengchi University's Research Center for Mind, Brain, and Learning.

References

- Ai, L., Ro, T. (2014). The phase of prestimulus alpha oscillations affects tactile perception. *Journal of Neurophysiology*, 111(6), 1300-1307
- Anderson AK, Christoff K, Stappen I, Panitz D, Ghahremani DG, Glover G, Gabrieli JD, Sobel N (2003) Dissociated neural representations of intensity and valence in human olfaction. *Nat Neurosci* 6(2): 196-202.
- Barbas H (2000) Complementary roles of prefrontal cortical regions in cognition, memory, and emotion in primates. *Adv Neurol*.84:87-110.
- Barbas H (2004) Dead tissue, living ideas: facts and theory from neuroanatomy, *Cortex* 40(1): 205–206.
- Barry, R.J., Clarke, A.R., Johnstone, S.J., and Brown, C.R. (2009). EEG differences in children between eyes-closed and eyes-open resting conditions. *Clinical Neurophysiology* 120, 1806-1811.
- Barry, R.J., Clarke, A.R., Johnstone, S.J., Magee, C.A., and Rushby, J.A. (2007). EEG differences between eyes-closed and eyes-open resting conditions. *Clinical Neurophysiology* 118, 2765-2773.
- Bechara A (2004) The role of emotion in decision-making: evidence from neurological patients with orbitofrontal damage. *Brain Cogn.* 55(1):30-40
- Bechara A, Naqvi N (2004) Listening to your heart: interoceptive awareness as a gateway to feeling. *Nat Neurosci.* 7(2):102-3.
- Berridge KC (2003) Pleasures of the brain. *Brain Cogn.* 52(1):106-28.

Block N (1996) How can we find the neural correlate of consciousness? *Trends in Neurosciences* 19 (11): 456-459.

Brown RA, Seligman R. (2009) Anthropology and cultural neuroscience: creating productive intersections in parallel fields. *Prog Brain Res.* 178:31-42. Review.

Buckner, R. L., Andrews-Hanna, J. R., and Schacter, D. L. (2008). The brain's default network: Anatomy, function, and relevance to disease. *Annals of the New York Academy of Sciences* 1124, 1-38.

Campbell BC, Garcia JR. (2009) Neuroanthropology: evolution and emotional embodiment. *Front Evol Neurosci.* 1:4.

Carmicheal , Price CJ (1996) Connectional networks within the orbital and medial prefrontal cortex of macaque monkeys. *Journal of Comparative Neurology.* 371. 179-207

Chen, A.C.N., Feng, W., Zhao, H., Yin, Y., and Wang, P. (2008). EEG default mode network in the human brain: Spectral regional field powers. *NeuroImage* 41, 561-574.

Cohen, M. X. (2014). *Analyzing Neural Time Series Data.* Cambridge, MA: MIT Press.

Chiao, J. Y., Harada, T., Komeda, H., Li, Z., Mano, Y., Saito, D. N., Parrish, T. B., Sadato, N., & Iidaka, T. (2009a). Neural basis of individualistic and collectivistic views of self. *Human Brain Mapping*, 30, 2813–2820.

Chiao, J. Y., Harada, T., Komeda, H., Li, Z., Mano, Y., Saito, D. N., Parrish, T. B., Sadato, N., & Iidaka, T. (2009b). Dynamic cultural influences on neural representations of the self. *Journal of Cognitive Neuroscience.* 22, 1-11.

Choudhury, S. Kirmayer, L. J. (2009). Cultural neuroscience and psychopathology: prospects for cultural psychiatry. *178:263-283*

- Craig AD (2002) How do you feel? Interoception: the sense of the physiological condition of the body. *Nat Rev* 3: 656-666
- Craig AD (2003) Interoception: the sense of the physiological condition of the body. *Curr Opinion Neurobiol* 13:500-505
- Craig AD (2004) Human feelings: why are some more aware than others? *Trends Cog Sci* 8(6):239-241
- Craig AD (2005) Forebrain emotional asymmetry: a neuroanatomical basis? *Trends Cogn Sci*. 9(12):566-571
- Critchley HD (2005) neural Mechanisms of Autonomic, affective, and Cognitive Intergration. *J Comp Neur*,493: 154-166
- Critchley HD, Wiems SW, Rotshtein P, Oehman A, Dolan RJ (2004) Neural systems supporting interoceptive awareness. *Nature Neuroscience*, 7(2): 189-195
- Damasio A (2003a) Mental self: The person within. *Nature*.15;423(6937):227
- Damasio A (2003b) Feelings of emotion and the self. *Ann N Y Acad Sci*. 1001:253-61
- Damasio AR (1999) *The Feeling of What Happens: Body and Emotion in the Making of Consciousness*. New York, Harcourt Brace
- Damasio AR, Grabowski TJ, Bechara A, Damasio H, Ponto LL, Parvizi J, Hichwa RD (2000) Subcortical and cortical brain activity during the feeling of self-generated emotions. *Nat Neurosci*. 3(10):1049-56
- D'Argembeau A, Collette F, Van der Linden M, Laureys S, Del Fiore G, Degueldre C, Luxen A (2005) Self-referential reflective activity and its relationship with rest: a PET study. *NeuroImage* 25 (2):

- Davidson RJ (2000) Emotion, plasticity, context, and regulation: Perspectives from affective neuroscience. *Psychological Bulletin* 126: 890-909
- Davidson RJ (2004) What does the prefrontal cortex "do" in affect: perspectives on frontal EEG asymmetry research. *Biol Psychol* 67(1-2): 219-234
- de Greck, M. Zhenhao Shi, Gang Wang, iangyu Zuo, Xuedong Yang, Xiaoying Wang, G Northoff, Shihui Han (2011) Culture modulates brain activity during empathy with anger. *Neuroimage*, in press
- de Greck, M., Wang, G., Yang, X., Wang, X., Northoff, G., Han, S. (2011) Neural substrates underlying intentional empathy. *Social Cognitive and Affective Neuroscience*, in press.
- Delorme, A., and Makeig, S. (2004). EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *J Neurosci Methods* 134, 9-21.
- Delorme, A., Sejnowski, T., and Makeig, S. (2007). Enhanced detection of artifacts in EEG data using higher-order statistics and independent component analysis. *Neuroimage* 34, 1443-1449.
- Domínguez D JF, Lewis ED, Turner R, Egan GF. (2009) The brain in culture and culture in the brain: a review of core issues in neuroanthropology. *Prog Brain Res.* 178:43-64.
- Domínguez Duque JF, Turner R, Lewis ED, Egan G. (2010) Neuroanthropology: a humanistic science for the study of the culture-brain nexus. *Soc Cogn Affect Neurosci.* 5(2-3):138-47.
- Edelman GM, Tononi G (2000) *A universe of consciousness: how matter becomes imagination*, Basic Book.
- Ellis RD (2005) *Curious emotions. Roots of consciousness and personality in motivated action.* John Benjamin Publisher, New York/Amsterdam.

Enzi B, Duncan NWN, Kaufmann J, Tempelmann C, Wiebking C., Northoff, G. (2012). Glutamate Modulates Resting State Activity in the Perigenual Anterior Cingulate Cortex - A combined fMRI-MRS study. *Neuroscience* 227: 102–109.

Esslen, M., Metzler, S., Pascual-Marqui, R., Jancke, L. (2008) Pre-reflective and reflective self-reference: A spatiotemporal EEG analysis. *Neuroimage*, 42, 437-449

Falkenberg LE, Westerhausen R, Specht K, Hugdahl K. (2012). Resting-state glutamate level in the anterior cingulate predicts blood-oxygen level-dependent response to cognitive control. *Proc Natl Acad Sci U S A*, 109(13), 5069-73

Gallagher S, Marcel AJ. (1999) The self in contextualized action. *Journal of Consciousness Studies*

Geng, H., Zhang, S., Li, Q., Tao, R., & Xu, S. (2012). Dissociations of subliminal and supraliminal self-face from other-face processing: behavioral and ERP evidence. *Neuropsychologia*, 50(12), 2933–42.

Gillihan, S.J., Farah, M.J. (2005) Is self special? A critical review of evidence from experimental psychology and cognitive neuroscience. *Psychol Bull* 131, 76-97.

Goldie (2002) Emotions, feelings and intentionality. *Phenomenology and the Cognitive Sciences* 1, 235-254.

Grimm S, Ernst J, Boesiger P, Schuepbach D, Boeker H, Northoff G.(2011) Reduced negative BOLD responses in the default-mode network and increased self-focus in depression. *World J Biol Psychiatry*. In Press.

Grimm S, Schmidt CF, Bermpohl F, Heinzl A, Dahlem Y, Wyss M, Hell D, Boesiger P, Boeker H, Northoff G (2006) Segregated neural representation of distinct emotion dimensions in the

prefrontal cortex-an fMRI study. *Neuroimage*, 30(1):325-40.

Grimm, S., Boeker, H., Beck, J., Bermpohl, F., Heinzl, A., Boesiger, P., Northoff, G. (2009) Abnormal functional deactivation in the default-mode network in depression. *Neuropsychopharmacology* 34: 932-943.

Han, S., Gu, X., Mao, L., Ge, J., Wang, G., Ma, Y. (2010). Neural substrates of self-referential processing in Chinese Buddhists. *Social Cognitive and Affective Neuroscience*, 5, 332-339.

Han, S., Mao, L., Gu, X., Zhu, Y., Ge, J., Ma, Y. (2008). Neural consequences of religious belief on self-referential processing. *Social Neuroscience*, 3, 1-15.

Han, S., Mao, L., Qin, J., Friederici, A. D., Ge, J. (2011). Functional roles and cultural modulations of the medial prefrontal and parietal activity associated with causal attribution. *Neuropsychologia*, 49, 83-91.

Han, S., Northoff, G. (2008). Culture-sensitive neural substrates of human cognition: A transcultural neuroimaging approach. *Nature Review Neuroscience*, 9, 646-654.

Han, S., Northoff, G. (2009). Understanding the self: a cultural neuroscience approach. *Progress in Brain Research*, 178, 203-212.

Hsieh PJ, Colas JT, Kanwisher NG. (2012) Pre-stimulus pattern of activity in the fusiform face area predicts face percepts during binocular rivalry. *Neuropsychologia*, 50(4), 522-9

James W (1884) What is an emotion? *Mind* 9:185-205

Jensen, O., Mazaheri, A. (2010). Shaping Functional Architecture by Oscillatory Alpha Activity: Gating by Inhibition. *Frontiers in Human Neuroscience*, 4, 186

Justen, C., Herbert, C., Werner, K., Raab, M. (2014) Self vs. other: Neural correlates underlying agent

identification based on unimodal auditory information as revealed by electrotopography (sLORETA). *Neuroimage*, 259, 25-34

Kirmayer, L. J. (2001). Cultural variations in the clinical presentation of depression and anxiety: implications for diagnosis and treatment. *J Clin Psychiatry*. 62:22-28

Kirmayer, L. J. Groleau, D. (2001). Affective disorders in cultural context *Psychiatr Clin North Am*. 24(3):465-478.

Kitayama S, Park J. (2010) Cultural neuroscience of the self: understanding the social grounding of the brain. *Soc Cogn Affect Neurosci*. 5(2-3):111-29.

Kitayama S, Uskul AK. (2011) Culture, mind, and the brain: current evidence and future directions. *Annu Rev Psychol*. 62:419-49.

Klimesch, W., Sauseng, P., Hanslmayr, S. (2007) EEG alpha oscillations: The inhibition–timing hypothesis. *Brain Research Reviews*, 53, 63-88

Knyazev, G. (2013). EEG Correlates of Self-Referential Processing. *Front Hum Neurosci*, 7, 264.

Kringelbach ML (2005) The human orbitofrontal cortex: linking reward to hedonic experience. *Nat Rev Neurosci*. 6(9):691-702.

Lally, N., Mullins, P., Roberts, M., Price, D., Gruber, T., Haenschel, C. (2014) Glutamatergic correlates of gamma-band oscillatory activity during cognition: A concurrent ER-MRS and EEG study. *NeuroImage* 85 (2):823-833

Lambie JA, Marcel AJ (2002) Consciousness and the varieties of emotion experience: a theoretical framework. *Psychol Rev* 109(2): 219-259.

Lane RD, Reiman EM, Axelrod B, Yun LS, Holmes A, Schwartz GE (1998) Neural correlates of

levels of emotional awareness. Evidence of an interaction between emotion and attention in the anterior cingulate cortex. *J Cogn Neurosci*. 10(4):525-35.

Lang et al. (1993). Looking at pictures: Affective, facial, visceral, and behavioral reactions. *Psychophysiology*, 30, 261-273.

LeDoux JE (1996) *The emotional brain: the mysterious underpinnings of emotional life*. New York: Simon and Schuster.

LeDoux JE (2002) *Synaptic self : how our brains become who we are*, Viking.

Legrand, D., Ruby, P. (2009) What is self-specific? Theoretical investigation and critical review of neuroimaging results. *Psychol Rev* 116, 252-282.

Lemogne, C. Delaveau, P., Freton, M., Guionnet, S., Fossati, P. Medial prefrontal cortex and the self in major depression. *J Affect Disord*. In press.

Lemogne, C., Ie Bastard, G., Mayberg, H., Volle, E., Bergouignan, L, Leherich, S., Allilaire, J.F., Fossati, P., (2009). In search of the depressive self: extended medial prefrontal network during self-referential processing in major depression. *Soc Cogn Affect Neurosci*. 4(3):305-312.

Lemogne, C., Mayberg, H., Bergouignan, L, Volle, E., Delaveau, P., Leherich, S., Allilaire, J.F., Fossati, P., (2010) Self-referential processing and the prefrontal cortex over the course of depression: a pilot study. 24(1-2): 196-201.

Libkuman, T., Otani, H. Kern, R., Viger, S., Novak, N. (2007). Multidimensional normative ratings for the International Affective Picture System. *Behavioral Research Methods*, 39 (2): 326-334.

Linkenkaer-Hansen K, Nikulin VV, Palva S, Ilmoniemi RJ, Palva JM.(2004) Prestimulus oscillations enhance psychophysical performance in humans. *Journal of Neuroscience*, 24(45), 10186-90

Logothetis NK, Murayama Y, Augath M, Steffen T, Werner J, Oeltermann A. (2009) How not to study spontaneous activity. *Neuroimage*. 45(4):1080-9.

Ma, Y., Han, S. (2010). Why respond faster to the self than others? An implicit positive association theory of self advantage during implicit face recognition. *Journal of Experimental Psychology: Human Perception and Performance*, 36, 619-633.

Markus, H. R. & Kitayama, S. (1991), Culture and the self: implications for cognition, emotion and motivation. *Psychol. Rev.* 98, 224–253.

Markus, H. R. & Kitayama, S. (2003). Culture, self, and the reality of the social. *Psychol. Inq.* 14, 277–283.

Mayberg, H.S., 2003. Modulating dysfunctional limbic-cortical circuits in depression: towards development of brain-based algorithms for diagnosis and optimised treatment. *Br. Med. Bull.* 65, 193–207.

Min, B., Herrman, C. (2007) Prestimulus EEG alpha activity reflects prestimulus top-down processing. *Neuroscience Letters*, 422, 131-135

Mu, Y., Fan, Y., Mao, L., Han, S (2008) Event-related theta and alpha oscillations mediate empathy for pain. *Brain Research*, 1234, 128-136

Mu, Y., & Han, S. (2010). Neural oscillations involved in self-referential processing. *NeuroImage*, 53, 757–768.

Muthukumaraswamy, S., Singh, K., Swettenham, J., Jones, D. (2009) Visual gamma oscillations and evoked responses: Variability, repeatability and structural MRI correlates, *Neuroimage*, 49(4), 3349-57

Morales-Villagrán A, Medina-Ceja L, López-Pérez SJ. (2008) Simultaneous glutamate and EEG activity measurements during seizures in rat hippocampal region with the use of an electrochemical biosensor. *Journal of Neuroscience Methods*, 168(1), 48-53

Nagai Y, Critchley HD et al. (2004) Activity in ventromedial prefrontal cortex covaries with sympathetic skin conductance level: a physiological account of a "default mode" of brain function. *Neuroimage* 22(1): 243-51.

Nakao, T., Bai, Y., Nashiwa, H., and Northoff, G. (2013). Resting-state EEG power predicts conflict-related brain activity in internally guided but not in externally guided decision-making. *NeuroImage* 66, 9-21.

Ng, S. H., Han, S., Mao, L., & Lai, J. C. L. (2010). Dynamic bicultural brains: A fMRI study of their flexible neural representation of self and significant others in response to culture priming. *Asian Journal of Social Psychology*, 13, 83-91.

Niedenthal PM (2007) Embodying emotion. *Science*. 316(5827):1002-5.

Northoff G (2004) Why do we need a philosophy of the brain? *Trends Cogn Sci*. 8(11):484-5.

Northoff G (2014a) *Unlocking the brain. Volume I: Coding*. Oxford University Press, New York, Oxford.

Northoff G (2014b) *Unlocking the brain. Volume II: Consciousness*. Oxford University Press, New York, Oxford.

Northoff G and Bermpohl F (2004) Cortical midline structures and the self. *Trends Cogn Sci* 8(3): 102-107.

Northoff G, Heinzl A, de Greck M, Bermpohl F, Dobrowolny H, Panksepp J (2006) Self-referential

processing in our brain--a meta-analysis of imaging studies on the self. *Neuroimage* 15;31(1):440-57

Northoff G, Qin P, Feinberg TE. (2010) Brain imaging of the self--conceptual, anatomical and methodological issues. *Conscious Cogn.* 20(1):52-63.

Northoff G, Qin P, Nakao T. (2010) Rest-stimulus interaction in the brain: a review. *Trends Neurosci.* 33(6):277-84.

Northoff G, Wiebking C, Feinberg T, Panksepp J. (2011) The 'resting-state hypothesis' of major depressive disorder—A translational subcortical–cortical framework for a system disorder. *Neuroscience and Biobehavioral Reviews.* In Press.

Northoff, G. (2007) Psychopathology and pathophysiology of the self in depression neuropsychiatric hypothesis. *J. Affect. Disord.* 104, 1–14.

Northoff, G., Bermpohl, F., (2004). Cortical midline structures and processing of the self. *Trends in Cognitive Science.* 8(3):102-7.

Northoff, G., DeGreck, M., Bermpohl, F., Panksepp, J. (2006) Self-referential processing in our brain – a metaanalysis of imaging studies on the self. *Neuroimage.* 31:440-57.

Northoff, G. Schneider, F. Rotte, M., Matthias, C., Tempelmann, C., Wiebking, C., Panksepp, J. (2009). Differential parametric modulation of self-relatedness and emotions in different brain regions. *Human Brain Mapping,* 30(2), 369-382.

Panksepp J (1998) *Affective neuroscience: The foundations of human and animal emotions,* Oxford University Press.

Panksepp J (2005) *Affective consciousness: Core emotional feelings in animals and humans.*

Conscious Cogn 14(1): 30-80.

Park H, Rugg MD. (2010) Prestimulus hippocampal activity predicts later recollection. *Hippocampus*, 20(1), 24-8

Phan KL, Wager T, Taylor SF, Liberzon I (2002) Functional neuroanatomy of emotion: a meta-analysis of emotion activation studies in PET and fMRI. *Neuroimage* 16(2): 331-348.

Phillips ML, Williams LM, Heining M, Herba CM, Russell T, Andrew C, Bullmore ET, Brammer MJ, Williams SC, Morgan M, Young AW, Gray JA (2004) Differential neural responses to overt and covert presentations of facial expressions of fear and disgust. *Neuroimage*. 21(4):1484-96.

Pollatos O, Herbert BM, Matthias E, Schandry R (2007) Heart rate response after emotional picture presentation is modulated by interoceptive awareness. *Int J Psychophysiol*, 63(1):117-124

Pollatos O, Kirsch W, Schandry R (2005) On the relationship between interoceptive awareness, emotional experience and brain processes. *Brain Res Cogn Brain Res* 25(3) : 948-962.

Pollatos O, Traut-Mattausch E, Schroeder H, Schandry R (2006) Interoceptive awareness mediates the relationship between anxiety and the intensity of unpleasant feelings. *J Anxiety Disord*. 30.

Qin P, Liu Y, Shi J, Wang Y, Duncan N, Gong Q, Weng X, Northoff G. (2011) Dissociation between anterior and posterior cortical regions during self-specificity and familiarity: A combined fMRI-meta-analytic study. *Human Brain Mapping*, 33(1), 154-164.

Qin P, Northoff G. (2011) How is our self related to midline regions and the default-mode network? *Neuroimage*. 57(3):1221-33.

Raichle, M. E. et al. (2001) Inaugural article : a default mode of brain function. *Proceedings of the National Academy of Sciences* 98 (2): 678-682.

Rilling JK. (2008) Neuroscientific approaches and applications within anthropology. *Am J Phys Anthropol.* 47:2-32.

Sadaghiani S, Hesselmann G, Friston KJ, Kleinschmidt A (2010) The relation of ongoing brain activity, evoked neural responses, and cognition. *Frontiers in Systems Neuroscience*, 4:20

Salmon E (2005) Self-referential reflective activity and its relationship with rest: a PET study. *Neuroimage.* 2005 Apr 1;25(2):616-24.

Scheidegger M, Walter M, Lehmann M, Metzger C, Grimm S, et al. (2012) Ketamine Decreases Resting State Functional Network Connectivity in Healthy Subjects: Implications for Antidepressant Drug Action. *PLoS ONE* 7(9): e44799

Schneider F, Bermpohl F, Heinzl A, Rotte M, Walter M, Tempelmann C, Wiebking C, Dobrowolny H, Heinze HJ, Northoff G. (2008) The resting brain and our self: self-relatedness modulates resting state neural activity in cortical midline structures. *Neuroscience.* 11;157(1):120-31.

Seligman R, Brown RA. (2010) Theory and method at the intersection of anthropology and cultural neuroscience. *Soc Cogn Affect Neurosci.* 5(2-3):130-7.

Shibata K, Yamagishi N, Goda N, Yoshioka T, Yamashita O, Sato MA, Kawato M. (2008) The effects of feature attention on prestimulus cortical activity in the human visual system. *Cereb. Cortex*, 18(7), 1664-75

Sui, J., Liu, C. H., Han, S. (2009). Cultural difference in neural mechanisms of self-recognition. *Social Neuroscience*, 4, 402-411.

Thase, E. (2005) Bipolar depression: issues in diagnosis and treatment. *Harv. Rev. Psychiatry* 13, 257–271.

- Thompson E, Varela FJ (2001) Radical embodiment: Neural dynamics and consciousness. *Trends Cogn Sci* 5: 418–425
- van Dijk, H., Schoffelen, J., Oostenveld, R., Jensen, O. (2008) Prestimulus Oscillatory Activity in the Alpha Band Predicts Visual Discrimination Ability. *The Journal of Neuroscience*, 28(8), 1816-1823
- van Gelder TJ, Port RF (1995) *Mind in Motion: Exploration in the Dynamics of Cognition*. MIT Press, Cambridge/MASS.
- Vogeley K, Roepstorff A. (2009) Contextualising culture and social cognition. *Trends Cogn Sci*. 13(12):511-6.
- Wang, G., Mao, L., Ma, Y., Yang, X., Cao, J., Liu, X., Wang, J., Wang, X., Han, S. (2012) Neural representations of close others in collectivistic brains. *Social Cognitive and Affective Neuroscience*, in press.
- Weisz, N., Wühleb, A., Monittolaa, G., Demarchia, G., Freya, J., Popovd, T., Braun, C. (2014) Prestimulus oscillatory power and connectivity patterns predispose conscious somatosensory perception. *Proceedings of the National Academy of Sciences*, 111(4)
- Whitfield-Gabrieli S, Moran JM, Nieto-Castañón A, Triantafyllou C, Saxe R, Gabrieli JD. (2011) Associations and dissociations between default and self-reference networks in the human brain. *Neuroimage*. 1;55(1):225-32.
- Wicker B, Keysers C, Plailly J, Royet JP, Gallese V, Rizzolatti G (2003) Both of us disgusted in My insula: the common neural basis of seeing and feeling disgust. *Neuron* 40(3): 655-664.
- Williams LM, Liddell BJ, Rathjen J, Brown KJ, Gray J, Phillips M, Young A, Gordon E (2004) Mapping the time course of nonconscious and conscious perception of fear: an integration of

central and peripheral measures. *Hum Brain Mapp.* 21(2):64-74.

Yang Z, Zhao J, Jiang Y, Li C, Wang J, Weng X, Northoff G. (2011) Altered negative unconscious processing in major depressive disorder: an exploratory neuropsychological study. *PLoS One.* 6(7):e21881.

Yen, S., Robins, C.J., & Lin, N. (2000). A cross-cultural comparison of depressive symptom manifestation: China and the United States. *Journal of Consulting and Clinical Psychology*, 68, 993-999.

Zahavi D (2005) *Subjectivity and Selfhood. Investigating the first-person perspective.* Cambridge/MA, MIT Press.

Zhu, Y., Zhang, L., Fan, J., & Han, S. (2007). Neural basis of cultural influence on self representation. *Neuroimage*, 34, 1310-1317.

Figure Captions

Figure 1 Experimental design and paradigm

The figure illustrates the three stages of our experimental design (a) and paradigm (b).

(a) Our experimental design comprised three stages: the first stage consisted in measuring EEG during perception of stimuli and their subsequent assessment as high or low self-related. The second stage complemented the first by measuring resting state glutamate in MRS. For the third stage, resting state glutamate was correlated with pre-stimulus EEG. We need to detail the term ‘pre-stimulus ERSP’. The term ‘pre-stimulus’ refers to the period that immediately precedes the presentation, e.g., pure perception of the stimulus (rather than the period preceding the judgement). The term ERSP refers usually to stimulus-related changes; however, in order to describe the pre-stimulus power changes related to and that precede (and distinguish) high and low self-related stimuli, we use the term ERSP. The ERSP does then still refer to changes related to stimuli but those that precede (rather than occur during or follow) the presentation of the actual stimuli by distinguishing between low and high self-related stimuli.

(b) Experimental paradigm: After a 10-14s fixation cross (pre-stimulus duration), subjects were presented with well matched and selected (see methods) stimuli from the International Affective Picture System (IAPS) for three seconds. This was followed by a fixation cross and subsequent assessment of the stimuli as high or low self-related. Assessment of a given stimulus was followed by 1-2 s blink period.

Abbreviations

ERSP = Event-related spectral perturbation

MRS = Magneto-resonance spectroscopy

PACC = Perigenual anterior cingulate cortex

IAPS = International Affective Picture System

RT = Reaction time




Figure 2 Event-related potentials (ERP) during high and low self-related stimuli

The figure illustrates the ERP for high and low-related stimuli at Fz and Cz. The curves and analysis reveal marked difference between high and low self-related stimuli, during the 150ms to 400ms period.




Figure 3 Event-related spectral perturbations (ERSP) during high and low self-related stimuli

The figure illustrates the ERSP for high and low-related stimuli at Cz for stimulus onset (a) and the pre-stimulus period (e.g., prior to stimulus presentation or perception) (b) as well as the topographic maps of ERSP/Alpha power for -600- -400ms prior to the presentation of those stimuli that were subsequently perceived (and assessed) as high and low self-related stimuli (c). Both exhibit low and high self-related stimulus conditions. Note that the post-stimulus onset (e.g., during to stimulus presentation or perception) increase in ERSP in the alpha range (8-9hz) and theta range (4-7hz) at around 0-200ms during high-self related stimuli was larger than that during low self-related stimuli (3a). This post-stimulus data is complemented by a pre-stimulus (e.g., prior to stimulus presentation or perception) increase in alpha during high self-related stimuli, at about -600 to -400ms (3b). .




Figure 4 Glutamatergic modulation (MRS) of pre-stimulus event-related spectral perturbations (ERSP) during high and low self-related stimuli

The figure illustrates how the pre-stimulus (e.g., prior to stimulus presentation or perception) ERSP are related to glutamate concentration (set against creatine/Cr) in the PACC, in a topographic map (a) and a single channel (b). By plotting the concentration of glutamate in PACC (a, on the left) against ERSP of all channels at about -600 to -400ms, we obtained the following topographic map (a, on the right). The deeper the red colour, the more the PACC glutamate concentration influenced the pre-stimulus (e.g., prior to stimulus presentation or perception) ERSP at around -600 to -400ms; this influence was observed especially in frontal and central regions, thereby corresponding to the anatomical location of the PACC. Figure 4b shows the correlation of the PACC glutamate concentration with the degree of pre-stimulus (-600 to -400ms) (e.g., prior to stimulus presentation or perception) ERSP in a single channel, F7: the higher the subject's PACC concentration of glutamate, the higher its pre-stimulus (e.g., prior to stimulus presentation or perception) ERSP at -600 to -400ms, and the more likely the subsequent stimulus will be assessed by the subject as high self-related. Each dot in the graph corresponds to one subject.

