


**POSTILLA**  
**PEABODY MUSEUM**  
**YALE UNIVERSITY**

**NUMBER 157**

**19 JULY 1972**

**NEW EVIDENCE ON THE EVOLUTION OF  
THE PAIRED FINS OF RHIPIDISTIA AND THE  
ORIGIN OF THE TETRAPOD LIMB, WITH  
DESCRIPTION OF A NEW GENUS OF  
OSTEOLEPIDAE**

**KEITH STEWART THOMSON**



SCOTT BRANDE  
DIV. OF GEOL. AND PLANETARY SCIENCE  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
PASADENA, CALIFORNIA 91109



## POSTILLA

Published by the Peabody Museum of Natural History, Yale University

*Postilla* includes results of original research on systematic, evolutionary, morphological, and ecological biology, including paleontology. Syntheses and other theoretical papers based on research are also welcomed. *Postilla* is intended primarily for papers by the staff of the Peabody Museum or on research using material in this Museum.

Editors: Zelda Edelson, Elizabeth G. Weinman, Elise K. Kenney.

*Postilla* is published at frequent but irregular intervals. Manuscripts, orders for publications, and all correspondence concerning publications should be directed to:

Publications Office  
Peabody Museum of Natural History  
New Haven, Conn., 06520, U.S.A.

Lists of the publications of the Museum are available from the above office. These include *Postilla*, *Bulletin*, *Discovery*, and special publications. *Postilla* and the *Bulletin* are available in exchange for relevant publications of other scientific institutions anywhere in the world.

Inquiries regarding back numbers of the discontinued journal, *Bulletin of the Bingham Oceanographic Collection*, should be directed to:

Walter J. Johnson, Inc.  
111 Fifth Avenue  
New York, N.Y. 10003.

NEW EVIDENCE  
RHIPIDISTIA  
DESCRIPTION

KEITH STEWART

Department of  
Yale University

ABSTRACT

A specimen of  
the Upper Devonian  
internal structure  
the Family Osteichthyes  
thought to be  
also be considered  
that the known  
restrial locomotion  
proposed according  
evolved with a

POSTILLA 15

**NEW EVIDENCE ON THE EVOLUTION OF THE PAIRED FINS OF  
RHIPIDISTIA AND THE ORIGIN OF THE TETRAPOD LIMB, WITH  
DESCRIPTION OF A NEW GENUS OF OSTEOLEPIDAE**

KEITH STEWART THOMSON

Department of Biology and Peabody Museum of Natural History  
Yale University, New Haven, Connecticut 06520

(Received April 25, 1972)

**ABSTRACT**

A specimen of *Sterropterygion brandei*, gen. et sp. nov., a rhipidistian from the Upper Devonian of Pennsylvania, shows for the first time the detailed internal structure of the pectoral and pelvic fins and girdles in a member of the Family Osteolepidae. The structure conforms to the general pattern once thought to be directly antecedent to that of tetrapods but which now must also be considered an ancient feature of rhipidistian fishes. It is contended that the known Rhipidistia could not support their own weight during terrestrial locomotion through fin action alone and a scheme of evolution is proposed according to which the paired fins of osteolepids and tristicopterids evolved with a dual function: in locomotion and support of lung ventilation.

POSTILLA 157: 7p.

19 JULY 1972

## INTRODUCTION

In studies of the fish-tetrapod transition, great attention has been paid to the paired fins and girdles of the ancestral group of fishes — the Devonian Rhipidistia. Andrews and Westoll (1970a,b) have recently summarized our knowledge of the postcranial skeleton of Rhipidistia and contributed greatly to our understanding of the early evolution of the pentadactyl limb. However, all such studies have had to depend heavily upon a single fish — the lowermost Upper Devonian *Eusthenopteron foordi* Whiteaves (Family Tristicopterae, in the classification used by Thomson, 1969) and comparative materials have been scarce. The recent discovery of a well-preserved member of the rhipidistian Family Osteolepidae (Figs. 1-5), showing for the first time both the pectoral and pelvic fins, with girdles, provides an important source of new data.

The specimen with which the present notice is concerned was collected by Mr. Scott Brande at a highway cut outcropping of an Upper Devonian formation in Lycoming County, Pennsylvania. The present preliminary account will be followed at a later date by a full description and analysis. The new osteolepid fish represents a new genus and species. A brief taxonomic diagnosis is necessary.

CLASS OSTEICHTHYES  
ORDER CROSSOPTERYGII  
FAMILY OSTEOLEPIDAE  
Genus *Sterropterygion* nov.

TYPE SPECIES. *Sterropterygion brandei* Thomson, nov.

PRELIMINARY DIAGNOSIS. Rhipidistian with dermal bones and rhomboid scales bearing complete external covering of enameloid and dentine typical of the Osteolepidae. Estimated total length 380 mm. First dorsal fin inserted slightly behind the level of the pelvic insertion, and second dorsal in front of, or level with, the anal. Number of dorsolateral scale rows in front of first




FIG. 1. Reconstruction of the general features of *Sterropterygion brandei* gen. et sp. nov., Holotype.

PAIRED FIN EV

dorsal, second dors  
lateral scale row v  
Dorsomedian scale  
length of anterior d  
length of the ethmo

HORIZON AND LOCAL  
Catskill Formation,

DERIVATIO NOMINIS

HOLOTYPE and sole

DIAGNOSIS AND OCCU

DERIVATIO NOMINIS

The right pectoral f  
The internal fin ske  
dyle, a long thin  
elements. The ventr  
of projections mark  
pectoralis. The dors  
insertion of the d  
*Eusthenopteron* in t  
opposed to approxi  
ternal skeleton, the  
a massive entepicon  
ridge into a series o  
and elongate radius  
lated with the ulna.

The pelvic struct  
The internal fin ske  
and shorter, particu  
half the total length  
The osteolepid pelv  
of *Sterropterygion*  
long anterior iliac p  
to that of *Eusthenop*  
dorsal public proces

dorsal, second dorsal and pelvic fins: 41, 53, 50 respectively. Tenth dorso-lateral scale row with 6 scales above and 5 below the lateral line scale. Dorsomedian scale row count: anterior division  $41 \pm 1$ , median division 6; length of anterior division approximately 4.7 times longer than the estimated length of the ethmoidal division of the skull roof.

**HORIZON AND LOCALITY.** The Upper Devonian Susquehanna Group, probably Catskill Formation, of Northern Lycoming County, Pennsylvania.

**DERIVATIO NOMINIS:** from the Greek for "firm-fin."

*Sterropterygion brandei* sp. nov.

**HOLOTYPE** and sole known specimen: YPM 6721, incomplete fish.

**DIAGNOSIS AND OCCURRENCE.** As for the genus above.

**DERIVATIO NOMINIS:** after the discoverer, Mr. Scott Brande.

DISCUSSION


The right pectoral fin of *Sterropterygion brandei* (Fig. 4) is almost complete. The internal fin skeleton consists of a stout humerus with a large entepicondyle, a long thin radius, and a short ulna supporting four more distal elements. The ventral surface of the humerus is marked by a prominent row of projections marking the insertion of the ventral flexor musculature and M. pectoralis. The dorsal side bears only moderately developed processes for the insertion of the deltoid muscles. The pectoral fin differs from that of *Eusthenopteron* in that the lobed part occupies just slightly less than half (as opposed to approximately one third) the total length of the fin. In the internal skeleton, the new fish is unique in its short and broad humerus with a massive entepicondylar platform, the specialization of the ventral humeral ridge into a series of large discrete processes for muscle insertion, the slender and elongate radius, and the great breadth of the postaxial elements articulated with the ulna.

The pelvic structures of *Sterropterygion brandei* are shown in Figure 5. The internal fin skeleton differs from that of *Eusthenopteron* in being broader and shorter, particularly in the distal elements. The pelvic fin lobe occupies half the total length of the fin (as opposed to one third in *Eusthenopteron*). The osteolepid pelvic girdle previously has been completely unknown; that of *Sterropterygion brandei* consists of a pair of arch-shaped elements with long anterior iliac processes coming close together in the midline. It is similar to that of *Eusthenopteron*, differing only in the slightly greater breadth of the dorsal public process.

s been paid to the  
e Devonian Rhip-  
summarized our  
contributed greatly  
actyl limb. How-  
single fish — the  
whiteaves (Family  
1969) and com-  
a well-preserved  
-5), showing for  
provides an im-

was collected by  
Devonian forma-  
nary account will  
analysis. The new  
brief taxonomic

s and rhomboid  
d dentine typical  
dorsal fin inserted  
dorsal in front of,  
in front of first



*on brandei* gen. et

The discovery of *Sterropterygion brandei* reveals the remarkable fact that there are no fundamental points of difference in pectoral or pelvic structure between two families of Rhipidistia, the Osteolepidae and Tristicopterae, that have been distinct since least the Middle Devonian. (This confirms the observation of Andrews and Westoll, 1970b, based on fragmentary pectoral material of the Carboniferous osteolepid *Megalichthys*.) Moreover, there is evidence that the osteolepids range back to the Lower Devonian (undescribed material from China reported to be in the process of description, several years ago, by Mi-man Chang). From the earliest fossil record of the Rhipidistia there seem to have been two distinct patterns of paired fin development. The holoptychoids (Porolepidae and Holoptychidae) had a monoseriate pattern similar to that of lungfish and coelacanth and it is extremely unlikely that they gave rise to any tetrapods (Schultze, 1970). The biseriata condition seen in osteolepids and tristicopterids is equally ancient and must be considered a primitive, rather than derived, character in these lines. Thus, the biseriata condition of the paired fins, which shows great similarity to that of the Amphibia in the proximal elements (but not the distal elements) must be primarily an adaptation for life in water. It is not a specialization of the immediate tetrapod ancestors and it is extremely unlikely that this condition evolved in response to selection involving specific factors of semiterrestrial existence.

The following interpretation of the evolution of the tetrapod limb is suggested by this new evidence. The paired fins of Rhipidistia seem to be adapted primarily for slow swimming movements, much as in the modern lobe-finned chondrosteian *Polypterus*. Presumably, at an early point in their history, the Rhipidistia began to make excursions overland. The possible environmental and adaptive contents have been reviewed by Thomson (1969) and Andrews and Westoll (1970a). But it is extremely doubtful that the pectoral and pelvic limbs were sufficiently developed that these fish could "walk," that is, move through the actions of the paired fins alone. Furthermore, in all known Rhipidistia there was a large fin web that would have been a severe hindrance to attempts at rapid terrestrial movement, primarily by means of limb movements with the fin sharply flexed ventrally to raise the trunk from the ground.

It has been pointed out previously (Thomson, 1969) that, in their experimental movements overland, the prime new mechanical problem facing the lobe-finned fishes was not locomotion but lung ventilation. Locomotion was always possible by lateral body undulation, but lung ventilation required that the anterior trunk (at least) be raised up so that the weight of the body did not crush the lungs. While the pectoral limbs of Rhipidistia do not seem to have been strong enough to contribute greatly to active terrestrial locomotion, they were probably strong enough to lift up the anterior trunk when the animal was at rest. Thus the paired limbs of Rhipidistia may have been subject to two separate selective pressures. First, the fins were used as relatively immobile props in terrestrial locomotion. The mode of progression

was based on the normal musculature could also be used if the fin touched the ground. The opposite fin was raised in addition to this action. The fins themselves would not be a net raising of the trunk. The fish was at rest, however, and could be through the direct action of the considerably less strength of the locomotion and perhaps the evolution. Any increase in strength noting that the latter function and this may be a partial adaptation of the pentadactyl condition.

Only significantly later and further modification of the carpus and tarsus for weight support. The robustness of *brandei* may indicate that but it remains essentially the same as be found in animals of this organization.

I am deeply grateful to Mr. M. and its donation to the F. Bowen, University of specimen and kindly been possible without the by Mr. Peter Whybrow. and the study supported

Andrews, S. M. and T. S. *opteron foordi* Whiteav. 1970b. The postcranium of *opteron*. Trans. Roy. Soc. Edinb. 102: 1-10.  
Schultze, H-P. 1970. Fossil fishes. Novitates 2408: 1-10.  
Thomson, K. S. 1969. The

kable fact that pelvic structure risticopterae, is confirms the entary pectoral reover, there is n (undescribed rription, several record of the of paired fin chidae) had a nths and it is hultze, 1970). rids is equally ed, character in ich shows great s (but not the ater. It is not a remely unlikely specific factors

od limb is sug- n to be adapted ern lobe-finned eir history, the environmental ) and Andrews oral and pelvic ' that is, move in all known been a severe y by means of the trunk from

at, in their ex- problem facing n. Locomotion ilation required ght of the body ia do not seem restrial locomo- or trunk when may have been were used as of progression

was based on the normal sinuous movement of the body, but the trunk musculature could also apply force around the fixed point where each paired fin touched the ground. In this way a lever effect was produced and also the opposite fin was raised off the ground as it was being swung forward. Any addition to this action that could be made by the muscles and skeleton of the fins themselves would be favored. Since this type of locomotion involves a net raising of the trunk, lung ventilation would be facilitated. When the fish was at rest, however, the only means for raising the anterior trunk would be through the direct action of the fins themselves. This would require considerably less strength and mobility than a similar action during active locomotion and perhaps was possible from an early stage in rhipidistian evolution. Any increase in this capacity would be favored, but it is worth noting that the latter function applies more to the pectoral than pelvic fins and this may be a partial explanation of their differential development toward the pentadactyl condition.

Only significantly later (Late Devonian) came reduction of the fin web and further modification of the internal fin skeleton (including development of the carpus and tarsus) allowing greater mobility and more extensive weight support. The robust development of the paired fins in *Sterropterygion brandei* may indicate that it belongs to an advanced stage in this sequence, but it remains essentially a fish and one must expect that the final stages will be found in animals that were more amphibian than piscine in overall organization.

#### ACKNOWLEDGMENTS

I am deeply grateful to Mr. Scott Brande for his careful collection of the fish and its donation to the Peabody Museum of Natural History. Professor Z. D. Bowen, University of Rochester, also recognized the importance of the specimen and kindly brought it to my attention. The study would not have been possible without the extraordinarily delicate and painstaking preparation by Mr. Peter Whybrow. The photographs were taken by Mr. A. H. Coleman and the study supported by National Science Foundation Grant GB 28823X.

#### LITERATURE CITED

- Andrews, S. M. and T. S. Westoll. 1970a. The postcranial skeleton of *Eusthenopteron foordi* Whiteaves. *Trans. Roy. Soc. Edinb.* 68: 207-329.  
 1970b. The postcranial skeleton of rhipidistian fishes excluding *Eusthenopteron*. *Trans. Roy. Soc. Edinb.* 68: 391-489.  
 Schultze, H-P. 1970. Folded teeth and the monophyletic origin of tetrapods. *Novitates* 2408: 1-10.  
 Thomson, K. S. 1969. The biology of lobe-finned fishes. *Biol. Rev.* 44: 91-154.




FIG. 2. General view of the right flank of *Sterropterygion brandei* gen. et sp. nov., Holotype.  $\times 0.55$ .




FIG. 3. Detail of the right pectoral fin of *Sterropterygion brandei* gen. et sp. nov., Holotype.  $\times 1.2$ .

PAIRED FIN EV



FIG. 4. The pectoral fin of *Sterropterygion brandei*. Scales removed to show structure.



FIG. 5. The pelvic fin of *Sterropterygion brandei*. Prepared specimen.





FIG. 4. The pectoral fin of *Sterropterygion brandei* gen. et sp. nov., Holotype. Scales removed to show internal elements, ventral.  $\times 1.0$ .




FIG. 5. The pelvic fin and girdle of *Sterropterygion brandei* gen. et sp. nov., Holotype. Prepared specimen showing both girdles and the left pelvic fin.  $\times 1.0$ .

## INFORMATION FOR AUTHORS

- REVIEW** The Publications Committee of the Peabody Museum of Natural History reviews and approves manuscripts for publication. Papers will be published in approximately the order in which they are accepted; delays may result if manuscript or illustrations are not in proper form. To facilitate review, the original and one carbon or xerox copy of the typescript and figures should be submitted. The author should keep a copy.
- STYLE** Authors of biological papers should follow the *Style Manual for Biological Journals*, Second Edition (Amer. Inst. Biol. Sci.). Authors of paleontological manuscripts may choose to follow the *Suggestions to Authors of the Reports of the U.S. Geological Survey*, Fifth Edition (U.S. Govt. Printing Office).
- FORM** Maximum size is 80 printed pages including illustrations (= about 100 manuscript pages including illustrations). Manuscripts must be typewritten, with wide margins, on one side of good quality 8½ x 11" paper. *Double space everything. Do not underline anything except genera and species.* The editors reserve the right to adjust style and form for conformity.
- TITLE** Should be precise and short. Title should include pertinent key words which will facilitate computerized listings. Names of new taxa are not to be given in the title.
- ABSTRACT** The paper must begin with an abstract. Authors must submit completed BioAbstract forms; these can be obtained from the *Postilla* editors in advance of submission of the manuscripts.
- NOMENCLATURE** Follow the International Codes of Zoological and Botanical Nomenclature.
- ILLUSTRATIONS** Must be planned for reduction to 4¼ x 7" (to allow for running head and two-line caption). If illustration must go sideways on page, reduction should be to 4 x 7¼". All illustrations should be called "Figures" and numbered in arabic, with letters for parts within one page. It is the author's responsibility to see that illustrations are properly lettered and mounted. Captions should be typed double-spaced on a separate page.
- FOOTNOTES** Should not be used, with rare exceptions. If unavoidable, type double-spaced on a separate page.
- TABLES** Should be numbered in arabic. Each must be typed on a separate page. Horizontal rules should be drawn lightly in pencil; vertical rules must not be used. Tables are expensive to set and correct; cost may be lowered and errors prevented if author submits tables typed with electric typewriter for photographic reproduction.
- REFERENCES** The style manuals mentioned above must be followed for form and for abbreviations of periodicals. Double space.
- AUTHOR'S COPIES** Each author receives 50 free copies of his *Postilla*. Additional copies may be ordered at cost by author when he returns galley proof. All copies have covers.
- PROOF** Author receives galley proof and manuscript for checking printer's errors, but extensive revision cannot be made on the galley proof. Corrected galley proof and manuscript must be returned to editors within seven days.
- COPYRIGHT** Any issue of *Postilla* will be copyrighted by Peabody Museum of Natural History only if its author specifically requests it.