



Future directions in bladder cancer immunotherapy: towards adaptive immunity

The clinical management of bladder cancer has not changed significantly in several decades. In particular, intravesical bacillus Calmette–Guérin (BCG) immunotherapy has been a mainstay for high-risk nonmuscle invasive bladder cancer since the late 1970s/early 1980s. This is despite the fact that bladder cancer has the highest recurrence rates of any cancer and BCG immunotherapy has not been shown to induce a tumor-specific immune response. We and others have hypothesized that immunotherapies capable of inducing tumor-specific adaptive immunity are needed to impact bladder cancer morbidity and mortality. This article summarizes the preclinical and clinical development of bladder cancer immunotherapies with an emphasis on the last 5 years. Expected progress in the near future is also discussed.

First draft submitted: 30 October 2015; Accepted for publication: 16 December 2015; Published online: 10 February 2016

Keywords: adaptive • BCG • bladder cancer • checkpoint inhibitors • cytokines • IL-12 • immunotherapy • memory • Toll-like receptors

Bladder cancer overview

Bladder cancer is the 5th most common cancer in the United States with an estimated 74,000 new cases, 16,000 deaths and a total domestic prevalence of more than 570,000 in 2015 [1,2]. Most new cases (70–80%) are diagnosed as nonmuscle invasive bladder cancer (NMIBC) with an associated 15-year survival of 62–95% [3]. Low-risk NMIBC, typically represented by small, papillary tumors confined to the urothelium, is well-managed via transurethral resection of the bladder tumor (TURBT) often followed by intravesical instillation of mitomycin C. High-risk NMIBC, including high-grade carcinoma *in situ* (CIS) and tumors invading the lamina propria, is treated via TURBT followed by intravesical immunotherapy with *mycobacterium bovis* bacillus Calmette–Guérin (BCG) [4–6]. A significant minority of patients (20–30%) initially present with one or more tumors that have invaded the muscle layer of the bladder. Because muscle

invasive bladder cancer (MIBC) can rapidly progress to metastatic disease, cystectomy, or surgical removal of the bladder, is standard of care.

Despite the fact that bladder cancer is diagnosed relatively early in the course of disease, it has the highest recurrence rate of any malignancy at 50–80%. These high recurrence rates necessitate long-term maintenance therapy and regular surveillance [7,8] which in turn causes bladder cancer to have the highest lifetime treatment costs per patient of all cancers [9].

Immunologic challenges in bladder cancer

Bladder cancer, with the third highest rate of somatic mutations, is one of the most immunogenic cancers [10]. Yet bladder cancer is able to evade immune-mediated elimination even in the presence of antigen-specific immune cell infiltration. Understanding the immunoevasive strategies employed by blad-

Sean G Smith¹ & David A Zaharoff^{*1}

¹Department of Biomedical Engineering, University of Arkansas, Fayetteville, AR, USA

*Author for correspondence:

Tel.: +1 479 575 5221

Fax: +1 479 575 4346

zaharoff@uark.edu

der tumors is key to developing therapies capable of inducing adaptive responses. This section will provide a high level overview of some of the challenges associated with generating adaptive immunity in the bladder as well as some of the means employed by bladder tumors to evade destruction.

The first challenge in inducing an adaptive response to bladder cancer is the unique immunological milieu within the organ itself. The bladder epithelium is infiltrated by mast cells, macrophages, dendritic cells and T cells, but the overall organization of the immune system within the bladder is the result of a balance between conflicting needs. On one hand, the lumen is routinely exposed to noncommensal bacteria and must implement strategies to subvert infection. On the other hand, the bladder must be able to store high concentrations of self-antigens and toxins without eliciting an immune response. The bladder's balanced strategy includes a thick mucin layer, secretion of antibacterial agents and rapid micturition to limit penetration of invading microorganisms while establishing an immunosuppressive environment to limit undesirable immune responses [11]. Because of this unique milieu, the establishment of a tumor-specific adaptive immunity in the bladder is difficult, although not impossible as will be discussed later in this review.

Another obstacle to immunotherapy is the immunosuppressive bladder tumor microenvironment. Bladder tumors can anergize tumor infiltrating lymphocytes (TILs) [12] and promote the accumulation of immunosuppressive myeloid cells. Bladder tumors have also been associated with high levels of regulatory T-cells (Tregs) and T_H1 inhibitory cytokines such as IL-10 [13]. Horn *et al.* showed that higher FOXP3:CD3 and FOXP3:CD8 ratios in bladder tumor infiltrates correspond to worse overall survival in patients who underwent radical cystectomy, suggesting that infiltration by Tregs may support tumor invasion [14]. However, a retrospective study by Winerdal *et al.* showed that a higher FOXP3⁺ TIL density correlated with improved outcomes while FOXP3⁺ tumor cells corresponded with poorer outcomes [15]. A recent review covers the influence of TILs in bladder cancer in greater detail [16].

Bladder cancers, like many cancers, utilize immune checkpoints to modulate immunity. Specifically, bladder tumors can promote immune tolerance by overexpressing immune checkpoint ligands capable of inhibiting activated T cells. The most commonly investigated immune checkpoint molecules are PD-1, PD-L1 and CTLA-4, but there are multiple other regulatory molecules, such as LAG-3 and TIM-3, that are potential targets for bladder cancer immunotherapy [17]. Several clinical studies have shown that bladder tumors and infiltrating immune cells exhibiting increased expres-

sion of PD-L1 and PD-1 are associated with poorer outcomes [18–21]. Investigations into the effects of CTLA-4 expression have been limited with conflicting opinions as to whether the CTLA-4 +49 A>G polymorphism increases or decreases bladder cancer risk [22–24].

An additional challenge is that bladder tumors, like many other cancers, can produce escape variants by removing their surface expression of major histocompatibility complex I (MHC I) molecules [25]. MHC I is essential for displaying tumor antigens vital for recognition by T-cells. Several studies have shown that bladder tumors can be heterogeneous in their display of MHC I [26]. Some express normal levels of MHC I and are associated with better outcomes. Other, so called 'soft', tumors have low MHC I expression that can be upregulated to normal levels through IFN-mediated mechanisms. Still others, termed "hard" tumors, have permanent alterations in MHC I expression due to a genetic mutation in the processing machinery or one of the MHC I components. There is some evidence that while both normal and soft tumors are BCG responsive, MHC I negative hard tumors may be BCG resistant leading to tumor escape [27,28]. It should be noted that any truly MHC I negative tumor should be susceptible to natural killer (NK) cell mediated cytotoxicity yet the mechanism by which these tumors might escape NK cell surveillance has not yet been described.

Limitations of BCG immunotherapy

Since the pioneering work of Morales *et al.* in 1976 [29], BCG has served as the standard intravesical immunotherapy for high-risk NMIBC. On one hand, its longevity can be used to highlight its effectiveness in the management of NMIBC. However, as mentioned previously, bladder cancer has the highest recurrence rate of all cancers and the four-decade-long lack of progress in the treatment of NMIBC underscores the need for new and more effective therapies.

Intravesically administered BCG promotes an inflammatory reaction in the bladder wall that is known to reduce recurrence rates. However, the exact mechanism(s) by which BCG exerts its antitumor effects is unknown. A suggested model of BCG immunotherapy is outlined in an excellent review by Redelman-Sidi *et al.* [30]. Briefly, BCG attaches to urothelial cells via fibronectin and integrin $\alpha 5\beta 1$ before internalization by bladder cancer cells through micropinocytosis. BCG infection can decrease the proliferation of bladder cancer cells and is cytotoxic only at high concentrations. Thus, it is likely that the majority of BCG's antitumor activity is accomplished indirectly. Infected bladder cancer cells secrete a number of cytokines including IL-6, IL-8, GM-CSF and TNF- α . BCG infection also causes the upregulation

| Table 1. Summary of new therapeutics being investigated for bladder cancer. | | | | | |
|---|---|---------------|-------------------------------|------------------|----------------------------|
| Therapeutic strategy | Therapeutic form | Delivery site | Furthest stage of development | Disease targeted | Ref. |
| TLR agonists | TLR-9 agonist CpG | IT | Preclinical | NMIBC | [36] |
| | TLR-2 agonist HP-NAP | IVes, IT | Preclinical | NMIBC | [37] |
| | TLR-7 agonist TMX-101 | IVes | Phase II | NMIBC, CIS | [38] |
| Cancer vaccines | DC-targeted tumor antigens | SC | Phase II | MIBC | [45,46] |
| | Natural adjuvants + <i>HER2/NEU</i> plasmid | SC | Preclinical | NMIBC | [39–42] |
| | Peptide vaccine | SC | Phase II | NMIBC | [44] |
| | shRNA for FOXO3 + <i>HER2/NEU</i> plasmid | SC | Preclinical | NMIBC, MIBC | [43] |
| | Fowlpox vector, PANVAC | SC | Phase II | NMIBC | NCT02015104 |
| | HS-410 cell line | SC | Phase I/II | NMIBC | NCT02010203 |
| IFN- α | Recombinant | IVes | Phase II | NMIBC | [50–55] |
| | rBCG | IVes | Preclinical | NMIBC | [56,58] |
| | Virus | IVes | Phase II | NMIBC | [57] |
| GM-CSF | Vaccine | IVes, SC | Preclinical | NMIBC, Met | [59–62] |
| | Oncolytic virus | IVes | Phase II | NMIBC | [63] |
| IL-2 | Recombinant | IVes | Phase I/II | NMIBC | [67,70] |
| | Vaccine | IVes, SC | Preclinical | NMIBC, Met | [61,68,69] |
| | p53 TCR fusion protein | IV | Phase II | NMIBC, MIBC | NCT01326871 NCT01625260 |
| IL-12 | Recombinant | IVes | Phase I | NMIBC | [74] |
| | Recombinant + chitosan | IVes | Preclinical | NMIBC | [75,76] |
| | Plasmid | IVes, IT | Preclinical | NMIBC | [72,73,78] |
| | Retrovirus | SC | Preclinical | NMIBC | [77] |
| IL-15 | Plasmid | IVes | Preclinical | NMIBC | [79] |
| | Fusion protein (ALT-803) + BCG | IVes | Phase I/II | NMIBC | [80] |
| Checkpoint inhibitors | α CTLA-4 | IV | Phase II | Met | [24,81] |
| | α PD-L1 | IV | Phase II | Met | [82] |
| | α CTLA-4, α PD-L1 + CpG | IT | Preclinical | NMIBC | [36] |
| Anti-CD40 | Anti-CD40 | IT | Phase I | NMIBC, Met | [83] |
| Anti-IL-10 | Anti-IL-10 | IV | Preclinical | NMIBC, Met | [84–86] |

IT: Intratumoral; IV: Intravenously or systemically; Ives: Intravesically; Met: Metastatic; MIBC: Muscle-invasive bladder cancer; NMIBC: Non-muscle-invasive bladder cancer; SC: Subcutaneous.

of MHC II and ICAM-1 on bladder cancer cells, thus enhancing their antigen presenting capabilities. The initial cytokine burst leads to the recruitment of granulocytes, monocytes and macrophages and the subsequent release of additional cytokines including IL-1, IL-2, IL-5, IL-6, IL-8, IL-12, IL-18, TNF- α , IFN- γ and GM-CSF. This second cytokine burst leads to the recruitment and activation of NK cells, CD4⁺ helper T cells (T_H) and CD8⁺ cytotoxic T cells (CTLs). Tumor cell killing is likely primarily accomplished by NK cells, CD8⁺ T cells, macrophages and granulocytes through the release of TRAIL. Despite the involvement

of T lymphocytes, BCG immunotherapy has not been shown to induce a tumor-specific immune response.

Overall, 20–30% of patients will fail initial BCG therapy and 30–50% of BCG responders will develop recurrent tumors within 5 years [31]. An incomplete understanding of BCG's antitumor mechanisms has made it difficult: 1) to predict responses to BCG; 2) to optimize BCG immunotherapy; and 3) to understand why some patients ultimately fail BCG immunotherapy. Separately, BCG is a live pathogen and causes severe infections, including life threatening sepsis, in approximately 5% of patients [32]. At least seven

patients have died from BCG sepsis following bladder instillations [32] and 30–40% of patients discontinue intravesical BCG immunotherapy due to local and systemic toxicities [33]. Lastly, despite four decades of experience, it is still not clear which of the 6+ strains of BCG is most effective at treating NMIBC.

Novel strategies for bladder cancer immunotherapy

The aforementioned challenges associated with initiating an adaptive immune response within the bladder and bladder cancer milieu can be overcome with appropriate strategies. An effective strategy should: 1) reverse the immunosuppressive environment of the bladder tumor; 2) initiate an inflammatory response at the tumor site capable of eliminating the primary tumor; 3) activate tumor antigen-specific T-cells capable of eliminating secondary local or distant lesions; and 4) establish a reservoir of memory T-cells with polyclonal specificity to monitor for recurrent tumors. There is a long history of strategies that have attempted to overcome one or more of these challenges. The goal of this section is to present an overview of recent (published within the last 5 years) strategies being investigated in preclinical and clinical settings for the treatment of superficial and metastatic bladder cancer and to provide commentary on those strategies' ability or potential to establish an adaptive memory response. A summary of immunotherapy strategies is given in [Table 1](#).

TLR agonists

Perhaps inspired by BCG, one of the most common strategies investigated in preclinical studies is the activation of innate immune pathways [34,35]. Toll-like receptor (TLR) pathways are of particular interest in cancer immunotherapy. TLRs are a family of receptors that bind to common components of many pathogens as well as signals released by damaged cells. TLRs are expressed on many innate immune cells, including dendritic cells. The most potent of all antigen presenting cells, dendritic cells play a pivotal role in bridging the innate and adaptive responses. Thus, targeting dendritic cell TLRs is a common strategy for enhancing adaptive immune responses. TLRs are also present on a large portion of bladder tumors where higher TLR expression is correlated with less invasive tumors [35]. With their wide expression on a variety of cells and their ability to link adaptive and innate immune responses, TLR ligands make attractive candidates for potential bladder cancer therapeutics. As such, several TLR ligands have been investigated at both the clinical and preclinical level. There are more than 15 years of studies regarding TLR ligands against bladder cancer, and for a more complete picture please see the excellent

review by LaRue *et al.* [35]. The following will provide a brief update of findings within the last 5 years.

TLR9 agonistic CpG oligodeoxynucleotides were investigated preclinically in conjunction with checkpoint inhibitors against a subcutaneous (s.c.) MB49 bladder tumor model. CpG, administered peritumorally, in combination with intraperitoneal (i.p.) anti-CTLA-4, anti-PD-1 or anti-PD-L1 antibodies resulted in 6/7, 5/7 and 2/7 long-term survivors, respectively. All combinations of CpG and checkpoint inhibition were more effective than combinations of BCG plus checkpoint inhibition as well as checkpoint inhibition alone [36]. The TLR2 agonist HP-NAP, a bacterial protein produced by *helicobacter pylori*, was tested against both subcutaneous and orthotopic MB49 tumors in mice. In these studies, HP-NAP increased both the numbers and percentages of CD4⁺ and CD8⁺ T cells in local lymph nodes and limited tumor growth more efficiently than BCG [37].

In the clinic, the TLR7 agonist imiquimod (TMX-101) was well tolerated by 7 NMIBC patients in a Phase I study [38]. A Phase II trial with TMX-101 was recently completed against CIS (NCT01731652). No data from this study have been published at the time this review was published. Overall, TLR agonists and other stimulants of the innate immune system hold promise as adjuvants for initiating the antitumor response but will likely need to be used in conjunction with other therapeutics.

Cancer vaccines

Therapeutic cancer vaccines are under development to induce adaptive immune responses to one or more tumor-associated antigens. Several different forms of cancer vaccines have been explored recently. Vaccines comprised of tumor lysates are easy to produce and have the potential to induce polyclonal responses against both known and unknown tumor antigens. However, because tumor lysates are poorly immunogenic, the evaluation and inclusion of effective adjuvants is critical. The antimicrobial peptide shrimp anti-lipopolysaccharide factor (SALF) has been evaluated preclinically as an adjuvant for a tumor lysate vaccine. Mice were given SALF mixed with MBT-2 tumor cell lysate on days 7, 14 and 21 before subcutaneous inoculation with MBT-2 on day 28. Tumors developed in mice treated with MBT-2 lysate alone (5/7) and SALF alone (7/7), but not in those treated with the mixture (0/7). Tumor-specific memory was shown via CTL assays using splenocytes from treated mice [39]. A similar study investigated a different antimicrobial peptide, GE33. Using the same treatment schedule, tumor development was more pronounced in mice treated with MBT-2 lysate alone (5/7) and SALF alone

(7/7) than those treated with the mixture (2/7) [40]. In the last 5 years, there have been no clinical trials using tumor lysate-based vaccines against bladder cancer.

Similarly, tumor antigen encoding plasmids are a useful source of tumor antigen, however, adjuvants are needed to induce a robust immune response. Extracts from the mushroom *Clitocybe nuda* have been shown to activate dendritic cells. Mice bearing s.c. MBT-2 which express p185neu received three weekly intramuscular injections of *HER2/NEU*-encoding plasmids with or without *C. Nuda* extract [41]. The addition of *C. Nuda* extract increased the number of survivors at 60 days from 2/7 to 5/7. Untreated mice and mice receiving the extract alone were moribund by day 35. The spleens of treated mice were analyzed for response to *HER2* peptide stimulation. There were increased percentages of activated CD8⁺ T-cells in the combination group compared to either therapy alone. An *in vitro* stimulation of bone marrow-derived dendritic cells demonstrated that *C. Nuda* was able to enhance immunomodulatory markers (CD40, CD80, CD86, MHC I and MHC II) at levels similar to LPS.

In a similar study, Ling Zhi-8 (LZ-8) isolated from *Ganoderma lucidum* (Lingzhi mushroom) was used as an adjuvant for *HER2/NEU*-encoding plasmids against s.c. MBT-2 tumors [42]. LZ-8 was found to stimulate dendritic cells through TLR4 and enhanced the ability to induce an antigen-specific immune response in an *in vitro* experiment. In antitumor studies, LZ-8 was administered i.p. while the *HER2/NEU* gene was given once a week for three weeks. LZ-8 did not improve long-term survival, however, plasmid alone or plasmid plus LZ-8 showed extended survival relative to untreated mice or LZ-8 alone.

Another strategy to enhance immune responses to plasmid-based vaccines utilized small hairpin RNA (shRNA) to silence *FOXO3*, an immunosuppressive transcription factor. Both silencing and *HER2/NEU* plasmids were administered once a week for 3 weeks to mice bearing MBT-2 tumors. The inclusion of *FOXO3* silencing increased the number of mice surviving >70 days from 1/7 to 5/7. Antitumor responses were mediated by CD8⁺, but not CD4⁺ cells. In an experimental pulmonary metastases model, lung weights were significantly lower in both *HER2/NEU* and *HER2/NEU* plus *FOXO3* shRNA groups when compared with untreated controls [43]. To our knowledge, no clinical trial for bladder cancer in the last 5 years has evaluated a plasmid-based cancer vaccine.

Peptide-based cancer vaccines are useful for generating immune responses against unique and well-defined epitopes. In a Phase I study in Japan, peptides from bladder cancer oncoproteins, MPHOSPH1 and DPEPDCI, most likely to bind to the most common

HLA-A type in the Japanese population were identified via BIMAS HLA peptide binding predictions [44]. Six patients with unresectable bladder cancer were vaccinated with 1 mg of each peptide in incomplete Freund's adjuvant once a week for 4 weeks. The peptide vaccines were well-tolerated with no severe adverse events. There were peptide-specific CD8⁺ responses against MPHOSPH1–278 and DEPDC1–294 peptides in 3/4 and 4/6 evaluated patients, respectively. A subsequent Phase II study is ongoing (NCT00633204).

CDX-1307 is a fusion molecule in which a mannose receptor targeted antibody is linked to a tumor antigen, β -hCG. Targeting the mannose receptor on antigen presenting cells is expected to increase the likelihood that β -hCG is taken up, processed and displayed to antigen-specific T cells. Two Phase I studies evaluating CDX-1307 in patients with advanced cancers, including bladder cancer, were recently completed, however, no published data could be found. A Phase II study (NCT01094496) to investigate the efficacy of CDX-1307 given against muscle invasive disease began recruitment in 2010, but was recently terminated due to slow enrollment [45].

A related molecule, CDX-1401, which is comprised of human monoclonal antibody specific for DEC-205, a dendritic cell receptor, fused to the full-length tumor antigen NY-ESO-1, was the subject of a recently completed Phase I study in patients with advanced malignancies, including advanced bladder cancer [46]. The vaccine was well tolerated and 2/45 patients experienced a >20% tumor regression while 13/45 patients exhibited stable disease. It could not be determined if any of these responses were in bladder cancer patients.

Two additional bladder cancer vaccines of interest are being evaluated in clinical studies with no published results at the time this review was written. An ongoing Phase I study will evaluate a fowlpox vector expressing carcinoembryonic antigen, mucin-1 and three costimulatory molecules with and without BCG in high-grade NMIBC patients who have failed BCG (NCT02015104). An ongoing Phase I/II study will assess HS-410, a vaccine comprised of irradiated bladder cancer cells engineered to secrete gp96, alone or in combination with BCG in NMIBC patients (NCT02010203)

Cytokines

Cytokines are potent nodes of communication within the immune cell network. Their potential to act as immunotherapeutics has long been recognized but translation of cytokines into the clinic has faced some significant obstacles. In particular, widespread clinical application of cytokine-based cancer immunotherapies has been prevented by inappropriate systemic deliv-

ery of proinflammatory cytokines. Nearly all clinical trials, both past and present, evaluating cytokine monotherapies utilize systemic, that is, intravenous or s.c., injections. However, cytokines function primarily through paracrine and autocrine mechanisms and thus are rarely measurable in the circulation of healthy individuals. In addition, cytokine communication networks are far more complex than initially conceived. Cytokines are pleiotropic and often redundant which can lead to off target effects. In sum, the systemic delivery of multifunctional cytokines risks unintentional signaling and dose-limiting toxicities.

Fortunately, cytokine-based bladder cancer immunotherapies have significant potential due to the bladder's accessibility and ability to sequester high concentrations of cytokines with limited systemic exposure. The following sections will cover recent studies in which cytokines in a variety of formats including recombinant protein, viral vectors, plasmids, transfected BCG and transfected tumor cells are used for the treatment of bladder cancer. Many of the cytokines investigated for intravesical therapy have been inspired by downstream molecules released after BCG treatment including IFN- α , GM-CSF, IL-2 and IL-12.

IFN- α

IFN- α is a type 1 interferon produced primarily by innate immune cells in response to invading microbes, particularly viruses [47]. IFN- α 2b is approved as a monotherapy for malignant melanoma, hairy cell leukemia and Kaposi's sarcoma [48]. IFN- α binds to a heterodimeric receptor complex that is expressed on the surface of most cell types, including many cancer cells. Thus, IFN- α can act directly upon cancer cells to induce a range of effects including apoptosis and upregulation of tumor cell-surface antigens. Indirect antitumor activities of IFN- α include dendritic cell maturation.

IFN- α , as both a monotherapy and in combination with BCG, has been widely explored against bladder cancer including as the subject of a number of clinical trials [49]. Despite promising preclinical results, the consensus of clinical studies is that IFN- α alone is no more effective than current treatments at preventing recurrence of NMIBC and in some instances is less effective. One recent example showed that alternating IFN- α with BCG during maintenance therapy was less effective than BCG alone at preventing recurrence with 80% and 45% experiencing a recurrence within 15 years, respectively [50]. Another study showed that BCG was more effective than IFN- α + epirubicin with recurrence-free survival rates of 38% and 59% at a follow-up time of 5 years [51]. A retrospective study found that patients given a salvage therapy of IFN- α + BCG

had reduced recurrence-free survival when compared with those patients treated with BCG alone [52].

Only a single study has directly compared IFN- α plus BCG against BCG alone in a controlled, randomized clinical trial [53]. This study found no significant difference between treatment groups in the percentage of patients that were tumor free after 2 years. However, preclinical and clinical data have shown that IFN- α can enhance the therapeutic efficacy of reduced doses of BCG, a strategy that could potentially limit the toxic side effects seen in many patients treated with BCG [54,55].

Several studies have explored the use of BCG which has been genetically modified to express IFN- α or T_H1 cytokines [56–58]. These studies have shown an enhanced ability to stimulate immune cells *in vitro* against human bladder cancer cell lines, but cytokine-expressing BCG has not progressed to clinical trials. A related strategy is the direct transfection of the tumor or the urothelium via plasmids or viruses. While there have been numerous preclinical efforts using this strategy, clinical efforts have thus far been limited to a single Phase I trial (NCT01162785) in which an adenovirus encoding IFN- α was delivered intravesically [57]. IFN- α was produced for up to 5 days after administration and was well tolerated. A Phase II trial (NCT01687244) is ongoing to further investigate the efficacy of this strategy.

Overall, IFN- α has been the most common cytokine used as either a monotherapy or a combination therapy with BCG. Despite the number of investigations, IFN- α has shown limited effectiveness. While IFN- α exhibits potent cytostatic/cytotoxic activity and can enhance tumor immunogenicity, it may not be able to effectively engage the adaptive immune system.

GM-CSF

GM-CSF is a hematopoietic growth factor and immune modulator that is secreted by a range of cells including endothelial cells, macrophages, T-cells and some tumor cells. As implied by its name, GM-CSF stimulates the production of granulocytes and monocytes from hematopoietic progenitors. In addition, GM-CSF activates macrophages and promotes dendritic cell development. GM-CSF has been widely used in cancer vaccine and immunotherapy approaches due to its ability to enhance dendritic cell function.

Preclinically in bladder cancer, the delivery of recombinant GM-CSF to the bladder tumor environment can be enhanced via streptavidin and biotin-based targeting [59–62]. For example, 1-day old orthotopic MB49 bladder tumors treated intravesically first with biotin for 30 min followed by streptavidin tagged GMCSF (SA-GMCSF) for 1 h resulted in elimination

of 6/16 tumors versus 1/10 for recombinant GM-CSF alone [59]. These results suggest that maintenance of GM-CSF at the tumor site increases its efficacy. In subsequent studies, ethanol fixed MB49 cells were labeled with SA-GMCSF *in vitro* before being applied as both therapeutic and preventative vaccines against s.c. and lung metastasized MB49 [60,62]. Mice treated with the vaccine showed extended survival while cured mice in each of these studies demonstrated tumor-specific immunity.

In recent clinical studies, CG0070, an oncolytic adenovirus virus expressing GM-CSF and programmed to replicate only in cells with retinoblastoma (Rb) pathway defects was evaluated in a Phase I trial for NMIBC (NCT00109655) [63,64]. Patients enrolled in this study had all experienced at least one failed treatment with BCG. Different doses and treatment schedules were tested intravesically on a total of 35 patients with no reports of severe adverse events. Response rates were promising with 17/35 (48.6%) of patients exhibiting a complete response. CG0070 is currently being evaluated in a Phase II trial (NCT01438112).

It should be noted that GM-CSF can be a double-edged sword. On the one hand, GM-CSF drives the proliferation of leukocytes and the development of antigen presenting cells which are needed for a robust adaptive response. On the other hand, GM-CSF has been shown to expand immature myeloid subsets, such as myeloid derived suppressor cells, that can act as immune suppressors. Knowledge of proper dosing and delivery strategies is needed to drive appropriate responses [65].

IL-2

IL-2 induces the proliferation and activation of lymphocytes and is approved for the treatment of metastatic renal cell carcinoma and metastatic melanoma. Unfortunately, IL-2 also promotes the growth of tumor-supporting Tregs. The history of IL-2 in the treatment of bladder cancer is quite long, stretching back to intravesical treatments of NMIBC in 1984 [66]. However, despite promising preclinical data, results from clinical studies have been mixed.

Clinical study of IL-2 immunotherapy for bladder cancer in the last 5 years has been limited. Most studies have been small with less than 30 participants and have not compared IL-2 against alternate therapeutics. In a recent pilot study, 36 patients with NMIBC were treated with intravesical IL-2 immunotherapy after complete or intentionally incomplete TURBT to assess the importance of residual marker lesions [67]. Marker lesions are a potentially important source of tumor-associated antigens. Intravesical IL-2 immunotherapy was found to be more effective at preventing recurrence

when a marker lesion was left behind than complete TURBT with median recurrence-free survival times of 20 and 7 months, respectively [67]. One other clinical-stage investigation with IL-2 involves a fusion protein, ALT-801, composed of IL-2-linked to a p53-specific T-cell receptor. ALT-801 has shown effectiveness in other tumors and is currently being investigated in clinical trials against both MIBC (NCT01326871) and NMIBC (NCT01625260).

Preclinically, a few studies have investigated the utility of MB49 bladder cancer cells treated with biotin and streptavidin-IL-2 as a vaccine. In one study, vaccinated mice exhibited both therapeutic and preventive immune responses against MB49 lung metastases [68]. A similar study showed that sequential vaccination with GM-CSF-bound MB49 cells followed by IL-2-bound MB49 cells was more effective than either alone at prolonging survival and inducing memory [61]. Another study investigated the ability of streptavidin-IL-2 to bind to biotin-pretreated orthotopic MB49 tumors in mice. This delivery approach resulted in prolonged persistence of IL-2 on the bladder surface and led to enhanced survival over IL-2 alone. Nine of 25 mice with 2-day old tumors treated with the biotin/streptavidin-IL-2 approach survived longer than 60 days. Perhaps more importantly, 5/9 long-term survivors rejected an intravesical rechallenge with MB49 cells [69].

Another study investigated IL-2, IFN- α and gemcitabine as single agents and in combination with BCG against orthotopic AY-27 tumors in rats [70]. BCG was given at low (5×10^5 cfu/ml), medium (5×10^6 cfu/ml) and high (5×10^7 cfu/ml) doses. Recombinant IL-2 (5×10^5 units) was combined with low-dose BCG. Two out of 10, 4/10, 5/10 and 5/10 rats were alive at day 90 with 0, 3, 4 and 3 being completely tumor-free in the low, medium, high and combination treatment group, respectively. IFN α and gemcitabine combined with BCG showed no additional survival benefit over any of the agents alone. As single agents, IFN- α and gemcitabine were as effective as BCG.

Overall, IL-2 remains a promising candidate for bladder cancer immunotherapy due to its ability to induce adaptive responses and protective immunity. As with IL-2-based immunotherapies in other cancers, immune-enhancing benefits must be weighed against the potential for amplification of tumor-supporting Tregs. To assess the full potential of IL-2-based immunotherapy for bladder cancer, additional clinical studies are needed to either combine or compare IL-2 and BCG

IL-12

IL-12 is a proinflammatory T_H1 polarizing cytokine produced primarily by antigen presenting cells

in response to danger signals. IL-12 activates NK cells, promotes proliferation of cytotoxic T-cells and increases the production of IFN- γ , a key downstream effector molecule. IFN- γ has direct cytotoxic and/or cytostatic activity on a range of cells, including cancer cells. In addition, IFN- γ promotes antitumor responses indirectly through antiangiogenic responses as well as by increasing MHC expression on tumor cells leading to enhanced immune recognition [71]. In several older preclinical studies, intravesical administration of IL-12 has shown the ability to eradicate lesions and generate local immunity when delivered against orthotopic MBT-2 tumors [72,73].

In the clinic, only one trial has investigated intravesical IL-12 immunotherapy in 15 patients with recurrent superficial bladder cancer [74]. Intravesical administration of IL-12 was well tolerated, however, a maximum tolerated dose was not found and no responses were recorded. A likely reason for the lack of response was the lack of an effective delivery strategy. The urothelium is the least permeable epithelial barrier in the body and IL-12 is a large (70 kDa) protein. It is likely that IL-12 was not able to penetrate into the tumor and surrounding urothelium to reach resident immune cells. Our group has hypothesized that enhancing the delivery of IL-12 will improve its antitumor efficacy.

Our experiments have focused on using chitosan as an intravesical delivery vehicle. Chitosan is a large polymer that forms a viscous and mucoadhesive solution capable of transiently opening the tight junctions of the urothelium. This allows for increased and sustained penetration of IL-12 into the wall of the bladder. In addition, the mucoadhesive feature of chitosan allows it to stick to bladder surface mucin. We have shown that recombinant IL-12 (5 μ g) alone can eliminate 50% of established orthotopic MB49 tumors. However, when IL-12 was co-formulated with chitosan solution (chitosan/IL-12) the proportion of complete responders was increased to 88–100% [75]. Of particular interest to the bladder cancer community, 100% of cured mice rejected a subsequent intravesical tumor rechallenge for a least 18 months after the primary treatment, suggesting a local memory response.

A follow-up study investigated chitosan/IL-12, at reduced doses, against established MB49 and MBT-2 tumors [76]. Chitosan/IL-12 was able to eradicate 100% of MB49 tumors at a dose of 1 μ g IL-12 and 50% of MBT-2 tumors at a dose of 2.5 μ g IL-12. In this instance cured mice rejected both local (intravesical) and distant s.c. tumor rechallenges, demonstrating for the first time that chitosan/IL-12 can induce a systemic tumor-specific immune response from a localized, intravesical therapy. Another seminal finding from this study was that intravesical chitosan/

IL-12 immunotherapy was capable of controlling a distant, s.c. tumor. This potentially clinically significant result suggests that intravesical chitosan/IL-12 can induce abscopal responses against metastatic lesions. Therefore, intravesical chitosan/IL-12 could represent a bladder-sparing alternative for MIBC.

Other groups have also demonstrated the potential of IL-12 in preclinical studies. One study showed that IL-12 produced in a gamma-retrovirus targeted to HER2 was able to inhibit tumor growth and enhance survival for 2/12 mice when given intratumorally against s.c. MBT-2 [77]. Another study showed that IL-12 has antitumor properties even in the absence of T cells. Nude mice were given human EJ bladder cancer cells subcutaneously and then treated with chemotherapy, IL-12 plasmids, or a combination [78]. The combination therapy was the most effective at decreasing the tumor size, but was unable to eliminate the tumors.

Despite strong preclinical evidence for IL-12's ability to induce a sustained, adaptive antitumor response, there do not appear to be any IL-12-based clinical trials for bladder cancer on the horizon. It is likely that toxicity concerns remaining after two widely publicized IL-12-related deaths in a Phase II advanced renal cell carcinoma study nearly two decades ago have somewhat curbed enthusiasm for IL-12. However, IL-12's toxicity is associated with frequent systemic exposure to the cytokine. A recent focus on the development of controlled, localized delivery strategies capable of limiting systemic exposure promises to reduce toxicity while harnessing the antitumor potential of this powerful cytokine.

IL-15

IL-15 is a T-cell growth factor that shares several features with IL-2. However, among several key differences, IL-15 has been shown to specifically increase the proliferation and maintenance of CD8⁺ memory T-cells. In bladder cancer, IL-15 has been investigated in two preclinical studies. Matsumoto *et al.* delivered six intravesical doses of liposomes containing IL-15 plasmids in mice bearing established orthotopic MBT-2 tumors [79]. Treated mice experienced reduced tumor burdens, increased CD8⁺ T-cell infiltration and enhanced survival with 8/20 mice surviving long term. Surviving mice also showed the ability to delay, but not reject, the growth of a s.c. tumor rechallenge. These data suggest that IL-15, like IL-2 and IL-12, is capable of inducing a systemic antitumor immune response.

Another group investigated the efficacy of a mutated IL-15/IL-15R α -FC fusion protein (termed ALT-803) in conjunction with BCG against chemically induced orthotopic tumors in rats [80]. Treatment with ALT-

803 plus BCG resulted in reduced bladder weights, increased levels of NK cells and enhanced infiltration of CD3⁺ cells but not CD8⁺ or CD4⁺ cells when compared with BCG alone. IL-1 α and IL-1 β were the primary effector cytokines at the systemic level while RANTES was the primary local cytokine following treatment with ALT-803 plus BCG. Memory responses were not evaluated in this model, but ALT-803 had previously generated a memory response against melanoma in mice. A clinical study inspired by this data is currently recruiting NMIBC patients (NCT02138734).

In sum, IL-15-based immunotherapies are promising due to their ability to induce tumor-specific adaptive immune responses. Like IL-2 and IL-12, in order for IL-15 to reach its full potential, a well-considered delivery strategy is needed. In addition, combinations and dosing schedules with other therapeutics should also be explored. In particular, IL-15 may be most effective in sustaining or augmenting a memory response initiated by some other inductive immunotherapy.

Antibodies

Checkpoint inhibitors

Anti-CTLA-4 antibody (α CTLA-4) therapy, which is approved for the treatment of unresectable or metastatic melanoma, has been investigated in a 2010 clinical trial of 12 patients with localized bladder cancer prior to cystectomy [81]. The primary goal of this study was to identify biomarkers associated with a positive response to α CTLA-4 (ipilimumab). Patients received two doses, 3 mg/kg or 10 mg/kg, of ipilimumab 3 weeks apart with cystectomy 4 weeks after the second dose. A sustained increase in the level of CD4⁺ICOS^{high} lymphocytes was found to be associated with enhanced survival [81]. A Phase II clinical trial evaluating ipilimumab in patients with metastatic bladder cancer is underway and expected to be completed by June 2016 [24]. Ipilimumab is also under investigation in combination with the anti-PD-1 antibody (α PD-1), nivolumab, in three clinical studies.

Anti-PD-L1 (α PD-L1) therapies are under clinical investigation for a number of cancer indications and have demonstrated some recent successes in bladder cancer [24]. MPDL3280A was evaluated in 67 patients with metastatic bladder cancer in a Phase I trial [82]. Patients were scored for PD-L1 expression on both infiltrating immune cells and tumors. Treatment with intravenous MPDL3280A led to an objective response rate of 43% in tumors with high levels of PD-L1 expression but only 11% for low expressing tumors, indicating that PD-L1 expression can be used as a predictive biomarker for the success of PD-L1-based therapies. Based on these results, MPDL3280A

received breakthrough status by the US FDA and is currently under investigation in multiple clinical trials. Regarding anti-PD-1 immunotherapy, at the time of this review, four clinical trials were recruiting bladder cancer patients to evaluate the safety and activity of nivolumab with an additional nine clinical trials exploring pembrolizumab.

Because of the clinical success of checkpoint inhibitors in other cancers, clinical studies have outpaced preclinical studies in bladder cancer. In the only preclinical study to date, Mangsbo *et al.* administered anti-CTLA-4 or anti-PD-1 antibodies i.p. in combination with peritumoral CpG or BCG in mice bearing day 7 or 8 s.c. MB49 tumors [36]. CpG plus α CTLA-4, α PD-1 or α PD-L1 induced complete regression of large tumors in 6/7, 5/7 and 2/7 mice, respectively. CpG plus check point inhibition (13/21 mice became tumor-free) was superior to BCG plus checkpoint inhibition (4/18), CpG alone (1/7), or checkpoint inhibition alone (10/33). CTLA-4 blockade was superior to PD-1 blockade and the combination was not better than α CTLA-4 alone. Mechanistically, CpG plus α CTLA-4 or α PD-1 increased levels of tumor-reactive T cells and reduced numbers of Tregs at the tumor site.

While the future of checkpoint inhibition in the treatment of bladder cancer appears promising, there is much work to be done at the preclinical level to delineate antitumor mechanisms and explore additional treatment paradigms including novel combinations. It is not clear, from the literature in bladder cancer or any cancer, if checkpoint inhibition alone is sufficient to induce durable antitumor immunity or if combination with other inductive treatments, such as CpG, radiotherapy or cytokine immunotherapy, is needed for durable antitumor immunity. An additional area of exploration is local/intravesical delivery of checkpoint inhibitors. Checkpoint inhibitors cause severe immune-related adverse events in a significant minority of patients. Similar to our IL-12 studies, intravesical delivery of checkpoint inhibitors could minimize the adverse events associated with high systemic concentrations while maintaining high local concentrations at the tumor site.

Agonistic antibodies

CD40 is a costimulatory molecule found on antigen presenting and phagocytic cells, such as macrophages and dendritic cells. The binding of CD40 to CD40L located on T_H cells results in activation of, as well as enhanced phagocytosis and cytokine production by, antigen presenting cells. CD40 is also present on some tumors where anti-CD40 antibodies can have a direct antitumor effect by inducing apoptosis or antibody

dependent cytotoxicity. For these reasons, anti-CD40 antibodies have been investigated as a potential immunotherapeutic both clinically and preclinically, but its use against bladder cancer has been limited.

Mangso *et al.* showed that peritumoral administration of anti-CD40 antibodies was able to eradicate established s.c. MB49 tumors as effectively as systemic (i.p.) administration [83]. Local delivery also reduced systemic inflammation as measured by haptoglobin levels. Importantly, local delivery of anti-CD40 antibodies was able to control distant tumors while cured mice were protected from rechallenge in a tumor-specific manner. These data demonstrate that local anti-CD40 therapy is capable of inducing systemic, adaptive immunity.

A follow-up study using fully humanized anti-CD40 (ADC-1013) showed that locally administered ADC-1013 initiated an antitumor response against human s.c. EJ bladder tumors in immunodeficient mice whose immune systems were reconstituted with human dendritic cells and T-cells [52]. This study also demonstrated that ADC-1013 given peritumorally in one MB49 tumor was able to control a second distant MB49 tumor. A tumor-specific memory response mediated by CD8⁺ and CD4⁺ T cells was capable of rejecting MB49 rechallenge up to 5 months after treatment. A Phase I trial to assess the safety of ADC-1013 is currently recruiting patients with advanced solid tumors, including bladder tumors (NCT02379741).

While investigations into agonistic antibodies for the treatment of bladder tumors are quite new, preclinical results are promising. The demonstration of a sustained memory as well as abscopal responses via local delivery is particularly encouraging. More preclinical and clinical investigations of agonistic antibodies are warranted to assess their effectiveness in intravesical delivery as single agents and in combination with other therapies.

Anti-IL-10

IL-10 is an anti-inflammatory cytokine associated with Treg infiltration and poor tumor prognosis in bladder cancer [13]. Several preclinical studies have explored the consequences of blocking IL-10 or its receptor in the context of bladder cancer [84–86]. Blocking IL-10 function via systemically administered antibodies against the IL-10 receptor increased the IFN γ :IL-10 ratio, enhanced the efficacy of BCG against 1-day old lesions, prevented lung metastases and established systemic antitumor immunity as measured via tumor-specific cytotoxic T lymphocyte activity. Overall, the strategy of blocking the effects of IL-10 is promising especially when paired with an inductive treatment such as BCG, TLR agonists, or inflammatory cytokines.

Future perspective

After decades of research and unfulfilled promises, the era of cancer immunotherapy has finally arrived. Within the last 6 years, there have been a number of breakthrough approvals including the first therapeutic cancer vaccine, sipuleucel-T, three monoclonal antibodies targeting immune checkpoints, ipilimumab, nivolumab and pembrolizumab, and an oncolytic virus encoding GM-CSF, talimogene laherparepvec. In particular, the approvals of CTLA-4 (ipilimumab) and PD-1 (nivolumab and pembrolizumab) blockade for advanced melanoma and metastatic lung cancer have created substantial momentum. Currently, there are more than 470 clinical studies evaluating one or more of these checkpoint inhibitors, with and without additional therapies, in various cancers including bladder cancer. As mentioned previously, checkpoint blockade may induce stronger, more durable antitumor responses when combined with an inductive immunotherapy that can initiate a robust adaptive immune response.

With regard to bladder cancer, over the next 5–10 years, the expanding indications of CTLA-4 and PD-1 blockade are likely to include muscle invasive and/or metastatic bladder carcinomas. There are currently 12 clinical studies investigating CTLA-4 or PD-1 checkpoint inhibition in advanced, MIBC or high-risk bladder cancer. The 5-year survival rate for MIBC is 30 to 50% depending on tumor status and lymph node involvement. Given that advanced melanoma has similar survival rates and is similarly immunogenic as MIBC in terms of mutational load, it would not be surprising if clinical studies demonstrate similar clinical benefits. However, like in melanoma, checkpoint blockade is not expected to induce complete responses in the majority of patients with MIBC. As such, it does not appear likely that checkpoint blockade alone could prevent life altering cystectomy in a majority of patients. Perhaps combining checkpoint blockade with another inductive immunotherapy could be bladder sparing.

Another combination approach worthy of future consideration is chemotherapy plus immunotherapy. Bladder cancer is responsive to cytotoxic agents. Intravesical mitomycin C is routine for low grade NMIBC following TURBT while systemic cisplatin is standard for metastatic bladder cancer. In addition, a growing literature in diverse cancers demonstrates that cytotoxic agents can enhance tumor recognition by the immune system, in some cases revealing neoantigens. Furthermore, certain chemotherapy agents can eliminate or inhibit tumor-supporting immunosuppressive cells. Examples include cyclophosphamide for inhibition of Tregs and oxaliplatin for inhibition of regulatory B cells.

In addition to combination immunotherapies, there are two additional approaches for bladder cancer that should be considered. First, it may be more effective and safer to administer checkpoint inhibitors intravesically instead of intravenously. As described in this review, a number of preclinical studies have demonstrated that local immunomodulation of the bladder through intravesical delivery of cytokines and agonists can result in durable local and systemic adaptive immune responses. The intravesical route provides direct access to bladder tumors and a unique opportunity to reduce adverse events associated with systemic delivery of immunotherapy. Our group is currently investigating the efficacy of localized checkpoint blockade in preclinical models. Preliminary data indicate that α CTLA-4 exhibits greater orthotopic bladder tumor inhibition than α PD-1 when administered intravesically.

Secondly, although the vast majority of clinical studies investigating checkpoint blockade are in advanced cancers, it is expected that checkpoint blockade will be most effective against smaller tumor burdens such as NMIBC. As discussed below, overcoming BCG entrenchment is a significant challenge. However, combining checkpoint blockade with BCG is a logical intermediate step. Preclinical studies have shown that checkpoint blockade plus BCG is superior to BCG alone, however, combinations of checkpoint blockade with other immunotherapies are more effective.

Beyond the inevitable evaluation of checkpoint inhibition, there are a number of other immunotherapies, including IL-12, IL-15 and anti-CD40 that are deserving of clinical consideration in bladder cancer. Each of these immunotherapies, unlike BCG, have demonstrated the ability to induce tumor-specific immunity. Like most novel cancer treatments, development and clinical translation of these immunotherapies will likely require investment from the biopharmaceutical industry.

Finally, we speculate on the future of BCG immunotherapy in the treatment of high-risk NMIBC. While BCG has been touted by some as the most successful cancer immunotherapy to date, it is not without significant concerns. The immunological limitations of BCG were presented in a previous section and can be summarized by the dogma that BCG is incapable of inducing tumor-specific adaptive immunity. In order to significantly impact recurrence rates, we believe that a tumor-specific adaptive response is essential.

It has been frustrating to many, researchers, physicians and patients alike, that the use of intravesical BCG for high-risk or high-grade NMIBC is largely unchanged since Morales' study nearly four decades ago [29]. The lack of progress in the management of NMIBC is disconcerting and causes us to ask why have we not been able to find a better immunotherapy than BCG for bladder cancer? The answer to this question is likely multifactorial. First, funding for bladder cancer research is dismal. The National Cancer Institute

Executive summary

Bladder cancer overview

- The clinical management of bladder cancer has not changed significantly in several decades.
- Immunotherapy for bladder cancer is dominated by intravesical BCG for high-risk NMIBC.
- Bladder cancer, due to high recurrence rates, requires continuous and costly follow-up.

Immunologic challenges in bladder cancer

- The immunologic milieu of the bladder is uniquely suited to prevent chronic inflammation in the face of constant antigen exposure.
- Immunosuppressive strategies used by the normal bladder are likely obstacles to effective bladder cancer immunotherapy.

Limitations of BCG immunotherapy

- Intravesical BCG immunotherapy for NMIBC has remained unchanged for nearly four decades.
- BCG has not been shown to establish tumor-specific adaptive immunity.
- The exact mechanisms by which BCG exerts antitumor responses remain elusive.

Novel strategies for bladder cancer

- IFN- α has been widely explored, however, preclinical and clinical results show limited effectiveness.
- Several promising immunotherapies, including IL-12, IL-15 and anti-CD40, have demonstrated the ability to induce tumor-specific immunity.
- Checkpoint blockade has skipped preclinical evaluation and is the subject of intense clinical investigation.

Future perspective

- Checkpoint blockade will continue to be evaluated, with and without additional therapies, in advanced bladder cancer.
- Local delivery of checkpoint inhibitors may be as effective as systemic delivery but with less toxicity.
- The dominant stance held by BCG will have to be softened in order for other immunotherapies to breakthrough for the treatment of NMIBC.

(NCI) invests only about \$20M per year in bladder cancer research [87]. This translates to about \$280 per newly diagnosed bladder cancer patient. For comparison, breast cancer averages about \$2700 in NCI-sponsored research support per newly diagnosed patient. Not to mention, breast cancer receives considerable support from a number of high profile foundations. Second, bladder cancer clinical studies are notoriously difficult to accrue to and comparative studies require large numbers of patients. Third, while BCG is not perfect, there may be a sense that it is 'good enough', and that new therapies are too risky. Such a conservative and risk-averse outlook discourages innovation. Furthermore, clinical studies investigating novel immunotherapies are typically limited to enrolling BCG refractory patients. However, the oft-cited mantra that bladder cancer treatment has not changed in 40 years may be encouraging change. Over the next 5 years, we expect the bladder cancer community's appe-

tite for novel immunotherapies capable of replacing BCG will grow. We expect to see a greater tolerance for clinical studies in BCG naïve patients where immunotherapies have a greater chance for both accrual and better clinical outcomes. Indeed, the next 5 years can be transformational and the hope is that by focusing on immunotherapies capable of establishing adaptive tumor-specific memory, we can significantly impact bladder cancer morbidity and mortality.

Financial & competing interests disclosure

The authors have no relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript. This includes employment, consultancies, honoraria, stock ownership or options, expert testimony, grants or patents received or pending, or royalties.

No writing assistance was utilized in the production of this manuscript.

References

Papers of special note have been highlighted as: • of interest; •• of considerable interest

- Howlader N, Noone AM, Krapcho M *et al.* SEER Cancer Statistics Review (2015). http://seer.cancer.gov/csr/1975_2012/
- Siegel R, Miller K, Jemal A. Cancer statistics, 2015. *CA Cancer J. Clin.* 65(1), 5–29 (2015).
- Kaufman D, Shipley W, Feldman A. Bladder cancer. *Lancet* 374(9685), 239–249 (2009).
- Kresowik T, Griffith T. Bacillus Calmette–Guerin immunotherapy for urothelial carcinoma of the bladder. *Immunotherapy* 1(2), 281–288 (2009).
- Bostrom P, Rhijn B van, Fleshner N *et al.* Staging and staging errors in bladder cancer. *Eur. Urol. Suppl.* 9(1), 29 (2010).
- Herr H, Dotan Z, Donat M, Bajorin D. Defining optimal therapy for muscle invasive bladder cancer. *J. Urol.* 177(2), 437443 (2007).
- Chamie K, Saigal CS, Lai J *et al.* Compliance with guidelines for patients with bladder cancer: variation in the delivery of care. *Cancer* 117(23), 5392–5401 (2011).
- Chamie K, Litwin M, Bassett J *et al.* Recurrence of high-risk bladder cancer: a population-based analysis. *Cancer* 119(17), 3219–3227 (2013).
- Sievert, Amend, Nagele, *et al.* Economic aspects of bladder cancer: what are the benefits and costs? *World J. Urol.* 27(3), 295–300 (2009).
- Kandoth C, McLellan MD, Vandin F *et al.* Mutational landscape and significance across 12 major cancer types. *Nature* 502(7471), 333–339 (2013).
- Ingersoll MA, Albert ML. From infection to immunotherapy: host immune responses to bacteria at the bladder mucosa. *Mucosal Immunol.* 6(6), 1041–1053 (2013).
- Excellent review describing the oft-overlooked immunology of the bladder.
- deLeeuw R, Kost S, Kakal J, Nelson B. The prognostic value of FoxP3+ tumor-infiltrating lymphocytes in cancer: a critical review of the literature. *Clin. Cancer Res.* 18(11), 3022–3029 (2012).
- Loskog A, Ninalga C, Paul-Wetterberg G, Torre M de la, Malmström P-UU, Tötterman TH. Human bladder carcinoma is dominated by T-regulatory cells and Th1 inhibitory cytokines. *J. Urol.* 177(1), 353–358 (2007).
- Horn T, Laus J, Seitz AK *et al.* The prognostic effect of tumour-infiltrating lymphocytic subpopulations in bladder cancer. *World J. Urol.* (2015) (Epub ahead of print).
- Winerdal M, Marits P, Winerdal M *et al.* FOXP3 and survival in urinary bladder cancer. *BJU Int.* 108(10), 1672–1678 (2011).
- Liakou CI, Narayanan S, Ng Tang D, Logothetis CJ, Sharma P. Focus on TILs: prognostic significance of tumor infiltrating lymphocytes in human bladder cancer. *Cancer Immun.* 7, 10 (2007).
- Baksh K, Weber J. Immune checkpoint protein inhibition for cancer: preclinical justification for CTLA-4 and PD-1 blockade and new combinations. *Semin. Oncol.* 42(3), 363–367 (2015).
- Boorjian S, Sheinin Y, Crispen P *et al.* T-cell coregulatory molecule expression in urothelial cell carcinoma: clinicopathologic correlations and association with survival. *Clin. Cancer Res.* 14(15), 4800–4808 (2008).
- Xylinas E, Robinson BD, Kluth LA *et al.* Association of T-cell co-regulatory protein expression with clinical outcomes following radical cystectomy for urothelial carcinoma of the bladder. *Eur. J. Surg. Oncol.* 40(1), 121–127 (2014).
- Inman BA, Sebo TJ, Frigola X *et al.* PD-L1 (B7-H1) expression by urothelial carcinoma of the bladder and

- BCG-induced granulomata: associations with localized stage progression. *Cancer* 109(8), 1499–1505 (2007).
- 21 Nakanishi J, Wada Y, Matsumoto K, Azuma M, Kikuchi K, Ueda S. Overexpression of B7-H1 (PD-L1) significantly associates with tumor grade and postoperative prognosis in human urothelial cancers. *Cancer Immunol. Immunother.* 56(8), 1173–1182 (2007).
- 22 Jaiswal PK, Singh V, Mittal RD. Cytotoxic T lymphocyte antigen 4 (CTLA4) gene polymorphism with bladder cancer risk in North Indian population. *Mol. Biol. Rep.* 41(2), 799–807 (2014).
- 23 Wang L, Su G, Zhao X *et al.* Association between the cytotoxic T-lymphocyte antigen 4 +49A/G polymorphism and bladder cancer risk. *Tumor Biol.* 35(2), 1139–1142 (2013).
- 24 Carosella E, Ploussard G, LeMaout J, Desgrandchamps F. A systematic review of immunotherapy in urologic cancer: evolving roles for targeting of CTLA-4, PD-1/PD-L1, and HLA-G. *Eur. Urol.* 68(2), 267–279 (2015).
- 25 Seliger B. Novel insights into the molecular mechanisms of HLA class I abnormalities. *Cancer Immunol. Immunother.* 61(2), 249–254 (2011).
- 26 Garrido F, Cabrera T, Aptsiauri N. “Hard” and “soft” lesions underlying the HLA class I alterations in cancer cells: implications for immunotherapy. *Int. J. Cancer* 127(2), 249–256 (2010).
- 27 Carretero R, Cabrera T, Gil H *et al.* Bacillus Calmette–Guerin immunotherapy of bladder cancer induces selection of human leukocyte antigen class I-deficient tumor cells. *Int. J. Cancer* 129(4), 839–846 (2011).
- 28 Romero JM, Jiménez P, Cabrera T *et al.* Coordinated downregulation of the antigen presentation machinery and HLA class I/beta2-microglobulin complex is responsible for HLA-ABC loss in bladder cancer. *Int. J. Cancer* 113(4), 605–610 (2005).
- 29 Morales A, Eidinger D, Bruce AW. Intracavitary bacillus Calmette–Guerin in the treatment of superficial bladder tumors. *J. Urol.* 116(2), 180–183 (1976).
- 30 Redelman-Sidi G, Glickman M, Bochner B. The mechanism of action of BCG therapy for bladder cancer – a current perspective. *Nat. Rev. Urol.* 11(3), 153–162 (2014).
- **Provides an excellent description of the current understanding of BCG mechanisms.**
- 31 Askeland EJ, Newton MR, O'Donnell MA, Luo Y. Bladder cancer immunotherapy: BCG and beyond. *Adv. Urol.* 2012 181987 (2012).
- 32 Alexandroff AB, Jackson AM, O'Donnell MA, James K. BCG immunotherapy of bladder cancer: 20 years on. *Lancet* 353(9165), 1689–1694 (1999).
- 33 Smaldone M, Casella D, Welchons D, Gingrich J. Investigational therapies for non-muscle invasive bladder cancer. *Expert Opin. Investig. Drugs* 19(3), 371–383 (2010).
- 34 Goutagny N, Estornes Y, Hasan U, Lebecque S, Caux C. Targeting pattern recognition receptors in cancer immunotherapy. *Targ. Oncol.* 7(1), 29–54 (2012).
- 35 LaRue H, Ayari C, Bergeron A, Fradet Y. Toll-like receptors in urothelial cells – targets for cancer immunotherapy. *Nat. Rev. Urol.* 10(9), 537–545 (2013).
- 36 Mangsbo S, Sandin L, Anger K, Korman A, Loskog A, Tötterman T. Enhanced tumor eradication by combining CTLA-4 or PD-1 blockade with CpG therapy. *J. Immunother.* 33(3), 225–235 (2010).
- **One of the few preclinical investigations with checkpoint inhibitors, this study demonstrated a synergy with CpG and CTLA-4, PD-1 and PD-L1 inhibitors.**
- 37 Codolo G, Fassan M, Munari F *et al.* HP-NAP inhibits the growth of bladder cancer in mice by activating a cytotoxic Th1 response. *Cancer Immunol. Immunother.* 61(1), 31–40 (2012).
- 38 Arends TJ, Lammers RJ, Falke J *et al.* Pharmacokinetic, pharmacodynamic, and activity evaluation of TMX-101 in a multicenter Phase I study in patients with papillary non-muscle-invasive bladder cancer. *Clin. Genitourin Cancer* 13(3), 204.e2–209.e2 (2015).
- 39 Huang H-N, Rajanbabu V, Pan C-Y, Chan Y-L, Chen J-Y, Wu C-J. Enhanced control of bladder-associated tumors using shrimp anti-lipopolysaccharide factor (SALF) antimicrobial peptide as a cancer vaccine adjuvant in mice. *Marine Drugs* 13(5), 3241–3258 (2015).
- 40 Huang H-N, Rajanbabu V, Pan C-Y, Chan Y-L, Wu C-J, Chen J-Y. A cancer vaccine based on the marine antimicrobial peptide pardaxin (GE33) for control of bladder-associated tumors. *Biomaterials.* 34(38), 10151–10159 (2013).
- 41 Chen M-H, Li W-S, Lue Y-S *et al.* Clitocybe nuda activates dendritic cells and acts as a DNA vaccine adjuvant. *Evid. Based Complement. Alternat. Med.* 2013, 761454 (2013).
- 42 Lin C-C, Yu Y-L, Shih C-C *et al.* A novel adjuvant Ling Zhi-8 enhances the efficacy of DNA cancer vaccine by activating dendritic cells. *Cancer Immunol. Immunother.* 60(7), 1019–1027 (2011).
- 43 Wang S-T, Chang C-C, Yen M-C *et al.* RNA interference-mediated silencing of Foxo3 in antigen-presenting cells as a strategy for the enhancement of DNA vaccine potency. *Gene Ther.* 18(4), 372–383 (2010).
- 44 Obara W, Ohsawa R, Kanehira M *et al.* Cancer peptide vaccine therapy developed from oncoantigens identified through genome-wide expression profile analysis for bladder cancer. *Jpn J. Clin. Oncol.* 42(7), 591–600 (2012).
- 45 Morse MA, Bradley DA, Keler T *et al.* CDX-1307: a novel vaccine under study as treatment for muscle-invasive bladder cancer. *Expert Rev. Vaccines* 10(6), 733–742 (2011).
- 46 Dhodapkar MV, Sznol M, Zhao B *et al.* Induction of antigen-specific immunity with a vaccine targeting NY-ESO-1 to the dendritic cell receptor DEC-205. *Sci. Transl. Med.* 6(232), 232ra51 (2014).
- 47 Bekisz J, Schmeisser H, Hernandez J, Goldman N, Zoon K. Human interferons alpha, beta and omega. *Growth Factors* 22(4), 243–251 (2004).
- 48 Ferrantini M, Capone I, Belardelli F. Interferon-alpha and cancer: mechanisms of action and new perspectives of clinical use. *Biochimie* 89(6–7), 884–893 (2007).
- 49 Lamm D, Brausi M, O'Donnell MA, Witjes JA. Interferon alfa in the treatment paradigm for non-muscle-invasive bladder cancer. *Urol. Oncol.* 32(1), 35.e21–30.e21 (2014).

- 50 Järvinen R, Marttila T, Kaasinen E *et al.* Long-term outcome of patients with frequently recurrent non-muscle-invasive bladder carcinoma treated with one perioperative plus four weekly instillations of Mitomycin C followed by monthly bacillus Calmette–Guérin (BCG) or alternating BCG and interferon- α 2b instillations: prospective randomised Finnbladder-4 Study. *Eur. Urol.* 68(4), 611–617 (2015).
- 51 Hemdan T, Johansson R, Jahnson S, Hellström P, Tasdemir I, Malmström P. 5-year outcome of a randomized prospective study comparing bacillus Calmette–Guérin with epirubicin and interferon- α 2b in patients with T1 bladder cancer. *J. Urol.* 191(5), 1244–1249 (2014).
- 52 Prasad SM, Eyre S, Loughlin KR. Salvage combination intravesical immunotherapy with bacillus Calmette–Guérin and interferon- α 2B: impact on recurrence, progression, and survival. *Hosp. Pract. (1995)*. 41(4), 31–39 (2013).
- 53 Nepple KG, Lightfoot AJ, Rosevear HM, O'Donnell MA, Lamm DL. Bacillus Calmette–Guérin with or without interferon α -2b and megadose versus recommended daily allowance vitamins during induction and maintenance intravesical treatment of nonmuscle invasive bladder cancer. *J. Urol.* 184(5), 1915–1919 (2010).
- **The best clinical study regarding IFN- α , this paper showed that IFN- α plus BCG was no more effective than BCG alone.**
- 54 Chen X, O'Donnell MA, Luo Y. Dose-dependent synergy of Th1-stimulating cytokines on bacille Calmette–Guérin-induced interferon-gamma production by human mononuclear cells. *Clin. Exp. Immunol.* 149(1), 178–185 (2007).
- 55 Agarwal A, Agrawal U, Verma S, Mohanty NK, Saxena S. Serum Th1 and Th2 cytokine balance in patients of superficial transitional cell carcinoma of bladder pre- and post-intravesical combination immunotherapy. *Immunopharmacol. Immunotoxicol.* 32(2), 348–356 (2010).
- 56 Liu W, O'Donnell MA, Chen X, Han R, Luo Y. Recombinant bacillus Calmette–Guérin (BCG) expressing interferon-alpha 2B enhances human mononuclear cell cytotoxicity against bladder cancer cell lines *in vitro*. *Cancer Immunol. Immunother.* 58(10), 1647–1655 (2009).
- 57 Dinney CP, Fisher MB, Navai N *et al.* Phase I trial of intravesical recombinant adenovirus mediated interferon- α 2b formulated in Syn3 for bacillus Calmette–Guérin failures in nonmuscle invasive bladder cancer. *J. Urol.* 190(3), 850–856 (2013).
- 58 Luo Y, Henning J, O'Donnell MA. Th1 cytokine-secreting recombinant *Mycobacterium bovis* bacillus Calmette–Guérin and prospective use in immunotherapy of bladder cancer. *Clin. Dev. Immunol.* 2011 728930 (2011).
- 59 Hu Z, Tan W, Zhang L *et al.* A novel immunotherapy for superficial bladder cancer by intravesical immobilization of GM-CSF. *J. Cell. Mol. Med.* 14(6B), 1836–1844 (2010).
- 60 Zhu YT, Zhao Z, Fu XY *et al.* The granulocyte macrophage-colony stimulating factor surface modified MB49 bladder cancer stem cells vaccine against metastatic bladder cancer. *Stem Cell Res.* 13(1), 111–122 (2014).
- 61 Shi X, Zhang X, Li J *et al.* Sequential administration of GM-CSF and IL-2 surface-modified MB49 cells vaccines against the metastatic bladder cancer. *Urol. Oncol.* 31(6), 883–893 (2013).
- 62 Zhang X, Shi X, Li J *et al.* A novel therapeutic vaccine of mouse GM-CSF surface modified MB49 cells against metastatic bladder cancer. *J. Urol.* 187(3), 1071–1079 (2012).
- 63 Burke JM, Lamm DL, Meng MV *et al.* A first in human Phase 1 study of CG0070, a GM-CSF expressing oncolytic adenovirus, for the treatment of nonmuscle invasive bladder cancer. *J. Urol.* 188(6), 2391–2397 (2012).
- 64 Ramesh N, Ge Y, Ennist DL *et al.* CG0070, a conditionally replicating granulocyte macrophage colony-stimulating factor – armed oncolytic adenovirus for the treatment of bladder cancer. *Clin. Cancer Res.* 12(1), 305–313 (2006).
- 65 Clive KS, Tyler JA, Clifton GT *et al.* Use of GM-CSF as an adjuvant with cancer vaccines: beneficial or detrimental? *Expert Rev. Vaccines* 9(5), 519–525 (2010).
- 66 Pizza G, Severini G, Menniti D, De Vinci C, Corrado F. Tumour regression after intravesical injection of interleukin 2 (IL-2) in bladder cancer. Preliminary report. *Int. J. Cancer* 34(3), 359–367 (1984).
- 67 Den Otter W, Van Moorselaar RJ Van, Jacobs JJ *et al.* Role of marker lesion when applying intravesical instillations of IL-2 for non-muscle-invasive bladder cancer comparison of the therapeutic effects in two pilot studies. *Anticancer Res.* 33(5), 2099–2105 (2013).
- 68 Zhang X, Shi X, Li J *et al.* Novel immunotherapy for metastatic bladder cancer using vaccine of human interleukin-2 surface-modified MB 49 cells. *Urology.* 78(3), 722.e1–722.e6 (2011).
- 69 Huang X, Yu HS, Chen Z, Li JL, Hu ZM, Gao JM. A novel immunotherapy for superficial bladder cancer by the immobilization of streptavidin-tagged bioactive IL-2 on the biotinylated mucosal surface of the bladder wall. *Chin. J. Cancer* 29(6), 611–616 (2010).
- 70 Xiao Z, Hanel E, Mak A, Moore RB. Antitumor efficacy of intravesical BCG, Gemcitabine, Interferon- α and Interleukin-2 as mono- or combination-therapy for bladder cancer in an orthotopic tumor model. *Clin Med Insights Oncol.* 5, 315–323 (2011).
- 71 Trinchieri G. Interleukin-12 and the regulation of innate resistance and adaptive immunity. *Nat. Rev. Immunol.* 3(2), 133–146 (2003).
- 72 O'Donnell MA, Luo YI, Hunter SE, Chen X, Hayes LL, Clinton SK. Interleukin-12 immunotherapy of murine transitional cell carcinoma of the bladder: dose dependent tumor eradication and generation of protective immunity. *J. Urol.* 171(3), 1330–1335 (2004).
- 73 Horinaga M, Harsch K, Fukuyama R, Heston W, Larchian W. Intravesical interleukin-12 gene therapy in an orthotopic bladder cancer model. *Urology.* 66(2), 461–466 (2005).
- 74 Weiss GR, O'Donnell MA, Loughlin K, Zonno K, Laliberte RJ, Sherman ML. Phase 1 study of the intravesical administration of recombinant human interleukin-12 in patients with recurrent superficial transitional cell carcinoma of the bladder. *J. Immunother.* 26(4), 343–348 (2003).

- 75 Zaharoff DA, Hoffman BS, Hooper BH *et al.* Intravesical immunotherapy of superficial bladder cancer with Chitosan/ Interleukin-12. *Cancer Res.* 69(15), 6192-6199 (2009).
- 76 Smith SG, Koppolu BP, Ravindranathan S *et al.* Intravesical chitosan/interleukin-12 immunotherapy induces tumor-specific systemic immunity against murine bladder cancer. *Cancer Immunol. Immunother.* 64(6), 689–696 (2015).
- **First study to demonstrate robust systemic tumor-specific immunity from an intravesical IL-12-based immunotherapy.**
- 77 Tsai YS, Shiau AL, Chen YF, Tsai HT, Tzai TS, Wu CL. Enhancement of antitumor activity of gammaretrovirus carrying IL-12 gene through genetic modification of envelope targeting HER2 receptor: a promising strategy for bladder cancer therapy. *Cancer Gene Ther.* 17(1), 37–48 (2010).
- 78 Liu W, Cao Y, Fernández MI, Niu H, Xiu Y. Additive antitumoral effect of interleukin-12 gene therapy and chemotherapy in the treatment of urothelial bladder cancer *in vitro* and *in vivo*. *Int. Urol. Nephrol.* 43(3), 721–727 (2011).
- 79 Matsumoto K, Kikuchi E, Horinaga M *et al.* Intravesical interleukin-15 gene therapy in an orthotopic bladder cancer model. *Hum. Gene Ther.* 22(11), 1423–1432 (2011).
- 80 Gomes-Giacoaia E, Miyake M, Goodison S *et al.* Intravesical ALT-803 and BCG treatment reduces tumor burden in a carcinogen induced bladder cancer rat model; a role for cytokine production and NK cell expansion. *PLoS ONE.* 9(6), e96705 (2014).
- **An interesting study that demonstrated the immune effects of an IL-15 superagonist on NMIBC.**
- 81 Carthon BC, Wolchok JD, Yuan J *et al.* Preoperative CTLA-4 blockade: tolerability and immune monitoring in the setting of a presurgical clinical trial. *Clin. Cancer Res.* 16(10), 2861–2871 (2010).
- 82 Powles T, Eder JP, Fine GD *et al.* MPDL3280A (anti-PD-L1) treatment leads to clinical activity in metastatic bladder cancer. *Nature* 515(7528), 558–562 (2014).
- **Recent study showing that an anti-PD-L1 inhibitor can be effective against metastatic bladder cancer, paving the way for this therapy into other cancers.**
- 83 Sandin L, Orlova A, Gustafsson E *et al.* Locally delivered CD40 agonist antibody accumulates in secondary lymphoid organs and eradicates experimental disseminated bladder cancer. *Cancer Immunol Res.* 2(1), 80–90 (2013).
- **Investigated a murine agonistic antibody capable of inducing systemic immunity. A human version of this therapy is in clinical trials.**
- 84 Newton MR, Askeland EJ, Andresen ED *et al.* Anti-interleukin-10R1 monoclonal antibody in combination with bacillus Calmette–Guérin is protective against bladder cancer metastasis in a murine orthotopic tumour model and demonstrates systemic specific anti-tumour immunity. *Clin. Exp. Immunol.* 177(1), 261–268 (2014).
- 85 Bockholt NA, Knudson MJ, Henning JR *et al.* Anti-interleukin-10R1 monoclonal antibody enhances bacillus Calmette–Guérin induced T-helper type 1 immune responses and antitumor immunity in a mouse orthotopic model of bladder cancer. *J. Urol.* 187(6), 2228–2235 (2012).
- 86 Luo Y. Blocking IL-10 enhances bacillus Calmette–Guérin induced T helper type 1 immune responses and anti-bladder cancer immunity. *Oncoimmunology* 1(7), 1183–1185 (2012).
- 87 NCI. Research Funding by Cancer Type (2013). <http://fundedresearch.cancer.gov/nciportfolio/>