

LAB #:
PATIENT:
ID:
SEX:
DOB: AGE:

Microbiology Profile, stool

BACTERIOLOGY CULTURE

Expected/Beneficial flora	Commensal (Imbalanced) flora	Dysbiotic flora
4+ Bacteroides fragilis group 4+ Bifidobacterium spp. 4+ Escherichia coli 3+ Lactobacillus spp. NG Enterococcus spp.	3+ Alpha hemolytic strep 1+ Gamma hemolytic strep 4+ Mucoid Escherichia coli 1+ Staphylococcus aureus	
2+ Clostridium spp. NG = No Growth		

BACTERIA INFORMATION

Expected /Beneficial bacteria make up a significant portion of the total microflora in a healthy & balanced GI tract. These beneficial bacteria have many health-protecting effects in the GI tract including manufacturing vitamins, fermenting fibers, digesting proteins and carbohydrates, and propagating anti-tumor and anti-inflammatory factors.

Clostridia are prevalent flora in a healthy intestine. Clostridium spp. should be considered in the context of balance with other expected/beneficial flora. Absence of clostridia or over abundance relative to other expected/beneficial flora indicates bacterial imbalance. If *C. difficile* associated disease is suspected, a Comprehensive Clostridium culture or toxigenic *C. difficile* DNA test is recommended.

Commensal (Imbalanced) bacteria are usually neither pathogenic nor beneficial to the host GI tract. Imbalances can occur when there are insufficient levels of beneficial bacteria and increased levels of commensal bacteria. Certain commensal bacteria are reported as dysbiotic at higher levels.

Dysbiotic bacteria consist of known pathogenic bacteria and those that have the potential to cause disease in the GI tract. They can be present due to a number of factors including: consumption of contaminated water or food, exposure to chemicals that are toxic to beneficial bacteria; the use of antibiotics, oral contraceptives or other medications; poor fiber intake and high stress levels.

YEAST CULTURE

Normal flora	Dysbiotic flora
1+ Candida parapsilosis 1+ Rhodotorula mucilaginosa	2+ Saccharomyces cerevisiae/boulardii

MICROSCOPIC YEAST

Result: None	Expected: None - Rare
------------------------	---------------------------------

The microscopic finding of yeast in the stool is helpful in identifying whether there is proliferation of yeast. Rare yeast may be normal; however, yeast observed in higher amounts (few, moderate, or many) is abnormal.

YEAST INFORMATION

Yeast normally can be found in small quantities in the skin, mouth, intestine and mucocutaneous junctions. Overgrowth of yeast can infect virtually every organ system, leading to an extensive array of clinical manifestations. Fungal diarrhea is associated with broad-spectrum antibiotics or alterations of the patient's immune status. Symptoms may include abdominal pain, cramping and irritation. When investigating the presence of yeast, disparity may exist between culturing and microscopic examination. Yeast are not uniformly dispersed throughout the stool, this may lead to undetectable or low levels of yeast identified by microscopy, despite a cultured amount of yeast. Conversely, microscopic examination may reveal a significant amount of yeast present, but no yeast cultured. Yeast does not always survive transit through the intestines rendering it unviable.

Comments:

Date Collected:
Date Received:
Date Completed:


* *Aeromonas, Campylobacter, Plesiomonas, Salmonella, Shigella, Vibrio, Yersinia, & Edwardsiella tarda* have been specifically tested for and found absent unless reported.



LAB #:
 PATIENT:
 ID:
 SEX:
 AGE:

Yeast Susceptibilities: *Rhodotorula mucilaginosa*

NATURAL ANTIFUNGALS



Natural antifungal agents may be useful for treatment of patients when organisms display in-vitro sensitivity to these agents. The test is performed by using standardized techniques and filter paper disks impregnated with the listed agent. Relative activity is reported for each natural agent based upon the diameter of the zone of inhibition or no growth zone surrounding the disk. Data based on over 5000 individual observations were used to relate the zone size to the activity level of the agent. A scale of relative activity is defined for the natural agents tested.

NON-ABSORBED ANTIFUNGALS



Non-absorbed antifungals may be useful for treatment of patients when organisms display in-vitro sensitivity to these agents. The test is performed using standardized commercially prepared disks impregnated with Nystatin. Relative activity is reported based upon the diameter of the zone of inhibition or no growth zone surrounding the disk.

SAMPLE REPORT


Comments:
 Date Collected:
 Date Received:
 Date Completed:

Yeast antifungal susceptibility testing is intended for research use only.
 Not for use in diagnostic procedures.

LAB #:
 PATIENT:
 ID:
 SEX:
 AGE:

Yeast Susceptibilities: *Candida parapsilosis*

NATURAL ANTIFUNGALS




Natural antifungal agents may be useful for treatment of patients when organisms display in-vitro sensitivity to these agents. The test is performed by using standardized techniques and filter paper disks impregnated with the listed agent. Relative activity is reported for each natural agent based upon the diameter of the zone of inhibition or no growth zone surrounding the disk. Data based on over 5000 individual observations were used to relate the zone size to the activity level of the agent. A scale of relative activity is defined for the natural agents tested.

NON-ABSORBED ANTIFUNGALS



Non-absorbed antifungals may be useful for treatment of patients when organisms display in-vitro sensitivity to these agents. The test is performed using standardized commercially prepared disks impregnated with Nystatin. Relative activity is reported based upon the diameter of the zone of inhibition or no growth zone surrounding the disk.

AZOLE ANTIFUNGALS



Susceptible results imply that an infection due to the fungus may be appropriately treated when the recommended dosage of the tested antifungal agent is used.
Susceptible - Dose Dependent (S-DD) results imply that an infection due to the fungus may be treated when the highest recommended dosage of the tested antifungal agent is used.
Resistant results imply that the fungus will not be inhibited by normal dosage levels of the tested antifungal agent.

Standardized test interpretive categories established for *Candida* spp. are used for all yeast isolates.


Comments:
 Date Collected:
 Date Received:
 Date Completed:

Yeast antifungal susceptibility testing is intended for research use only.
 Not for use in diagnostic procedures.

LAB #:
 PATIENT:
 ID:
 SEX:
 AGE:

Yeast Susceptibilities: *Saccharomyces cerevisiae/boulardii*

NATURAL ANTIFUNGALS




Natural antifungal agents may be useful for treatment of patients when organisms display in-vitro sensitivity to these agents. The test is performed by using standardized techniques and filter paper disks impregnated with the listed agent. Relative activity is reported for each natural agent based upon the diameter of the zone of inhibition or no growth zone surrounding the disk. Data based on over 5000 individual observations were used to relate the zone size to the activity level of the agent. A scale of relative activity is defined for the natural agents tested.

NON-ABSORBED ANTIFUNGALS



Non-absorbed antifungals may be useful for treatment of patients when organisms display in-vitro sensitivity to these agents. The test is performed using standardized commercially prepared disks impregnated with Nystatin. Relative activity is reported based upon the diameter of the zone of inhibition or no growth zone surrounding the disk.

AZOLE ANTIFUNGALS



Susceptible results imply that an infection due to the fungus may be appropriately treated when the recommended dosage of the tested antifungal agent is used.
Susceptible - Dose Dependent (S-DD) results imply that an infection due to the fungus may be treated when the highest recommended dosage of the tested antifungal agent is used.
Resistant results imply that the fungus will not be inhibited by normal dosage levels of the tested antifungal agent.

Standardized test interpretive categories established for *Candida* spp. are used for all yeast isolates.

Comments:
 Date Collected:
 Date Received:
 Date Completed:

Yeast antifungal susceptibility testing is intended for research use only.
 Not for use in diagnostic procedures.

INTRODUCTION

This analysis of the stool specimen provides fundamental information about the overall gastrointestinal health of the patient. When abnormal microflora or significant aberrations in intestinal health markers are detected, specific interpretive paragraphs are presented. If no significant abnormalities are found, interpretive paragraphs are not presented.

Clostridium spp

Clostridia are expected inhabitants of the human intestine. Although most clostridia in the intestine are not virulent, certain species have been associated with disease. Clostridium perfringens is a major cause of food poisoning and is also one cause of antibiotic-associated diarrhea. Clostridium difficile is a causative agent in antibiotic-associated diarrhea and pseudomembranous colitis. Other species reported to be prevalent in high amounts in patients with Autistic Spectrum Disorder include Clostridium histolyticum group, Clostridium cluster I, Clostridium bolteae, and Clostridium tetani.

If these disease associations are a concern further testing may be necessary.

Washington W, Allen S, Janda W, Koneman E, Procop G, Schreckenberger P, Woods, G. Koneman's Color Atlas and Textbook of Diagnostic Microbiology, 6th edition. Lippincott Williams and Wilkins; 2006. pg 931-939

Song Y, Liu C, Finegold SM. Real-Time PCR Quantitation of Clostridia in Feces of Autistic Children. Applied and Environmental Microbiology. Nov. 2004, 6459-6465.

Parracho H, Bingham MO, Gibson GR, McCartney AL. Differences Between the Gut Microflora of Children with Autistic Spectrum Disorders and That of Healthy Children. Journal of Medical Microbiology. 2005;54, 987-991.

Imbalanced flora

Imbalanced flora are those bacteria that reside in the host gastrointestinal tract and neither injure nor benefit the host. Certain dysbiotic bacteria may appear under the imbalances category if found at low levels because they are not likely pathogenic at the levels detected. When imbalanced flora appear, it is not uncommon to find inadequate levels of one or more of the beneficial bacteria and/or a fecal pH which is more towards the alkaline end of the reference range (6 - 7.8). It is also not uncommon to find hemolytic or mucoid E. coli with a concomitant deficiency of beneficial E. coli and alkaline pH, secondary to a mutation of beneficial E. coli in alkaline conditions (DDI observations). Treatment with antimicrobial agents is unnecessary unless bacteria appear under the dysbiotic category.

Mackowiak PA. The normal microbial flora. N Engl J Med. 1982;307(2):83-93.

Cultured Yeast

Yeast, such as *Candida* are normally present in the GI tract in very small amounts. Many species of yeast exist and are commensal; however, they are always poised to create opportunistic infections and have detrimental effects throughout the body. Factors that contribute to a proliferation of yeast include frequent use of wide-spread antibiotics/low levels of beneficial flora, oral contraceptives, pregnancy, cortisone and other immunosuppressant drugs, weak immune system/low levels of sIgA, high-sugar diet, and high stress levels.

When investigating the presence of yeast, disparity may exist between culturing and microscopic examination. Yeast grows in colonies and is typically not uniformly dispersed throughout the stool. This may lead to undetectable or low levels of yeast identified by microscopy, despite a cultured amount of yeast. Conversely, microscopic examination may reveal a significant amount of yeast present, but no yeast cultured. Yeast does not always survive transit through the intestines rendering it unviable for culturing. Therefore, both microscopic examination and culture are helpful in determining if abnormally high levels of yeast are present.

Beneficial Flora

One or more of the expected or beneficial bacteria are low in this specimen. Normally abundant include lactobacilli, bifidobacteria, clostridia, *Bacteroides fragilis* group, enterococci, and some strains of *Escherichia coli*. The beneficial flora have many health-protecting effects in the gut, and as a consequence, are crucial to the health of the whole organism. Some of the roles of the beneficial flora include digestion of proteins and carbohydrates, manufacture of vitamins and essential fatty acids, increase in the number of immune system cells, break down of bacterial toxins and the conversion of flavinoids into anti-tumor and anti-inflammatory factors. Lactobacilli, bifidobacteria, clostridia, and enterococci secrete lactic acid as well as other acids including acetate, propionate, butyrate, and valerate. This secretion causes a subsequent decrease in intestinal pH, which is crucial in preventing an enteric proliferation of microbial pathogens, including bacteria and yeast. Many GI pathogens thrive in alkaline environments. Lactobacilli also secrete the antifungal and antimicrobial agents lactocidin, lactobacillin, acidolin, and hydrogen peroxide. The beneficial flora of the GI have thus been found useful in the inhibition of microbial pathogens, prevention and treatment of antibiotic associated diarrhea, prevention of traveler's diarrhea, enhancement of immune function, and inhibition of the proliferation of yeast.

In a healthy balanced state of intestinal flora, the beneficial flora make up a significant proportion of the total microflora. Healthy levels of each of the beneficial bacteria are indicated by either a 3+ or 4+ (0 to 4 scale). However, some individuals have low levels of beneficial bacteria and an overgrowth of nonbeneficial (imbalances) or even pathogenic microorganisms (dysbiosis). Often attributed to the use of antibiotics, individuals with low beneficial bacteria may present with chronic symptoms such as irregular transit time, irritable bowel syndrome, bloating, gas, chronic fatigue, headaches, autoimmune diseases (e.g., rheumatoid arthritis), and sensitivities to a variety of foods. Treatment may include the use of probiotic supplements containing various strains of lactobacillus and bifidobacterium species and consumption of cultured or fermented foods including yogurt, kefir, miso, tempeh and tamari sauce. Polyphenols in green and ginseng tea have been found to increase the numbers of beneficial

Lab number:
Patient:

Microbiology Profile

Page: 3

bacteria. If dysbiosis is present, treatment may also include the removal of pathogenic bacteria, yeast, or parasites.

Percival M. Intestinal Health. Clin Nutr In. 1997;5(5):1-6.

Fuller R. Probiotics in Human Medicine. Gut. 1991;32: 439-442.

Siitonen S, Vapaatalo H, Salminen S, et al. Effect of Lactobacilli GG Yoghurt in Prevention of Antibiotic Associated Diarrhea. Ann Med. 1990; 22:57-59.

Oksanen P, Salminen S, Saxelin M, et al. Prevention of Travelers' Diarrhea by Lactobacillus GG. Ann Med. 1990; 22:53-56.

Perdigon G, Alvarez M, et al. The Oral Administration of Lactic Acid Bacteria Increases the Mucosal Intestinal Immunity in Response to Enteropathogens. J Food Prot. 1990;53:404-410.

Valeur, N, et al. Colonization and Immunomodulation by Lactobacillus reuteri ATCC 55730 in the Human Gastrointestinal Tract. Appl Environ. Microbiol. 2004 Feb; 70(2):1176-81.

Elmer G, Surawicz C, and McFarland L. Brotherapeutic agents - a Neglected Modality for the Treatment and Prevention of Intestinal and Vaginal Infections. JAMA. 1996; 275(11):870-876.

Fitzsimmons N and Berry D. Inhibition of Candida albicans by Lactobacillus acidophilus: Evidence for Involvement of a Peroxidase System. Microbio. 1994; 80:125-133

Weisburger JH. Proc Soc Exp Biol Med 1999;220(4):271-5.

**SAMPLE
REPORT**

Lab number:
Patient:

Microbiology Profile

Page: 4

SAMPLE REPORT