

Kinetic Isotope Effects for Alkaline Phosphatase Reactions: Implications for the Role of Active-Site Metal Ions in Catalysis

Jesse G. Zalatan,[†] Irina Catrina,^{‡,§} Rebecca Mitchell,[‡] Piotr K. Grzyska,^{‡,||}
Patrick J. O'Brien,^{⊥,#} Daniel Herschlag,^{*,⊥,†} and Alvan C. Hengge^{*,‡}

Contribution from the Departments of Chemistry and Biochemistry, Stanford University, Stanford, California 94305, and Department of Chemistry and Biochemistry, Utah State University, Logan, Utah 84322

Received April 6, 2007; E-mail: herschla@stanford.edu; hengge@cc.usu.edu

Abstract: Enzyme-catalyzed phosphoryl transfer reactions have frequently been suggested to proceed through transition states that are altered from their solution counterparts, with the alterations presumably arising from interactions with active-site functional groups. In particular, the phosphate monoester hydrolysis reaction catalyzed by *Escherichia coli* alkaline phosphatase (AP) has been the subject of intensive scrutiny. Recent linear free energy relationship (LFER) studies suggest that AP catalyzes phosphate monoester hydrolysis through a loose transition state, similar to that in solution. To gain further insight into the nature of the transition state and active-site interactions, we have determined kinetic isotope effects (KIEs) for AP-catalyzed hydrolysis reactions with several phosphate monoester substrates. The LFER and KIE data together provide a consistent picture for the nature of the transition state for AP-catalyzed phosphate monoester hydrolysis and support previous models suggesting that the enzymatic transition state is similar to that in solution. Moreover, the KIE data provides unique information regarding specific interactions between the transition state and the active-site Zn²⁺ ions. These results provide strong support for a model in which electrostatic interactions between the bimetallo Zn²⁺ site and a nonbridging phosphate ester oxygen atom make a significant contribution to the large rate enhancement observed for AP-catalyzed phosphate monoester hydrolysis.

Introduction

Phosphoryl transfer reactions play a fundamental role in a wide range of biological processes including basic metabolism, energy transduction, gene expression, and cell signaling. Uncatalyzed phosphoryl transfer reactions are extremely slow, but enzymes can provide rate enhancements of >10²⁰-fold to allow these reactions to occur on a biologically relevant time scale.¹ As enzymatic rate enhancements can be considered to arise from preferential stabilization of the transition state relative to the ground state,^{2–6} characterizing the structure of the transition state is a critical step toward understanding how enzymes interact with and stabilize the transition state to provide the enormous rate enhancements typically associated with biological catalysis.

Escherichia coli alkaline phosphatase (AP) is an ideal enzyme for detailed structure–function analysis directed at understanding the nature of the enzymatic transition state. The AP active site contains two Zn²⁺ ions and an arginine residue that are positioned to interact with the substrate and a serine alkoxide that displaces the leaving group in the first step of the reaction to produce a covalent enzyme–phosphate intermediate (Scheme 1).^{7–9} This intermediate is hydrolyzed in the second step of the reaction.⁹ Of particular importance for mechanistic analysis is AP's relatively low substrate specificity, which has allowed linear free energy relationship (LFER) studies to be performed. These studies have suggested that the transition state for AP-catalyzed phosphate monoester hydrolysis is loose, similar to that in solution.^{10–13} The results obtained with AP are consistent with an emerging body of experimental data in a few well-studied enzyme systems suggesting that the transition states for enzyme catalyzed phosphoryl transfer reactions are similar to those in solution.¹⁴

[†] Department of Chemistry, Stanford University.

[‡] Department of Chemistry and Biochemistry, Utah State University.

[§] Present Address: Department of Biological Sciences, Hunter College.

^{||} Present Address: Department of Microbiology and Molecular Genetics, Michigan State University.

[⊥] Department of Biochemistry, Stanford University.

[#] Present Address: Department of Biological Chemistry, University of Michigan.

(1) Lad, C.; Williams, N. H.; Wolfenden, R. *Proc. Natl. Acad. Sci. U.S.A.* **2003**, *100*, 5607–5610.

(2) Polanyi, M. Z. *Elektrochem. Z.* **1921**, *27*, 142.

(3) Pauling, L. *Chem. Eng. News* **1946**, *24*, 1375–1377.

(4) Wolfenden, R. *Acc. Chem. Res.* **1972**, *5*, 10–18.

(5) Lienhard, G. E. *Science* **1973**, *180*, 149–154.

(6) Jencks, W. P., *Catalysis in Chemistry and Enzymology*; Dover Publications, Inc.: New York, 1987.

(7) Kim, E. E.; Wyckoff, H. W. *J. Mol. Biol.* **1991**, *218*, 449–464.

(8) Holtz, K. M.; Stec, B.; Kantrowitz, E. R. *J. Biol. Chem.* **1999**, *274*, 8351–8354.

(9) Coleman, J. E. *Annu. Rev. Biophys. Biomol. Struct.* **1992**, *21*, 441–483.

(10) O'Brien, P. J.; Herschlag, D. *J. Am. Chem. Soc.* **1999**, *121*, 11022–11023.


(11) Hollfelder, F.; Herschlag, D. *Biochemistry* **1995**, *34*, 12255–12264.

(12) O'Brien, P. J.; Herschlag, D. *Biochemistry* **2002**, *41*, 3207–3225.

(13) Nikolic-Hughes, I.; Rees, D. C.; Herschlag, D. *J. Am. Chem. Soc.* **2004**, *126*, 11814–11819.

(14) See references in Zalatan and Herschlag, 2006.⁴⁴

Scheme 1. Transition-State Model for Phosphoryl Transfer Catalyzed by AP Based on the Structure with a Bound Vanadate Ligand^{7,8}



Suggestions that enzymes alter the nature of the transition state for phosphoryl transfer reactions remain widespread,¹⁵ despite the increasing number of experimental studies suggesting otherwise. Thus, we sought to further test the conclusions of the LFER analysis by measuring kinetic isotope effects (KIEs) for AP-catalyzed reactions with several substrates. KIEs report on the nature of the transition state as well as interactions between the transition state and enzyme functional groups and provide information complementary to that obtained from LFER studies.¹⁶ The KIE approach involves the smallest possible perturbation to a specific site in the substrate, and thus circumvents many of the limitations of site-directed mutagenesis in providing information about specific active-site interactions.¹⁷

The results described herein are consistent with the conclusion from LFER studies suggesting that the transition state for AP-catalyzed phosphate monoester hydrolysis is loose, similar to that in solution. The KIE data further provide strong evidence in support of a model in which electrostatic interactions between the bimetallo Zn^{2+} site and a nonbridging phosphate ester oxygen atom make a significant contribution to the large rate enhancement observed for AP-catalyzed phosphate monoester hydrolysis.

Methods

Materials. The plasmids for expression of wild-type (wt) and R166S AP and the $phoA^-$ strain of *E. coli* (SM547) were provided by Evan Kantrowitz.¹⁸ Natural abundance and isotopically labeled *p*-nitrophenyl phosphate (pNPP), *p*-nitrophenyl phosphorothioate (pNPPS), and *m*-nitrobenzyl phosphate (mNBP) were synthesized as previously described.^{19–21} All other reagents were obtained from commercial sources.

Purification of Wild-Type and R166S AP. Wild-type and R166S AP were purified as previously described,²² except that the dialysis step was omitted. Protein concentration was determined by absorbance at 280 nm using a calculated extinction coefficient of $3.14 \times 10^4 \text{ M}^{-1} \text{ cm}^{-1}$ (in monomer units).²³

AP-Catalyzed mNBP Hydrolysis. Reactions were performed with 0.1 M NaMOPS, pH 8.0, 0.5 M NaCl, 1 mM $MgCl_2$, and 0.1 mM $ZnSO_4$ at 25 °C. Reactions were monitored using a malachite green assay to detect release of inorganic phosphate. Malachite green solutions were prepared and inorganic phosphate was assayed as previously described,²⁴ except that Triton X-100 was used in place of Sterox. Appearance of inorganic phosphate was monitored by removing 100 μL aliquots from the enzymatic reaction and quenching in 800 μL of the malachite green solution, which contains strong acid. After 1 min, 100 μL of a 34% sodium citrate solution was added. After 30 min, absorbance at 644 nm was measured in a Uvikon 9310 spectrophotometer. Rate constants for R166S AP-catalyzed pNPP hydrolysis obtained from the malachite green assay agreed within 10% of those obtained by continuously monitoring release of *p*-nitrophenolate at 400 nm.

Reactions of mNBP were typically followed to completion (≥ 10 half-lives). Rate constants were obtained by nonlinear least-squares fits to a single exponential to yield k_{obs} , which is the product of the enzyme concentration and the apparent second-order rate constant k_{cat}/K_M under the subsaturating reaction conditions used herein. Reactions were shown to be first order in substrate and enzyme by varying their concentrations over at least a 10-fold range. To ensure that reactions of wt AP were conducted under k_{cat}/K_M conditions, these reactions were conducted in the presence of 1 mM sodium tungstate, an inhibitor of AP.¹² The value of k_{cat}/K_M could then be calculated from the observed second-order rate constant with inhibition and the known inhibition constant for tungstate ($K_I = 1.1 \mu\text{M}$ at pH 8.0).¹²

Isotope Effect Determinations. ¹⁸O KIEs were measured using isotope ratio mass spectrometry by the remote label method, using the nitrogen atom as a reporter for isotopic fractionation at the bridge or nonbridge oxygen atoms. The experimental procedures used to measure these isotope effects were similar to those previously reported.¹⁶

All kinetic isotope effect experiments were run at least in triplicate. Tests were run to determine enzyme and substrate concentrations to ensure that the enzymatic rates of reaction were at least 50 times faster than the aqueous hydrolysis. Background nonenzymatic hydrolysis was a concern with pNPPS but not with mNBP, since the uncatalyzed hydrolysis of the latter substrate is extremely slow.

Isotope effects were calculated from the isotopic ratios at partial reaction in the product (R_p), in the residual substrate (R_s), and in the starting material (R_o).


$$\text{KIE} = [\log(1 - f)] / \{\log[1 - f(R_p/R_o)]\} \quad (1)$$

$$\text{KIE} = [\log(1 - f)] / \{\log[(1 - f)(R_s/R_o)]\} \quad (2)$$

Equations 1 or 2 were used to calculate the observed isotope effect either from R_p and R_o or from R_s and R_o , respectively, at the measured fraction of reaction.²⁵ The observed isotope effects from experiments to determine ¹⁸O isotope effects were corrected for the ¹⁵N effect and for incomplete levels of isotopic incorporation in the starting material as previously described.²⁶ The independent calculation of each isotope effect using R_p and R_o and using R_s and R_o from eqs 1 and 2, respectively, provided an internal check of the results.

- (15) See references in O'Brien and Herschlag, 1999, and Nikolic-Hughes, Rees, and Herschlag, 2004.^{10,13} See also: (a) Lahiri, S. D.; Zhang, G. F.; Dunaway-Mariano, D.; Allen, K. N. *Science* **2003**, *299*, 2067–2071. (b) Allen, K. N.; Dunaway-Mariano, D. *Trends Biochem. Sci.* **2004**, *29*, 495–503. (c) Krishnamurthy, H.; Lou, H. F.; Kimple, A.; Vielle, C.; Cukier, R. I. *Proteins* **2005**, *58*, 88–100. (d) Bellinzoni, M.; Haouz, A.; Grana, M.; Munier-Lehmann, H.; Shepard, W.; Alzari, P. M. *Protein Sci.* **2006**, *15*, 1489–1493. (e) Hartmann, M. D.; Bourenkov, G. P.; Oberschall, A.; Strizhov, N.; Bartunik, H. D. *J. Mol. Biol.* **2006**, *364*, 411–423. (f) Wittinghofer, A. *Trends Biochem. Sci.* **2006**, *31*, 20–23. Note that the data reported in ref 15a has recently been reexamined in: (g) Baxter, N. J.; Olguin, L. F.; Golicnik, M.; Feng, G.; Hounslow, A. M.; Bermel, W.; Blackburn, G. M.; Hoffelder, F.; Waltho, J. P.; Williams, N. H. *Proc. Natl. Acad. Sci. U.S.A.* **2006**, *103*, 14732–14737.
- (16) Hengge, A. C. *Acc. Chem. Res.* **2002**, *35*, 105–112.
- (17) Kraut, D. A.; Carroll, K. S.; Herschlag, D. *Annu. Rev. Biochem.* **2003**, *72*, 517–571.
- (18) Chaidaroglou, A.; Brezinski, D. J.; Middleton, S. A.; Kantrowitz, E. R. *Biochemistry* **1988**, *27*, 8338–8343.
- (19) Hengge, A. C.; Edens, W. A.; Elsing, H. *J. Am. Chem. Soc.* **1994**, *116*, 5045–5049.
- (20) Catrina, I. E.; Hengge, A. C. *J. Am. Chem. Soc.* **2003**, *125*, 7546–7552.
- (21) Grzyska, P. K.; Czyryca, P. G.; Purcell, J.; Hengge, A. C. *J. Am. Chem. Soc.* **2003**, *125*, 13106–13111.

- (22) O'Brien, P. J.; Herschlag, D. *Biochemistry* **2001**, *40*, 5691–5699.
- (23) Gill, S. C.; von Hippel, P. H. *Anal. Biochem.* **1989**, *182*, 319–326.
- (24) Lanzetta, P. A.; Alvarez, L. J.; Reinach, P. S.; Candia, O. A. *Anal. Biochem.* **1979**, *100*, 95–97.
- (25) Bigeleisen, J.; Wolfsberg, M. *Adv. Chem. Phys.* **1958**, *1*, 15–76.
- (26) Caldwell, S. R.; Raushel, F. M.; Weiss, P. M.; Cleland, W. W. *Biochemistry* **1991**, *30*, 7444–7450.

Scheme 2. Phosphate Monoester Hydrolysis through a Loose Transition State

KIEs for AP-Catalyzed pNPPS Hydrolysis. The reactions catalyzed by wt AP were performed in 10 mL of 0.1 M Tris·HCl, pH 9.0, 1 mM MgCl₂, 1 mM ZnSO₄, and 10 mM pNPPS at 35 °C. The reactions catalyzed by R166S AP were performed in a 2.5 mL volume of 0.5 M Tris·HCl, pH 8.0, 100 μM MgCl₂, 10 μM ZnSO₄, and 40 mM pNPPS at 25 °C. The enzyme concentration was 1 μM. After partial hydrolysis (ranging from 35 to 65%), the reactions were stopped by lowering the pH to 5.5 through the addition of cold 0.2 M glycine buffer (pH 2) and placing the mixture on ice. From each reaction mixture, two aliquots were taken; one 50 μL aliquot was added to a cuvette containing 3 mL of 0.1 N NaOH and the absorbance at 400 nm was used to determine the concentration of *p*-nitrophenol at partial hydrolysis. The second aliquot was adjusted to pH 2 in glycine buffer and allowed to react at 50 °C for 12 h to completely hydrolyze the remaining substrate. Control experiments showed that 12 h corresponds to >10 times the half-life for pNPPS hydrolysis under these conditions. Subsequently, a 50 μL aliquot of this sample was added to 3 mL of 0.1 N NaOH and the absorbance at 400 nm was used to determine the concentration of *p*-nitrophenol product at complete hydrolysis, which corresponds to the initial concentration of substrate. The concentration of *p*-nitrophenol at partial hydrolysis and the initial concentration of substrate were used to determine the fraction of reaction.

The reaction solutions at pH 5.5 were extracted three times with an equal volume of diethyl ether to quantitatively remove the product of the partial hydrolysis, *p*-nitrophenol. The water layers were titrated to pH 2 and allowed to react overnight at 50 °C to completely hydrolyze the residual substrate. The *p*-nitrophenol formed was then extracted using the same procedure as described above. The ether layers were dried over MgSO₄, filtered, and evaporated until dryness. The *p*-nitrophenol was sublimed under vacuum at 90 °C, and 1–1.5 mg samples were prepared for isotopic analysis using an ANCA-NT combustion system coupled with a Europa 20–20 isotope ratio mass spectrometer.

KIEs for AP-Catalyzed mNBP Hydrolysis. AP-catalyzed mNBP hydrolysis reactions were performed in 5 mL of NaMOPS, pH 8.0, 1 mM MgCl₂, 100 μM ZnSO₄, and 20 mM mNBP at 30 °C and an enzyme concentration of 0.3 μM. ³¹P NMR was used to monitor reaction progress. After partial reaction (ranging from 40 to 80%), the reactions were stopped by titration to pH 4. NMR was used to obtain the fraction of reaction from integrations of the signals from substrate and inorganic phosphate. The reaction mixtures were extracted with 25 mL of diethyl ether four times to obtain the *m*-nitrobenzyl alcohol product. The ether layers were dried over MgSO₄, filtered, and evaporated until dryness. The aqueous layer was titrated to pH 8 and about 1 mg of commercial alkaline phosphatase was added to hydrolyze the remaining substrate. After 12 h, this mixture was titrated to pH 4 and treated as before to recover the liberated *m*-nitrobenzyl alcohol. The *m*-nitrobenzyl alcohol samples were further purified by distillation under vacuum at ~105 °C onto a cold finger apparatus, and samples were then subjected to isotopic analysis, as described above.

Results and Discussion

Decades of extensive mechanistic studies have strongly suggested that phosphate monoester hydrolysis in solution proceeds through a loose transition state, with nearly complete bond cleavage to the leaving group and little bond formation to the nucleophile (Scheme 2).^{19,27–31} Results from LFER experiments suggest that AP-catalyzed phosphate monoester

hydrolysis also proceeds through a loose transition state, indistinguishable from that in solution.^{10–13,32} To gain further insight into the nature of the transition state for AP-catalyzed phosphate monoester hydrolysis, we have compared the reactivity of AP with several classes of phosphate ester substrates and determined KIEs for representative members of these classes. The KIEs were determined by the competitive method and are therefore effects on V/K (i.e., k_{cat}/K_M)¹⁶ that reflect differences between the enzymatic transition state and the ground state in which the enzyme and substrate are free in solution.

In addition to characterizing the nature of the transition state, the KIE data have allowed us to compare AP-catalyzed reactions of alkyl and aryl phosphates. AP catalyzes reactions of alkyl phosphates with significantly greater catalytic proficiency than reactions of aryl phosphates.³³ To probe the origin of this effect, we have obtained KIEs with two phosphate monoester substrates with significantly different leaving groups: *p*-nitrophenyl phosphate (pNPP) and *m*-nitrobenzyl phosphate (mNBP) (Figure 1). Both of these substrates have nitrogen containing functional groups that allow high precision KIE measurements using the remote-label method.¹⁶ The pK_a of the *p*-nitrophenyl leaving group of pNPP is 7.1,³⁴ and this substrate has been routinely used in studies of phosphatase enzymes such as AP.¹² In contrast, mNBP has not previously been studied as a substrate of AP. The high pK_a of the *m*-nitrobenzyl leaving group of mNBP (pK_a = 15)³⁵ is expected to lead to reactivity analogous to that of alkyl phosphates, which have similar leaving group pK_a values. The *m*-nitrobenzyl leaving group has been used previously in studies of both solution and enzyme-catalyzed phosphoryl transfer reactions and has been particularly valuable in allowing KIE comparisons between alkyl and aryl phosphates.^{21,35–37} We first compare the reactivity of aryl, alkyl, and benzyl phosphates in AP-catalyzed reactions and then analyze the KIE data in the context of these reactivity comparisons. Finally, we consider KIE data for AP-catalyzed *p*-nitrophenyl phosphorothioate (pNPPS) hydrolysis, and the results reinforce the conclusions from the analysis of phosphate monoester substrates.

- (27) Kirby, A. J.; Jencks, W. P. *J. Am. Chem. Soc.* **1965**, *87*, 3209–3216.
- (28) Kirby, A. J.; Varvoglis, A. G. *J. Am. Chem. Soc.* **1967**, *89*, 415–423.
- (29) Thatcher, G. R. J.; Kluger, R. *Adv. Phys. Org. Chem.* **1989**, *25*, 99–265.
- (30) Hengge, A. C. *Adv. Phys. Org. Chem.* **2005**, *40*, 49–108.
- (31) Cleland, W. W.; Hengge, A. C. *Chem. Rev.* **2006**, *106*, 3252–3278.
- (32) Holtz, K. M.; Catrina, I. E.; Hengge, A. C.; Kantrowitz, E. R. *Biochemistry* **2000**, *39*, 9451–9458.
- (33) The catalytic proficiency is defined as the ratio of the second order rate constants for enzymatic and solution reactions [(k_{cat}/K_M)/ k_W]. R166S AP has a catalytic proficiency of 2×10^{15} for pNPP²⁻ hydrolysis⁴³ and $>10^{23}$ for methyl phosphate hydrolysis.^{1,10} The overall difference in catalytic proficiency between pNPP²⁻ and methyl phosphate is $>10^8$ -fold.
- (34) Jencks, W. P.; Regenstein, J., *Ionization Constants of Acids and Bases. In Handbook of Biochemistry and Molecular Biology*; Fasman, G. D., Ed.; CRC: Cleveland, OH, 1976.
- (35) Sowa, G. A.; Hengge, A. C.; Cleland, W. W. *J. Am. Chem. Soc.* **1997**, *119*, 2319–2320.
- (36) Anderson, M. A.; Shim, H.; Raushel, F. M.; Cleland, W. W. *J. Am. Chem. Soc.* **2001**, *123*, 9246–9253.
- (37) Gerratana, B.; Sowa, G. A.; Cleland, W. W. *J. Am. Chem. Soc.* **2000**, *122*, 12615–12621.





Figure 1. Phosphate and phosphorothioate esters for which isotope effects are reported: red, nonbridge phosphoryl oxygen atoms [$^{18}(V/K)_{\text{nonbridge}}$]; blue, bridge oxygen atom, site of bond fission [$^{18}(V/K)_{\text{bridge}}$]; green, nitrogen atom of the leaving group [$^{15}(V/K)$]. The structures are depicted with partial double bonds to emphasize that the nonbridging oxygen atoms that contribute to $^{18}(V/K)_{\text{nonbridge}}$ are formally equivalent when the substrates are free in solution. For simplicity, the P–S and P–O bonds of pNPPS are depicted with the same partial double-bond character, although it is likely that the P–S bond has more single-bond character than the P–O bonds.^{43,83}

Scheme 3



Reactivity Comparisons of Nonenzymatic Reactions. Before comparing the reactivity of AP with alkyl, benzyl, and aryl phosphates, it is important to determine whether these compounds react via similar mechanisms in solution. The dianionic forms of the phosphate esters (Scheme 3) are the substrates for AP,¹² but kinetic constants for the corresponding nonenzymatic reactions of alkyl and benzyl phosphate dianions cannot be readily measured, as these reactions are extremely slow and can proceed via C–O bond cleavage instead of P–O cleavage.^{1,38–40} The nonenzymatic reactions of alkyl, aryl, and benzyl phosphate monoanions react via P–O cleavage, and rates can be measured, however.

Reactions of phosphate monoester monoanions have been proposed to proceed with proton transfer from the phosphoryl group to the leaving group in the transition state, thereby neutralizing the charge that would otherwise develop on the leaving group oxygen.^{19,21,28,38–41} The available literature data for alkyl, aryl, and benzyl phosphate monoanions suggest that each of these classes of compounds react via this mechanism. Hydrolysis reactions of alkyl and aryl phosphate monoanions follow the same leaving group dependence, with little change in the hydrolysis reaction rate with leaving group pK_a , consistent




Figure 2. Compilation of kinetic data for the reactions of wt and R166S AP, represented by closed and open symbols, respectively. Reactions of alkyl phosphates (■, □) give values of β_{lg} of -0.85 ± 0.1 and -0.66 ± 0.1 respectively for reactions with wt and R166S AP.^{10,12} Reactions of aryl phosphates (◆) with wt AP are limited by a nonchemical step (dashed line).¹² The rate constant for the reaction of the aryl phosphate pNPP²⁻ with R166S AP²² deviates negatively from the value expected by extrapolation from the line for alkyl phosphates (dotted line and single-headed arrow). Rate constants for the reactions of mNBP²⁻ with wt and R166S AP (●, ○) deviate positively from the values expected from the line for alkyl phosphates (double-headed arrows).

with the long-standing mechanistic proposal for proton transfer in the transition state.^{28,41} The hydrolysis of benzyl phosphate has a similar rate constant and pH-dependence with that observed for alkyl phosphate hydrolysis, except under strong acid conditions in which the neutral benzyl phosphate reacts via C–O cleavage.^{38,39,42} Thus, there appears to be no substantial difference in the intrinsic reactivity of alkyl, aryl, and benzyl phosphates, beyond effects from the differential stability of the leaving groups that is reflected in the different pK_a values.

Reactivity Comparisons of AP-Catalyzed Reactions. Figure 2 shows a compilation of kinetic data for the reactions of wt AP and a mutant of AP with the active-site Arg replaced by Ser (R166S AP), represented by closed and open symbols, respectively. R166S AP has been used in a number of mechanistic studies of AP,^{10,32,43,44} in part because it is not subject to the severe product inhibition that has often complicated mechanistic studies of wt AP.¹² The different substrate classes and the symbols used to represent them in Figure 2 are depicted in Scheme 3. Rate constants for reactions of mNBP²⁻ with wt and R166S AP (circles) were determined in this work, and all other rate constants are from previous work.^{10,12}

The rates of reaction of the alkyl, aryl, and benzyl phosphates with AP differ considerably, even after accounting for differential leaving group stability. Reactions of substrates with low pK_a leaving groups, such as aryl phosphates, with wt AP are limited by binding rather than the chemical step and are independent of leaving group pK_a (dashed line).^{12,45,46} Rate constants for alkyl phosphates with good leaving groups such as trifluoroethyl phosphate and propargyl phosphate (leaving

(38) Butcher, W. W.; Westheimer, F. H. *J. Am. Chem. Soc.* **1955**, *77*, 2420–2424.

(39) Kumamoto, J.; Westheimer, F. H. *J. Am. Chem. Soc.* **1955**, *77*, 2515–2518.

(40) Bunton, C. A.; Llewellyn, D. R.; Oldham, K. G.; Vernon, C. A. *J. Chem. Soc.* **1958**, 3574–3587.

(41) Bunton, C. A.; Fendler, E. J.; Humeres, E.; Yang, K. U. *J. Org. Chem.* **1967**, *32*, 2806–2811.

(42) Parente, J. E.; Risley, J. M.; Van Etten, R. L. *J. Am. Chem. Soc.* **1984**, *106*, 8156–8161.

(43) Nikolic-Hughes, I.; O'Brien, P. J.; Herschlag, D. *J. Am. Chem. Soc.* **2005**, *127*, 9314–9315.

(44) Zalatan, J. G.; Herschlag, D. *J. Am. Chem. Soc.* **2006**, *128*, 1293–1303.

(45) Labow, B. I.; Herschlag, D.; Jencks, W. P. *Biochemistry* **1993**, *32*, 8737–8741.

(46) Simopoulos, T. T.; Jencks, W. P. *Biochemistry* **1994**, *33*, 10375–10380.

group pK_a values of 12.4 and 13.6, respectively) also fall on the dashed line, suggesting that these reactions may not be limited by the chemical step. In contrast, the rate constants for less reactive alkyl phosphates are smaller and depend on leaving group pK_a , suggesting that the chemical step is rate-determining. The slope of the line correlating the log of the rate constant with the pK_a of the leaving group (the Brønsted coefficient, β_{lg}) is -0.85 .¹² The values of β_{lg} for alkyl phosphates are nearly the same for wt and R166S AP (Figure 2), suggesting that the interaction of the Arg side chain with the nonbridging phosphoryl oxygen atoms does not alter the nature of the transition state.¹⁰ Further, the values of β_{lg} for reactions of aryl phosphorothioate monoesters with both wt and R166S AP are the same and similar to those for the alkyl phosphates.^{11,32} These results are consistent with a similar loose transition state with substantial charge buildup on the leaving group oxygen for reactions of all phosphate monoester substrates, possibly with some lessening of the value of β_{lg} relative to solution reactions because of the electrostatic interaction of the leaving group oxygen atom with an active-site Zn^{2+} ion (see also below).^{12,13}

The enzymatic reactivity of $mNBP^{2-}$ can be directly compared with that of simple alkyl phosphates, as the leaving groups for these compounds fall in the same pK_a range. The rate constants for $mNBP^{2-}$ hydrolysis by wt and R166S AP are 1.8×10^7 and $2.3 \times 10^3 \text{ M}^{-1} \text{ s}^{-1}$, respectively. These values deviate positively from the correlation line for alkyl phosphates for both wt and R166S AP, falling ~ 10 -fold above their expected values (Figure 2; open and closed circles and double-headed arrows). The simplest model for the increased reactivity of $mNBP^{2-}$ is additional binding interactions with the larger benzyl leaving group (Scheme 3) within the AP active site. In this model, the charge buildup on the bridging oxygen in the transition state for $mNBP^{2-}$ hydrolysis is likely to be similar to that for simple alkyl phosphates.

An alternative explanation for the enhanced reactivity of $mNBP$ is a distinct catalytic mechanism involving the monoanion as the substrate for catalysis with intramolecular protonation of the leaving group, although previous studies of both alkyl and aryl phosphates with wt and R166S AP indicate that the dianion is the substrate for catalysis.^{12,22} To examine this possibility, we determined the pH-rate profile for the reaction of $mNBP$ with R166S AP over the pH range of 7–9. These data were sufficient to confirm that the pH dependence was identical to that observed previously for $pNPP$ and other phosphate monoester substrates, which react as the dianion,^{12,22} and strongly suggests that $mNBP$ also reacts as a dianion (Supporting Information). If $mNBP$ were to react as a monoanion, the pH dependence would have been predicted to deviate substantially from that observed for $pNPP$. Thus, there is no indication from the pH-rate profile of a change in rate-limiting step or mechanistic path for the $mNBP$ reaction from that previously observed for other AP-catalyzed reactions.

In contrast to the positive deviation observed with $mNBP^{2-}$, the rate constant for R166S AP-catalyzed $pNPP^{2-}$ hydrolysis falls $\sim 10^3$ -fold below the correlation line for the alkyl phosphate rate constants extrapolated to lower pK_a (Figure 2, dotted line and single-headed arrow). This comparison can only be made for R166S AP, as the reaction of $pNPP^{2-}$ with wt AP is limited by a nonchemical step (Figure 2, dashed line).¹² Because the leaving group oxygen for $pNPP^{2-}$ is attached to a secondary

carbon whereas the alkyl phosphate leaving groups are primary alkoxides, the deviation could arise from a steric effect. However, rate constants for substrates with *p*-nitrophenyl leaving groups also deviate negatively with respect to other aryl leaving groups in AP-catalyzed reactions of aryl phosphorothioates,^{11,32} aryl sulfates,¹³ methyl phenyl phosphate diesters,⁴⁴ and in several solution reactions.⁴⁴ This effect presumably arises from greater delocalization of the negative charge into the aromatic ring for *p*-nitrophenolate than for other aryl leaving groups.⁴⁴ Thus, the deviation of the rate constant for $pNPP^{2-}$ hydrolysis relative to that expected from alkyl phosphate reactions could arise because of differing amounts of negative charge localized on the leaving-group oxygen. Alkyl phosphates cannot delocalize charge into an aromatic ring as in aryl phosphates and are expected to have greater charge localization on the leaving-group oxygen in the transition state. Nevertheless, we cannot rule out steric contributions to the deviation observed for $pNPP^{2-}$.

Kinetic Isotope Effects for AP-Catalyzed Reactions. Determination of KIEs for reactions of alkyl and aryl phosphates with AP could provide insight into the nature of the transition state and active-site interactions for AP-catalyzed phosphate monoester hydrolysis that is complementary to the information obtained from LFER studies. Because a remote ^{15}N label is required to achieve the precision needed for these measurements,¹⁶ we used the benzyl phosphate $mNBP^{2-}$ (Figure 1), which contains a nitro group that can be used as the remote label. $mNBP^{2-}$ has a leaving group pK_a of 15, similar to most alkyl leaving groups and substantially greater than the pK_a of 7 for the aryl leaving group of $pNPP^{2-}$. As noted above, the electronic properties of the leaving group oxygen for $mNBP^{2-}$ are expected to be similar to alkyl phosphates, and the KIEs for $mNBP^{2-}$ reactions with AP should provide a reasonable model for the behavior of alkyl phosphates in the AP active site.

Heavy atom KIEs were determined at three different positions in each substrate (Figure 1). The ^{18}O nonbridging effect [$^{18}(\text{V}/K)_{\text{nonbridge}}$] is a secondary KIE reporting on changes in both hybridization and bond order. This effect can be diagnostic of the looseness or tightness of the transition state but can also be perturbed by interactions with the nonbridging oxygen atoms.¹⁶ The ^{18}O bridging effect [$^{18}(\text{V}/K)_{\text{bridge}}$] is a primary KIE that reports on the extent of bond cleavage to the leaving group, and is also sensitive to interactions that stabilize charge buildup such as protonation.¹⁶ Finally, the ^{15}N label allows ^{18}O KIE determination by the remote label method. With *p*-nitrophenyl leaving groups, the ^{15}N effect [$^{15}(\text{V}/K)$] also reports on charge delocalization into the aromatic ring.¹⁶ No bonding changes at the nitro group are expected during $mNBP^{2-}$ hydrolysis, however, and $^{15}(\text{V}/K)$ for $mNBP^{2-}$ should be close to unity and can serve as an internal control.²¹

KIEs for AP-catalyzed reactions and corresponding solution reactions are reported in Table 1. KIEs for the reaction of wt AP with $pNPP^{2-}$ have been previously reported and are all near unity, consistent with a nonchemical rate-determining step.¹⁹ The KIEs for the reaction of $mNBP^{2-}$ with wt AP differ from unity but are significantly smaller than the KIEs determined for the reaction with R166S AP (see later). The reaction of $mNBP^{2-}$ with wt AP is presumably partially limited by a nonchemical step, as the value of k_{cat}/K_M is close to the limit at which chemistry is no longer rate-determining (Figure 2) and

Table 1. Kinetic Isotope Effects^a

enzymatic reactions ^b	substrate	¹⁵ (V/K)	¹⁸ (V/K) _{bridge}	¹⁸ (V/K) _{nonbridge}
wt AP ¹⁹	pNPP ²⁻	1.0003(2)	1.0003(4)	0.9982(1)
wt AP	mNBP ²⁻	0.9999(4)	1.0072(7)	0.9988(4)
wt AP	pNPPS ²⁻	1.0005(2)	1.0094(4)	0.9760(22)
R166S AP ⁴⁷	pNPP ²⁻	1.0007(1)	1.0091(6)	0.9925(11)
R166S AP	mNBP ²⁻	n.d. ^c	1.0199(13)	0.9933(4)
R166S AP	pNPPS ²⁻	1.0006(1)	1.0098(3)	0.9768(25)
uncatalyzed reactions	substrate	¹⁵ k	¹⁸ k _{bridge}	¹⁸ k _{nonbridge}
At 95 °C ¹⁹	pNPP ²⁻	1.0028(2)	1.0189 (5)	0.9994(5)
estimated ^d	mNBP ²⁻	1.0000	~1.04 to 1.06	~1.00
At 50 °C ²⁰	pNPPS ²⁻	1.0027(1)	1.0237(7)	1.0135(13)
At 95 °C ¹⁹	pNPP ¹⁻	1.0004(2)	1.0087(3)	1.0184(5)
At 115 °C ²¹	mNBP ¹⁻	1.0000(1)	1.0157(9)	1.0151(2)
At 30 °C ²⁰	pNPPS ¹⁻	1.0005(1)	1.0091(7)	1.0221(4)

^a Standard errors for the last decimal place(s) are in parenthesis. The nonbridge KIEs are the overall effects per molecule. The corresponding per-atom isotope effects are the cube roots (in the case of phosphate esters) or the square roots (in the case of phosphorothioate esters) of the values reported above. Data for the reaction of wt and R166S AP with pNPP²⁻ and for solution reactions are from previous work.^{19–21,47} Data for all other reactions were obtained for this work. The value of ¹⁸k_{bridge} of 1.0157 for mNBP¹⁻ has been corrected from the value originally reported in Grzyska et al., 2003.^{21,81} ^b Enzymatic reactions with pNPP²⁻ were conducted at 25 °C, and enzymatic reactions with mNBP²⁻ were conducted at 30 °C. The reactions with pNPPS²⁻ were conducted at 35 °C for wt AP and 25 °C for R166S AP. Although differences in temperature can effect the observed KIEs, these effects are small and do not affect the conclusions drawn herein.^{20,82} ^c Not determined. ^d KIEs for mNBP²⁻ hydrolysis in solution cannot be measured because hydrolysis is dominated by alternative reaction pathways.^{1,38–40} The estimates shown were obtained as described in the main text.

the full isotope effects appear to be masked relative to the isotope effects with a slower, mutant enzyme, R166S AP. Reactions of R166S AP with pNPP²⁻ and mNBP²⁻ are likely to be limited by the chemical step, as the values for k_{cat}/K_M are well below the limiting values observed for wt AP (Figure 2). Consistent with this expectation, several of the observed KIEs for these substrates are significantly different from unity (Table 1).

KIEs for R166S AP-Catalyzed pNPP²⁻ Hydrolysis. KIEs for the reaction of R166S AP with pNPP²⁻ have been previously reported and discussed in the context of reactivity comparisons with *p*-nitrophenyl sulfate.⁴⁷ Here we discuss the implications of the pNPP²⁻ reaction KIEs for the nature of the transition state and for comparisons of alkyl and aryl phosphates.

The value of ¹⁸(V/K)_{bridge} KIE for R166S AP-catalyzed pNPP²⁻ hydrolysis is 1.0091, significantly reduced from the value of 1.0189 observed for pNPP²⁻ hydrolysis in solution (Table 1). Similarly, the value of ¹⁵(V/K) of 1.0007 is significantly reduced from its solution value of 1.0028. Both values are closer to those observed for monoanion than dianion hydrolysis in solution, but the pNPP dianion is the substrate for AP and no proton transfer to the leaving group in the transition state is expected.^{12,22} One possible interpretation of these results is that the transition state for AP-catalyzed pNPP²⁻ hydrolysis is tighter than that in solution, with less bond cleavage to the leaving group. However, previous LFER and KIE data suggest that the transition state for AP-catalyzed phosphate monoester hydrolysis is loose, similar to that in solution.^{10–13,32,48}

Furthermore, if the enzymatic transition state were tighter than that in solution, a normal contribution to the nonbridge KIE would be expected (Figure 3).¹⁶ Instead, the observed value of ¹⁸(V/K)_{nonbridge} for R166S AP-catalyzed pNPP²⁻ hydrolysis is 0.9925, significantly more inverse than the solution value of 0.9994 (Table 1). Thus, when taken together the KIEs for R166S AP-catalyzed pNPP²⁻ hydrolysis do not suggest a change in the nature of the transition state from that in solution.

We suggest an alternative model in which strong interactions between the substrate and the active-site Zn²⁺ ions (Scheme 1) result in the observed reductions in the KIEs for R166S AP-catalyzed pNPP²⁻ hydrolysis relative to the values for the uncatalyzed reaction. Interactions with the leaving group oxygen atom would be reflected in both ¹⁸(V/K)_{bridge} and ¹⁵(V/K), and interactions with the nonbridging oxygen atom would be reflected in ¹⁸(V/K)_{nonbridge}. The observed inverse contribution to the ¹⁸O isotope effects could arise if coordination to Zn²⁺ were to stiffen bending modes or introduce new vibrational modes that involve motions of the substrate oxygen atoms. Coordination of the leaving-group oxygen by Zn²⁺ could also decrease electron delocalization into the nitro group, leading to the observed decrease in the ¹⁵N isotope effect relative to that in solution (Table 1).

KIEs for R166S AP-Catalyzed mNBP²⁻ Hydrolysis. Interpretation of the KIEs for enzymatic mNBP²⁻ hydrolysis is complicated by the fact that corresponding KIEs for mNBP dianion hydrolysis in solution cannot be measured. The observed reaction of mNBP in solution is expected to be dominated by reaction of mNBP as a monoanion and reaction of the dianion with C–O rather than P–O cleavage.^{1,38–40} In the absence of measured KIEs for the uncatalyzed reaction, reasonable estimates can be made based on available literature data. The value of ¹⁸k_{nonbridge} for mNBP²⁻ is likely to be small and inverse, similar to that for pNPP²⁻ in solution, because the nonbridging P–O bonds are chemically equivalent in mNBP²⁻ and pNPP²⁻. The value of ¹⁸k_{bridge}, however, should be significantly larger for an alkyl leaving group than for an aryl leaving group. The maximum expected ¹⁸O isotope effect for cleavage of a *p*-nitrophenyl leaving group is ~1.03, based on the equilibrium isotope effect for *p*-nitrophenyl acetate hydrolysis.⁴⁹ This value is consistent with values of ¹⁸O KIEs of ~1.02 to 1.03 observed for pNPP²⁻ hydrolysis under a variety of solution and enzymatic conditions.¹⁶ In contrast to the behavior of aryl leaving groups, departure of a methoxide leaving group in the hydrazinolysis of methyl formate and methyl benzoate gives leaving-group ¹⁸O KIEs of 1.06 and 1.04, respectively.^{50,51} Furthermore, C–O cleavage reactions catalyzed by fumarase and crotonase give leaving-group ¹⁸O KIEs of 1.07 and 1.05, respectively.^{52,53} Finally, maximum KIE values for C–O and P–O cleavage reactions with alkyl leaving groups of ~1.05 have been predicted on the basis of vibrational stretching frequencies.^{50,54} Thus, the value of ¹⁸k_{bridge} for mNBP²⁻ hydrolysis in solution proceeding through a loose transition state with nearly complete leaving-

(47) Catrina, I.; O'Brien P, J.; Purcell, J.; Nikolic-Hughes, I.; Zalatan, J. G.; Hengge, A. C.; Herschlag, D. *J. Am. Chem. Soc.* **2007**, *129*, 5760–5765.
(48) Weiss, P. M.; Cleland, W. W. *J. Am. Chem. Soc.* **1989**, *111*, 1928–1929.

(49) Hengge, A. C.; Hess, R. A. *J. Am. Chem. Soc.* **1994**, *116*, 11256–11263.
(50) Sawyer, C. B.; Kirsch, J. F. *J. Am. Chem. Soc.* **1973**, *95*, 7375–7381.
(51) O'Leary, M. H.; Marlier, J. F. *J. Am. Chem. Soc.* **1979**, *101*, 3300–3306.
(52) Blanchard, J. S.; Cleland, W. W. *Biochemistry* **1980**, *19*, 4506–4513.
(53) Bahnsen, B. J.; Anderson, V. E. *Biochemistry* **1989**, *28*, 4173–4181.
(54) Gorenstein, D. G.; Lee, Y. G.; Kar, D. *J. Am. Chem. Soc.* **1977**, *99*, 2264–2267.




Figure 3. KIEs for AP-catalyzed reactions are different from the corresponding solution reaction. (A) Predicted direction of changes in KIEs relative to the solution reaction for two different models for behavior of phosphate monoester substrates in the AP active site. In model 1, the transition state is loose, similar to that in solution, and strong interactions with active-site metal ions lead to decreases in both $^{18}(V/K)_{\text{bridge}}$ and $^{18}(V/K)_{\text{nonbridge}}$. In model 2, AP catalyzes phosphate monoester hydrolysis through a tighter transition state than that in solution. This model predicts a decrease in $^{18}(V/K)_{\text{bridge}}$ and an increase in $^{18}(V/K)_{\text{nonbridge}}$ based on comparisons to corresponding solution reactions.¹⁶ (B) The observed values of both $^{18}(V/K)_{\text{bridge}}$ and $^{18}(V/K)_{\text{nonbridge}}$ for AP-catalyzed pNPP²⁻ hydrolysis decrease relative to the corresponding solution reaction, consistent with model 2. Isotope effects are expressed as a percentage difference from unity [KIE% = (KIE_{obs} - 1) × 100]. Values for the nonenzymatic reaction are shown in black, and values for the AP-catalyzed reaction are shown in gray.

group bond cleavage is likely to be in the range of ~1.04 to 1.06, significantly larger than $^{18}k_{\text{bridge}}$ for nonenzymatic pNPP²⁻ hydrolysis.

For R166S AP-catalyzed mNBP²⁻ hydrolysis, the value of $^{18}(V/K)_{\text{bridge}}$ is 1.0199. While this value is larger than the value of 1.0091 observed for R166S AP-catalyzed pNPP²⁻ hydrolysis, it represents a significant decrease from the estimated value of ~1.04 to 1.06 for the mNBP²⁻ hydrolysis in solution. As for pNPP²⁻, the reduced value of $^{18}(V/K)_{\text{bridge}}$ for the enzymatic reaction relative to the solution reaction probably reflects an inverse contribution to the observed KIE from interactions with a Zn²⁺ ion in the transition state (Figure 3).

The value of $^{18}(V/K)_{\text{nonbridge}}$ for R166S AP-catalyzed mNBP²⁻ hydrolysis is 0.9933, identical within error to the value of 0.9925 observed for R166S AP-catalyzed pNPP²⁻ hydrolysis. This result is consistent with the expectation that the values of

$^{18}k_{\text{nonbridge}}$ for the solution reactions of pNPP²⁻ and mNBP²⁻ should be similar. Moreover, the value of 0.9933 is significantly more inverse than the expected value in solution and may reflect an inverse contribution to the KIE from interactions with Zn²⁺ ions in the transition state, as noted above for the pNPP²⁻ reaction (Figure 3).

The Effect of Metal-Ion Coordination on KIEs. The data for AP-catalyzed reactions described herein suggest that coordination to metal ions can lead to stiffening of the vibrational environment of phosphoryl oxygen atoms and an inverse contribution to the observed KIEs.⁵⁵ It is well-established that coordination of oxygen to divalent metal ions can give rise to inverse isotope effects both in solution^{56,57} and in enzymatic reactions.⁵⁸ Moreover, vibrational spectroscopy studies have demonstrated that metal-ion coordination to phosphoryl oxygen atoms can lead to significant perturbations in the frequencies

of vibrational modes involving phosphoryl groups.^{59–62} This model does not require metal-ion coordination to lead to an increase in P–O bond order and a corresponding increase in the frequency of the P–O stretching vibration. Rather, the inverse contribution to the KIE due to metal-ion coordination arises from a stiffening of the overall vibrational environment of the phosphoryl oxygen atoms, and this stiffening can involve change in stretching, bending, and torsional vibrational modes.

While literature precedents support the idea that coordination to metal ions in the AP active site leads to inverse contributions to the observed KIEs, such effects may not be a general consequence of metal-ion coordination. For example, no significant isotope effect was observed for coordination of Mg^{2+} to inorganic phosphate in solution.⁶³ Suppression of the bridging isotope effect [$^{18}(\text{V}/\text{K})_{\text{bridge}}$] because of metal-ion interactions has not been previously reported in other systems, although it has been suggested as a possible explanation for the observed bridging KIE for λ protein phosphatase.⁶⁴ The available data for the nonbridging KIEs [$^{18}(\text{V}/\text{K})_{\text{nonbridge}}$] varies with the system studied. Only small inverse values of $^{18}(\text{V}/\text{K})_{\text{nonbridge}}$ were observed with λ protein phosphatase containing Mn^{2+} or Ca^{2+} in the active site.⁶⁴ However, such studies may not accurately model binding interactions with the bimetallo Zn^{2+} cluster in the AP active site. Furthermore, large inverse values of $^{18}(\text{V}/\text{K})_{\text{nonbridge}}$ were observed for the reactions of pNPP^{2-} with calcineurin containing Mn^{2+} or Mg^{2+} in the active site.^{65,66} From the available literature data, it is reasonable to suggest that the inverse contributions to KIEs observed for AP-catalyzed reactions arise from metal-ion coordination. Perhaps differences in isotope effects will ultimately aid in distinguishing between different possible transition-state coordination geometries in other binuclear metalloenzymes that catalyze phosphoryl transfer reactions.

KIEs for wt and R166S AP-Catalyzed pNPPS^{2-} Hydrolysis. We have also determined KIEs for the AP-catalyzed reaction of pNPPS^{2-} , an aryl phosphorothioate (Figure 1), and the results reinforce the above conclusions. Phosphorothioate monoesters

are analogues of phosphate monoesters with a single sulfur substitution at a nonbridging position (Scheme 3) and react through similar loose transition states, although bond cleavage to the leaving group may be more advanced.^{11,20,67} Comparisons of phosphate and phosphorothioate monoesters have previously provided valuable insights into the AP reaction mechanism.^{11,32,43}

The KIEs for AP-catalyzed pNPPS^{2-} hydrolysis are significantly different from unity (Table 1), confirming previous conclusions that the chemical step is rate-determining for phosphorothioate hydrolysis catalyzed by both wt and R166S AP.^{11,32} Furthermore, the KIEs for the wt and R166S AP-catalyzed reactions are identical within error. Indeed, mutation of Arg166 to Ser has no significant effect on $k_{\text{cat}}/K_{\text{M}}$ for pNPPS^{2-} hydrolysis.⁶⁸ A similar situation was observed for reactions of AP with phosphate diesters, where reactivity is unaffected by mutation of Arg166 to Ser.²² Both phosphate diesters and phosphorothioate monoesters have large functional groups (either a second ester group or a sulfur atom) that would result in unfavorable interactions with a positioned active-site arginine. The absence of rate effects upon mutation of Arg166 suggests that the arginine side chain can readily move away from the active site to allow diester or phosphorothioate binding and reaction. These data are consistent with pNPPS^{2-} binding to the AP active site with a nonbridging oxygen atom coordinated by the two Zn^{+} ions; the alternative in which the sulfur atom coordinates the Zn^{2+} ions is discussed below.

The values of $^{18}(\text{V}/\text{K})_{\text{bridge}}$ are 1.0094 and 1.0098 for wt and R166S AP-catalyzed pNPPS^{2-} hydrolysis, respectively (Table 1). These values are identical within error to that observed for R166S AP-catalyzed pNPP^{2-} hydrolysis, which has the same leaving group, and significantly smaller than the value of 1.0237 for uncatalyzed pNPPS^{2-} hydrolysis.²⁰ As for pNPP^{2-} and mNBP^{2-} , these data support a model in which coordination of the leaving group oxygen to an active-site Zn^{2+} suppresses the observed KIE.

The nonbridge ^{18}O KIE for the uncatalyzed hydrolysis of pNPPS^{2-} in solution is 1.0135.²⁰ This value is significantly larger than the small inverse values typically observed for phosphate monoester hydrolysis (Table 1),¹⁶ even though phosphate monoesters and phosphorothioate monoesters react through similar loose transition states.^{11,20,67} The large normal value of $^{18}k_{\text{nonbridge}}$ for pNPPS^{2-} has been attributed to a dominant contribution from loosening of bending modes in the transition state for phosphorothioate ester hydrolysis.²⁰

The values of $^{18}(\text{V}/\text{K})_{\text{nonbridge}}$ for pNPPS^{2-} hydrolysis catalyzed by wt and R166S AP are 0.9760 and 0.9768, respectively (Table 1). As for pNPP^{2-} and mNBP^{2-} , the enzymatic reaction results in a large inverse contribution to the nonbridging KIEs, consistent with Zn^{2+} coordination to a nonbridging oxygen atom restricting bending modes. However, the inverse contribution from AP to $^{18}(\text{V}/\text{K})_{\text{nonbridge}}$ for pNPPS^{2-} is far larger than that observed for pNPP^{2-} and mNBP^{2-} , and suggests that in addition to fixing the position of a nonbridging oxygen atom, the AP active site may more severely constrain bending and torsional vibrations that involve the large sulfur atom of pNPPS^{2-} .

(55) The value of $^{18}(\text{V}/\text{K})_{\text{nonbridge}}$ for wt AP-catalyzed hydrolysis of glucose 6-phosphate (G6P) has been reported to be 0.9994.⁴⁸ On the basis of the results reported herein, we would have expected a large inverse value of $^{18}(\text{V}/\text{K})_{\text{nonbridge}}$ for G6P hydrolysis similar to the values observed for pNPP^{2-} and mNBP^{2-} . However, it is possible that the chemical step is not rate-determining for AP-catalyzed G6P hydrolysis. The chemical step is rate-determining for simple alkyl phosphates with leaving group $\text{p}K_{\text{a}}$ values > 14 (Figure 2),¹² but mNBP^{2-} (leaving group $\text{p}K_{\text{a}}$, 14.9) reacts faster than expected, close to the limit at which a nonchemical step becomes rate determining (Figure 2). We have suggested above that the increased reactivity of mNBP^{2-} is due to additional active-site binding interactions with the large benzyl leaving group. G6P has a similarly large leaving group and may also benefit from active-site interactions that lead to a faster-than-expected reaction rate that is limited by a nonchemical step. The $\text{p}K_{\text{a}}$ of glucose at the 6-OH position has not been measured, but it is likely to be lower than 16, the value for ethanol.³⁴ because the neighboring hydroxyl group can provide an intramolecular hydrogen bond.

(56) Feder, H. M.; Taube, H. *J. Chem. Phys.* **1952**, *20*, 1335–1336.

(57) Taube, H. *J. Phys. Chem.* **1954**, *58*, 523–528.

(58) Cassano, A. G.; Anderson, V. E.; Harris, M. E. *Biochemistry* **2004**, *43*, 10547–10559.

(59) Nelson, B. N.; Exarhos, G. J. *J. Chem. Phys.* **1979**, *71*, 2739–2747.

(60) Stangret, J.; Savoie, R. *Can. J. Chem.* **1992**, *70*, 2875–2883.

(61) de la Fuente, M.; Hernanz, A.; Navarro, R. *J. Biol. Inorg. Chem.* **2004**, *9*, 973–986.

(62) Koleva, V. G. *Spectrochim. Acta A* **2007**, *66*, 413–418.

(63) Jones, J. P.; Weiss, P. M.; Cleland, W. W. *Biochemistry* **1991**, *30*, 3634–3639.

(64) Hoff, R. H.; Mertz, P.; Rusnak, F.; Hengge, A. C. *J. Am. Chem. Soc.* **1999**, *121*, 6382–6390.

(65) Hengge, A. C.; Martin, B. L. *Biochemistry* **1997**, *36*, 10185–10191.

(66) Martin, B. L.; Jurado, L. A.; Hengge, A. C. *Biochemistry* **1999**, *38*, 3386–3392.

(67) Hengge, A. C., Transfer of the PO_3^{2-} group. In *Comprehensive Biological Catalysis*; Sinnott, M. L., Ed.; Academic Press: London, 1998; pp 517–542.

(68) O'Brien, P. J.; Herschlag, D. Unpublished data.

An alternative possibility is that pNPPS²⁻ binds to the AP site with the sulfur atom coordinated by the two Zn²⁺ ions, which would lead to a different interpretation of the observed KIEs. However, the larger size and correspondingly longer bond lengths of sulfur relative to oxygen⁶⁹ suggest that sulfur coordination to the active-site Zn²⁺ ions would require significant rearrangements of the active site, including perturbations of Zn²⁺ interactions with the nucleophile and leaving group. If significant rearrangements of the active-site coordination environment were to occur, then the values of ¹⁸(V/K)_{bridge} for pNPPS²⁻ and pNPP²⁻ would be expected to differ, as this KIE reflects interactions of the leaving group with a Zn²⁺ ion. Instead, the values of ¹⁸(V/K)_{bridge} for pNPPS²⁻ and pNPP²⁻ are the same within error when the chemical step is rate-determining (Table 1). Further, structural studies of Klenow exonuclease, which has a bimetallo active site similar to that of AP, suggest that the metal ions are displaced from the active site upon the binding of a phosphorothioate substrate in which the sulfur atom is oriented toward the bimetallo site by remote binding interactions.⁷⁰ Thus, it seems likely that pNPPS²⁻ binds the AP active site with a nonbridging oxygen atom coordinated by the two Zn²⁺ ions.

Implications for Catalysis. The KIEs observed for AP-catalyzed reactions are consistent with previous conclusions from LFER experiments that the transition state for AP-catalyzed phosphate monoester hydrolysis is loose and similar to that in solution. Furthermore, the KIEs suggest that strong interactions with the active-site Zn²⁺ ions restrict vibrational motions of the substrate. These interactions occur at the position of bond cleavage, where extensive charge buildup is expected in a loose transition state, and at the nonbridging positions (Figure 3). Previous functional studies have implicated electrostatic interactions between a nonbridging phosphate ester oxygen atom and the active-site Zn²⁺ ions as being especially important for catalysis,^{43,47} and the KIE data described herein strengthen these conclusions.

AP catalyzes reactions of substrates with poor leaving groups with significantly greater catalytic proficiency than those with good leaving groups.⁷¹ The difference in catalytic proficiency is reflected in the difference between the solution and enzymatic values of β_{lg} , which are -1.23 and -0.85 , respectively.^{12,28} This difference corresponds to smaller catalytic proficiencies as the leaving group pK_a decreases (i.e., for a given decrease in the leaving group pK_a, the rate of the solution reaction increases more than the rate of the corresponding enzymatic reaction). A significant fraction of the difference in β_{lg} between the solution and enzymatic reaction has been suggested to arise from interactions with an active-site Zn²⁺ ion;^{11–13} that is, the large difference in catalytic proficiency could arise from competing effects between stabilization of charge buildup by the leaving group itself and stabilization by interaction with the active-site Zn²⁺. These stabilizing effects are not simply additive such that a leaving group that is better able to stabilize negative charge buildup would consequently interact more weakly with the active-site Zn²⁺ than a poor leaving group.

The KIE data for pNPP²⁻ and mNBP²⁻ suggest that there are strong interactions between the leaving-group oxygen and

an active-site Zn²⁺ ion, and it is possible that differences in the strength of this interaction could be responsible for the differing catalytic proficiencies observed for alkyl and aryl substrates. More direct support for this model could be provided if the decrease in the value of ¹⁸(V/K)_{bridge} relative to that in solution was larger for mNBP²⁻ than for pNPP²⁻. However, this comparison cannot be made because of the absence of experimental KIE data for the mNBP²⁻ reaction in solution.

Mechanistic studies of other enzymes support the idea that differences in catalytic proficiency for good and poor leaving groups could arise from differential interactions with active-site functional groups. The clearest evidence for this phenomenon has been obtained with glycosyl and phosphoryl transfer enzymes that utilize general acid catalysis.^{72–77} For these enzymes, mutation of the residue that acts as the proton donor has a larger deleterious effect for the reactions of substrates with poor leaving groups; that is, substrates with high pK_a leaving groups are more greatly impeded by the mutation than substrates with low pK_a leaving groups. These observations suggest that general acid catalysis makes a larger contribution to catalytic proficiency for substrates with poor leaving groups than for those with good leaving groups, as might be expected on the basis of differential proton affinities.

Enzymes that catalyze phosphoryl transfer reactions are generally observed to have general acids, divalent metal ions, or other charged groups positioned to stabilize negative charge buildup.^{67,78,79} However, it is difficult to assign a precise contribution to catalysis from specific active-site interactions, as perturbing features of the active site by mutagenesis often leads to disruption of multiple interactions and thus cannot be interpreted as evidence for the importance of a particular interaction.¹⁷ The KIE approach circumvents this limitation of site-directed mutagenesis by introducing the smallest possible perturbation to a specific site in the substrate and provides information on both the nature of the transition state and interactions between the active site and the transition state. The data support previous models suggested from extensive structural and functional data that the active-site Zn²⁺ ions in AP form strong interactions with oxygen atoms of the phosphate ester in the transition state.^{7–9,12,43,47,80} Nevertheless, challenges remain in quantitatively describing how the various active-site interactions contribute to catalysis and understanding how these interactions work together to lead to the enormous rate enhancements achieved by this and related enzymes.

Acknowledgment. This work was supported by a grant to D.H. from the NIH (Grant GM64798) and to A.H. from the

(69) Shannon, R. D. *Acta Crystallogr.* **1976**, A32, 751–767.
 (70) Brautigam, C. A.; Steitz, T. A. *J. Mol. Biol.* **1998**, 277, 363–377.
 (71) The overall difference in catalytic proficiency between pNPP²⁻ and methyl phosphate is >10⁸-fold.³³

(72) Richard, J. P.; Huber, R. E.; Lin, S.; Heo, C.; Amyes, T. L. *Biochemistry* **1996**, 35, 12377–12386.
 (73) Thompson, J. E.; Raines, R. T. *J. Am. Chem. Soc.* **1994**, 116, 5467–5468.
 (74) Hondal, R. J.; Bruzik, K. S.; Zhao, Z.; Tsai, M. D. *J. Am. Chem. Soc.* **1997**, 119, 5477–5478.
 (75) Hondal, R. J.; Zhao, Z.; Kravchuk, A. V.; Liao, H.; Riddle, S. R.; Yue, X.; Bruzik, K. S.; Tsai, M. D. *Biochemistry* **1998**, 37, 4568–4580.
 (76) Ostanin, K.; Van Etten, R. L. *J. Biol. Chem.* **1993**, 268, 20778–20784.
 (77) Whiteson, K. L.; Chen, Y.; Chopra, N.; Raymond, A. C.; Rice, P. A. *Chem. Biol.* **2007**, 14, 121–129.
 (78) Knowles, J. R. *Ann. Rev. Biochem.* **1980**, 49, 877–919.
 (79) Maegley, K. A.; Admiraal, S. J.; Herschlag, D. *Proc. Natl. Acad. Sci. U.S.A.* **1996**, 93, 8160–8166.
 (80) Murphy, J. E.; Stec, B.; Ma, L.; Kantrowitz, E. R. *Nat. Struct. Biol.* **1997**, 4, 618–622.
 (81) Grzyska, P. K.; Czyryca, P. G.; Purcell, J.; Hengge, A. C. *J. Am. Chem. Soc.* **2007**, 129, 5298.
 (82) Czyryca, P. G.; Hengge, A. C. *Biochim. Biophys. Acta* **2001**, 1547, 245–253.
 (83) Frey, P. A.; Sammons, R. D. *Science* **1985**, 228, 541–545.

NIH (Grant GM47297). J.G.Z. was supported in part by a Hertz Foundation Graduate Fellowship. We thank members of the Hengge and Herschlag labs for comments on the manuscript, Helen Wiersma for providing additional purified alkaline phosphatase for kinetic measurements, and an anonymous reviewer for helpful comments and suggestions.

Supporting Information Available: The pH rate profile for the reaction of mNBP with R166S AP and comparisons with previous data for AP-catalyzed reactions of pNPP and ethyl phosphate. This material is available free of charge via the Internet at <http://pubs.acs.org>.

JA072196+