




Geometry B Solutions

- Find 100 times the area of a regular dodecagon inscribed in a unit circle. Round your answer to the nearest integer if necessary.



Solution. 300. The area is:

$$12 \times \frac{1}{2} \times 1^2 \times \sin 30^\circ = 3$$

- A triangle has sides of lengths 5, 6, 7. What is 60 times the square of the radius of the inscribed circle?

Solution. 160.


Use Heron's Formula, and then the area of the triangle is

$$\sqrt{9 \times (9 - 5) \times (9 - 6) \times (9 - 7)} = 6\sqrt{6}$$

Hence radius of incircle is

$$\frac{2 \times 6\sqrt{6}}{5 + 6 + 7} = \frac{2\sqrt{6}}{3}$$

- A rectangular piece of paper $ABCD$ has sides of lengths $AB = 1$, $BC = 2$. The rectangle is folded in half such that AD coincides with BC and EF is the folding line. Then fold the paper along a line BM such that the corner A falls on line EF . How large, in degrees, is $\angle ABM$?





Solution. 30. Construct $NG \perp BC$ at G . Triangles ABM , NBM , NBG are all congruent. Hence $\angle ABM = 30^\circ$


4. Tetrahedron $ABCD$ has sides of lengths, in increasing order, 7, 13, 18, 27, 36, 41. If $AB = 41$, then what is the length of CD ?

Solution. 13. By triangle inequality, $AB + DB > 41$, and $AC + CB > 41$. Hence, one of the pairs AD , DB and $\{AC, CB\}$ must be $\{18, 27\}$, the other pair contains 36. WLOG, let $AC = 27$, $CB = 18$. Then $DB \neq 36$, otherwise, $CD > 18$. Hence $AD = 36$, $CD = 13$.


5. A polygon is called concave if it has at least one angle strictly greater than 180° . What is the maximum number of symmetries that an 11-sided concave polygon can have?

Solution. 1. An 11-gon can have only axes passing through a vertex and an opposite side. If it had exactly 2 such axes, they would have to be perpendicular and the polygon would have an even number of sides. If it had exactly 3 axes, each would have to be at an angle of 60° with the next one, and the number of sides of the polygon would be a multiple of 3. Since our polygon has a concave angle, any number of axes of symmetry above 3 would imply at least 12 sides.

6. In the following diagram (not to scale), A, B, C, D are four consecutive vertices of an 18-sided regular polygon with center O . Let P be the midpoint of AC and Q be the midpoint of DO . Find $\angle OPQ$ in degrees.



Solution. 30. Let R be the midpoint of AO . Connect RP , RQ . Then $RP = RO = OQ = RQ$. Hence triangle PQR is isosceles, then some simple calculation yields $\angle OPQ = \angle RPQ - \angle RPO = 50^\circ - 20^\circ = 30^\circ$.



7. Lines l and m are perpendicular. Line l partitions a convex polygon into two parts of equal area, and partitions the projection of the polygon onto m into two line segments of length a and b respectively. Determine the maximum value of $\lfloor \frac{1000a}{b} \rfloor$. (The floor notation $\lfloor x \rfloor$ denotes largest integer not exceeding x)

Solution. 2414. The greatest possible value of the ratio is $(1 + \sqrt{2})$. Let A and B be vertices of the convex polygon on different sides of l so that their distance from l is maximal on each side. Let K and L be the intersections of l with the sides of the polygon. Define the points K_1 and L_1 on extension of AK , AL respectively such that K_1L_1 is parallel to l . Since the polygon is convex, the part of the polygon on A 's side contains AKL , and the part of the polygon on B 's side is contained in K_1KLL_1 . Therefore $[K_1KLL_1] \geq [AKL]$. Let M , N be foot of perpendicular from A , B to line l respectively, then

$$\begin{aligned} \frac{[AKL]}{[AK_1L_1]} &\leq \frac{1}{2} \implies \frac{AM}{AM + BN} \leq \frac{1}{\sqrt{2}} \\ &\implies \frac{AM}{BN} \leq \frac{1}{\sqrt{2} - 1} = 1 + \sqrt{2} \end{aligned}$$

Equality is obtained when the polygon is a triangle with l parallel to one side.

8. Consider the solid with 4 triangles and 4 regular hexagons as faces, where each triangle borders 3 hexagons, and all the sides are of length 1. Compute the *square* of the volume of the solid. Express your result in reduced fraction and concatenate the numerator with the denominator (e.g., if you think that the square is $\frac{1734}{274}$, then you would submit 1734274).

Solution. 52972. Extend the edges that are common to two hexagons. We obtain a regular tetrahedron of side length 3. Hence the volume of original solid is a regular tetrahedron of side length 3 minus volume of 4 regular tetrahedrons of side length 1. The volume is

$$\frac{1}{3} \times \frac{9\sqrt{3}}{4} \times \sqrt{6} \times \frac{27 - 4}{27} = \frac{23\sqrt{2}}{12}$$