

## **Dry Forest Restoration and Unassisted Native Tree Seedling Recruitment at Auwahi, Maui**

Author(s): A. C. Medeiros , E. I. von Allmen , and C. G. Chimera

Source: Pacific Science, 68(1):33-45. 2014.

Published By: University of Hawai'i Press

DOI: <http://dx.doi.org/10.2984/68.1.3>

URL: <http://www.bioone.org/doi/full/10.2984/68.1.3>

---

BioOne ([www.bioone.org](http://www.bioone.org)) is a nonprofit, online aggregation of core research in the biological, ecological, and environmental sciences. BioOne provides a sustainable online platform for over 170 journals and books published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/page/terms\\_of\\_use](http://www.bioone.org/page/terms_of_use).

Usage of BioOne content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

# Dry Forest Restoration and Unassisted Native Tree Seedling Recruitment at Auwahi, Maui<sup>1</sup>

A. C. Medeiros,<sup>2,4</sup> E. I. von Allmen,<sup>3</sup> and C. G. Chimera<sup>3</sup>

**Abstract:** Efforts to restore highly degraded but biologically significant forests draw from a limited toolbox. With less than 10% of their former distribution remaining, Hawaiian dry forests, though critically endangered, remain important biological and cultural refugia. At restoration onset (1997), vegetation of restoration and control areas of degraded Auwahi dry forest, Maui Island, was similar, dominated by nonnative graminoids (restoration 78.3%; control 75.4%), especially *Cenchrus (Pennisetum) clandestinus*. In 2012, unrestored control area vegetation was basically unchanged. In contrast, in the restoration area in 2012, native shrub cover increased from 3.1% to 81.9%, and cover of nonnative graminoids declined from 75.4% to 3.3%. In 2012, nonplanted seedlings of 14 of 22 native tree species and six of seven native shrub species were observed in restoration plots; the majority (99%) were five native (*Dodonaea viscosa*, *Coprosma foliosa*, *Osteomeles anthyllidifolia*, *Chamaesyce celastoides*, *Nestegis sandwicensis*) and one nonnative species (*Bocconia frutescens*). By 2012, stem counts of native woody plants had increased from 12.4 to 135.0/100 m<sup>2</sup>, and native species diversity increased from 2.4 to 6.6/100 m<sup>2</sup>. By 2012, seven rare dry forest tree species, *Charpentiera obovata*, *Nothocestrum latifolium*, *Ochrosia haleakalae*, *Pleomele auwahiensis*, *Santalum ellipticum*, *S. haleakalae*, and *Streblus pendulinus*, had established seedlings and/or saplings within the restoration site, especially notable because natural reproduction is largely lacking elsewhere. Without development and implementation of appropriate management strategies, remaining Hawaiian dry forest will likely disappear within the next century. Multicomponent restoration incorporating ungulate exclusion, weed control, and outplanting as described here offers one strategy to conserve and restore tracts of high-value but degraded forests.

DISPROPORTIONATELY impacted by grazing, wildfire, and displacement by agriculture and human settlements, tropical dry forests are globally among the most threatened of eco-

systems (Janzen 1988, Olson and Dinerstein 2002, Miles et al. 2006). In the Hawaiian archipelago, dry forests, though diverse, are among the most fragmented, reduced, and ecologically degraded ecosystems, with less than 10% of their original pre-Polynesian contact area remaining (Bruegmann 1996). Loss and degradation of the patchwork of Hawaiian dry forest types that formerly occurred on leeward slopes to 1,500 m elevation began with more frequent fires associated with agriculture and *Rattus exulans* (Pacific rat) introduction associated with colonizing Polynesians ca. AD 1000–1200 (Athens 2009, Wilmshurst et al. 2011). Following European contact in 1778, the proliferation of nonnative species, particularly feral and domestic ungulates, fire-adapted grasses, and additional rodent species accelerated forest decline

---

<sup>1</sup> Manuscript accepted 22 May 2013. Project funding provided by the U.S. Fish and Wildlife Service.

<sup>2</sup> U.S. Geological Survey, Pacific Island Ecosystems Research Center, P.O. Box 369, Makawao, Hawai'i 96768.

<sup>3</sup> Pacific Cooperative Studies Unit, University of Hawai'i, Honolulu, Hawai'i 96822.

<sup>4</sup> Corresponding author (e-mail: artcmedeiros@gmail.com).

(Cuddihy and Stone 1990, D'Antonio and Vitousek 1992, Blackmore and Vitousek 2000, Cordell and Sandquist 2008). Though largely undocumented, introduced invertebrates and pathogens have undoubtedly reduced fitness of dry forest species, exemplified recently by the devastating impacts on the keystone dry forest tree *Erythrina sandwicensis* Degener by the invasive African gall-forming wasp *Quadrastichus erythrinae* (Rubinoff et al. 2010).

Despite their degraded state, Hawaiian dry forests remain important natural refugia harboring high numbers of threatened species (Rock 1913), including over 25% of Federally listed Endangered Hawaiian plant species (Cabin et al. 2002). Auwahi dry forest on Maui Island, the study site reported here, was previously known to be among the most diverse of Hawaiian ecosystems (Rock 1913). Currently, Auwahi has 13 species with World Conservation Union (IUCN) Red List status, exceeding New Caledonia dry forests, considered among the world's most endangered tropical dry forests (Gillespie and Jaffre 2003) (see Table 1). Culturally, dry forests are highly valued by native Hawaiians for ethnobotanical source materials, especially prized durable hardwoods for tools and weapons and species with utilitarian, medicinal, or religious significance (Medeiros et al. 1998). Despite being recognized in global conservation priorities (Olson and Dinerstein 2002), remaining Hawaiian dry forests will likely be lost in the next 50–100 yr unless effective management strategies can be developed to stabilize and restore them.

Beginning in 1845, Auwahi's forest understory was destroyed by cattle grazing and burning (C. Lennox, 1967, unpubl. typescript, Auwahi Forest report, in possession of the authors; S. Erdman, pers. comm.). The native understory was replaced by extensive stands of the invasive shrub *Ageratina adenophora* (Spreng.) King & Robinson that dominated until 1945 when eliminated by a biological control program coupled with severe drought (Medeiros et al. 1986). In response, ranchers planted slips of *Cenchrus* (*Pennisetum*) *clandestinus* (Hochst. ex Chiov.) Morrone [common name and hereafter referred to as kikuyu grass] throughout Auwahi ca. 1950 to enhance

cattle pasturage and reduce erosion (P. Erdman, pers. comm.). By 1965, kikuyu grass had spread extensively, developed rank mats, and was regarded as a primary threat to forest health at Auwahi and responsible for a dramatic decline of native trees (Lennox, 1967, unpubl. report).

Kikuyu grass, native to highland forest margins of central and East Africa at 1,950–2,700 m, is noted for its vigorous vegetative reproduction and allelopathic chemical production (Marais 2001). Though useful as forage in marginal situations, kikuyu grass is also invasive in Hawai'i, California, La Réunion, Australia, New Zealand, and South Africa, and is listed as a noxious weed with prohibited transport by the U.S. Department of Agriculture (Weber 2003). At Auwahi, kikuyu grass rarely produces seeds (S. Erdman, pers. comm.). Because seed-producing kikuyu grass cultivars in Hawai'i were developed or introduced after 1950 (Fukumoto and Lee 2003), the kikuyu grass cultivar at Auwahi is likely the "wild, unimproved type" reported to be largely sterile (Marais 2001).

By the late 1960s, approximately 95% of ca. 4,000 ha of Auwahi dry forest on leeward Haleakalā had been destroyed (Lennox, 1967, unpubl. report). The rarity of the forest type and diverse tree flora (49 species) combined with continued forest decline and the troubling long-term (50–100 yr) failure of native tree reproduction prompted a regional biological inventory to describe Auwahi as a "museum forest" (i.e., a high-diversity forest lacking recruitment) (Medeiros et al. 1986).

In the 1960s, forest protection efforts began at Auwahi by excluding domestic and feral ungulates with fencing, which unfortunately accelerated kikuyu grass growth and increased tree mortality (P. Erdman, pers. comm.). Despite effective kikuyu grass control in the 1990s in Auwahi enclosures with newly developed glyphosate-based herbicides, few native seedlings recruited, tree mortality continued, and minimal conservation benefit was realized.

In 1997, a multiphased restoration effort involving ungulate exclusion, herbicidal control of kikuyu grass mats, and mass planting native nurse shrub *Dodonaea viscosa* Jacq. seed-

TABLE 1  
Native Species of Auwahi Restoration Area

Taxa <sup>a</sup>	Hawaiian Name	Family	Guild	Status <sup>b</sup>
<i>Alectryon macrococcus</i> <sup>3</sup>	<i>Māhoe</i>	Sapindaceae	Tree	P
<i>Alphitonia ponderosa</i>	<i>Kauila</i>	Rhamnaceae	Tree	E, P
<i>Alyxia oliviformis</i>	<i>Maile</i>	Apocynaceae	Vine	E, P
<i>Bidens micranthra</i> subsp. <i>kalealaba</i> <sup>3</sup>	<i>Kōko'olau</i>	Asteraceae	Shrub	P
<i>Carex wahuensis</i>		Cyperaceae	Sedge	E
<i>Chamaesyce celastroides</i> var. <i>lorifolia</i>	<i>ʻAkoko</i>	Euphorbiaceae	Tree	E, P
<i>Charpentiera obovata</i>	<i>Pāpala</i>	Amaranthaceae	Tree	E, P
<i>Claoxylon sandwicense</i>	<i>Po'olā</i>	Euphorbiaceae	Shrub	P
<i>Cocculus trilobus</i>	<i>Huebue</i>	Menispermaceae	Vine	E
<i>Coprosma foliosa</i>	<i>Pilo</i>	Rubiaceae	Shrub	E, P
<i>Diospyros sandwicensis</i>	<i>Lama</i>	Droseraceae	Tree	E
<i>Dodonaea viscosa</i>	<i>ʻAʻaliʻi</i>	Sapindaceae	Shrub	E, P
<i>Leptocophylla tameiameia</i>	<i>Pūkiawe</i>	Epacridaceae	Shrub	E
<i>Mariscus billebrandii</i>		Cyperaceae	Sedge	E
<i>Melicope adscendens</i> <sup>3</sup>	<i>Alani</i>	Rutaceae	Shrub	E, P
<i>Metrosideros polymorpha</i>	<i>ʻŌhiʻa</i>	Myrtaceae	Tree	E
<i>Myoporum sandwicense</i>	<i>Naio</i>	Myoporaceae	Tree	E, P
<i>Myrsine lanaiensis</i>	<i>Kōlea</i>	Myrsinaceae	Tree	E, P
<i>Myrsine lessertiana</i>	<i>Kōlea lau nui</i>	Myrsinaceae	Tree	E, P
<i>Nestegis sandwicensis</i>	<i>Olopua</i>	Oleaceae	Tree	E
<i>Nothocestrum latifolium</i> <sup>1,4</sup>	<i>ʻAiea</i>	Solanaceae	Tree	E, P
<i>Ocrosia baleakalae</i> <sup>1,4</sup>	<i>Hōlei</i>	Apocynaceae	Tree	E, P
<i>Osteomeles anthyllidifolia</i>	<i>ʻŪlei</i>	Rosaceae	Shrub	E, P
<i>Panicum tenuifolium</i>	<i>Konakona</i>	Poaceae	Grass	P
<i>Peperomia blanda</i>	<i>ʻAlaʻala wai nui</i>	Piperaceae	Herb	E
<i>Pipturus albidus</i>	<i>Māmaki</i>	Urticaceae	Shrub	E
<i>Pisonia brunoniana</i>	<i>Pāpala kēpau</i>	Nyctaginaceae	Tree	P
<i>Planchonella sandwicensis</i>	<i>ʻAlaʻa</i>	Sapotaceae	Tree	E
<i>Pleomele auwahiensis</i> <sup>2</sup>	<i>Hala pepe</i>	Agavaceae	Tree	E, P
<i>Polyscias oahuensis</i>	<i>ʻOhe mauka</i>	Araliaceae	Tree	E
<i>Santalum ellipticum</i>	<i>ʻIiabioloʻe</i>	Santalaceae	Tree	E, P
<i>Santalum baleakalae</i> var. <i>lanaiensis</i> <sup>2,3</sup>	<i>ʻIiabi</i>	Santalaceae	Tree	E, P
<i>Sicyos pachycarpus</i>	<i>ʻAnunu</i>	Cucurbitaceae	Vine	E
<i>Sisyrinchium acre</i>	<i>Mauʻu bōʻulā ʻili</i>	Iridaceae	Herb	P
<i>Sophora chrysophylla</i>	<i>Māmame</i>	Fabaceae	Tree	E, P
<i>Streblus pendulinus</i>	<i>Aʻiaʻi</i>	Moraceae	Tree	E, P
<i>Vigna o-wahuensis</i> <sup>3</sup>		Fabaceae	Vine	P
<i>Wikstroemia monticola</i>	<i>ʻĀkia</i>	Rubiaceae	Tree	E
<i>Xylosma hawaiiense</i>	<i>Maua</i>	Flacourtiaceae	Tree	P
<i>Zanthoxylum hawaiiense</i> <sup>1,3</sup>	<i>Aʻe</i>	Rutaceae	Tree	P

<sup>a</sup> Endangerment status: <sup>1</sup>, Endangered IUCN Red List; <sup>2</sup>, Vulnerable IUCN Red List; <sup>3</sup>, Endangered USFWS; <sup>4</sup>, Candidate USFWS.

<sup>b</sup> E, extant within the restoration area; P, planted.

lings was initiated on a 4 ha tract of relict dry forest at Auwahi. Nurse planting involves outplanting selected nurse-plant species around target species, usually with the goal of stimulating target species seedling recruitment (Gomez-Aparicio 2009). Potential benefits associated with nurse plantings include sun and wind moderation, cooler and moister

soils, increased perch trees for seed-dispersing birds, and perhaps improved nutrient cycling and mycorrhizae (Padilla and Pugnaire 2006), as well as recovered hydrological functioning (Perkins et al. 2012). The primary objective of this paper was evaluation of a multiphased restoration technique in a highly degraded Hawaiian dry forest.

## MATERIALS AND METHODS

*Study Site*

Auwahi forest is located at 1,160–1,250 m elevation in Auwahi district on leeward flanks of Haleakalā volcano on 3,000- to 5,000-yr-old lava (20° 38' 24" N, 156° 20' 24" W) on privately owned 'Ulupalakua Ranch, Maui, Hawaiian Islands. The restoration area is 4 ha of relictual forest where an ungulate-proof enclosure was constructed with U.S. Fish and Wildlife Service (USFWS) funds. The terrain consists of a series of rocky ridges and interconnecting gullies (slope 20–30 degrees) with generally rocky substrate and sparse soil accumulations. Mean annual precipitation is ca. 730 mm with a pronounced dry season from April to September (Giambelluca et al. 2011) and mean monthly temperatures between 13.9°C and 18.3°C (Scholl et al. 2007).

*Restoration Methodology*

In 1997, restoration was initiated with construction of a perimeter fence 1.3 m high excluding domestic cattle and feral ungulates from the 4 ha site. Kikuyu grass mats were suppressed with one primary herbicide treatment (ca. 1.5% glyphosate) and one follow-up treatment several months later to treat resprouts. Treated grass mats were not removed but left to decay. Seedlings of the native *Dodonaea* (ca. 0.25 m height) were grown in tree planting tubes and planted at high densities (ca. 1 m spacing) by community volunteers. *Dodonaea* was selected as the primary nurse plant for its ease of propagation, hardiness, rapid growth, quick time to reproduction, and its historical presence as a primary component of the original understory vegetation (Lennox, 1967, unpubl. report). *Dodonaea* shrubs were planted in open areas where mats of kikuyu grass formerly occurred throughout the restoration site. Plantings of seedlings of other depleted native species (Table 1), also in tree planting tubes, were generally clustered to maximize particular habitat types or facilitate future outcrossing pollination.

*Vegetation Sampling*

To initially compare control and restoration areas, four randomly located 100 m transects were established in each area and vegetation sampled with point- and line-intercept in both 1997 and 2012. Understory vegetation < 0.5 m height was sampled with point-intercept and canopy shrub and tree species > 0.5 m height were recorded with line-intercept. In the control area, 68 100 m<sup>2</sup> plots were randomly located and woody species were counted and classified by basal diameter in 2012. In addition, to more closely track vegetation trajectories in the restoration area, 84 100 m<sup>2</sup> plots were randomly located within the restoration site and the number of stems for each woody species per plot recorded in size classes based on basal diameter. In a randomly selected subset (31 of 84 100 m<sup>2</sup> plots), estimates of cover were made visually to the nearest 5% (or to nearest 1% when cover was < 5%) for all vascular plant species. Count plots and cover estimates of species were assessed in 1997 before restoration and again in 2012. Methods follow Mueller-Dombois and Ellenberg (2003).

*Statistical Analyses*

This study is not an investigation of the efficacy of restoration treatments per se, but rather a comparison of spatiotemporal floristic differences between two specific areas within one site (inside versus outside the restoration area) over a 15-yr period. The restoration site was chosen with the landowner and funder to provide protection and attempt restoration of a particularly biologically important forest tract of an endangered plant community. From an experimental perspective, it would have been preferable to have multiple fenced and unfenced sites to evaluate multiple restoration treatments, but the rarity and conservation value of the studied forest type, and the lack of comparable unprotected areas, precluded such an experimental design. In essence, therefore, the study is pseudoreplicated, with only one restoration site that received a series of synchronized treatments (ungulate exclusion, herbicidal applications, and native

nurse-shrub planting) and one adjacent untreated area. Paired *t*-tests compared differences in cover of major vegetation categories and species as well as mean seedling recruitment within restoration site plots between 1997 and 2012. Mean differences in seedling recruitment between control plots and restoration site plots were compared with *t*-tests in 2012 only. Bonferroni corrections were used to adjust for multiple *t*-tests. Transformations were applied to cover data (arcsin) and seedling number data (square root + 0.05) to improve normality and meet assumptions of parametric statistical tests (Zar 1999). Data analyses were performed using Minitab 15 and Sigmaplot 10.

## RESULTS

### *Restoration versus Control Area in 1997 (Point- and Line-Intercept)*

Initially, vegetation of restoration and control areas was very similar. In 1997, understory vegetation (point-intercept) of both areas was similarly dominated by nonnative graminoids (restoration 78.3% ± 2.8; control 75.4% ± 4.2; *df* = 6; *P* = .27), especially kikuyu grass (restoration 75.4% ± 2.3; control 70.7% ± 4.3; *df* = 6; *P* = .18). Native tree cover (line-intercept) was also similar between restoration and control areas (restoration 5.7% ± 3.0; control 8.6% ± 4.6; *df* = 6; *P* = .69). Cover of native shrubs (line-intercept) was higher in control areas but not significantly so (restoration 3.1% ± 1.2; control 19.7% ± 9.8; *df* = 6; *P* = .20).

### *Control Area 1997–2012 (Point- and Line-Intercept)*

When resampled in 2012, understory vegetation of the control area was basically unchanged from 1997. Nonnative graminoid cover remained dominant (1997: 75.4% ± 4.2 versus 2012: 87.4% ± 4.3; *df* = 3; *P* = .18) with slight increases in kikuyu grass (70.6% ± 4.3 versus 77.6% ± 2.4; *df* = 3; *P* = .30). Native shrub cover (line-intercept) in the control area remained relatively constant (1997: 19.7% ± 9.8 versus 2012: 21.8% ± 13.9;

*df* = 3; *P* = .71), and native tree cover declined slightly (1997: 8.6% ± 4.6 versus 2012: 7.7% ± 2.0; *df* = 3; *P* = .84).

### *Restoration Area 1997–2012 (Point- and Line-Intercept)*

In contrast to control areas, by 2012 nonnative graminoids, especially kikuyu grass, had declined dramatically in the restoration area from 75.4% ± 2.3 to 3.3% ± 2.3 (*df* = 3, *P* < .0001). Correspondingly, cover of native shrubs (line-intercept) increased from 3.1% ± 1.2 to 81.9% ± 9.9 (*df* = 3, *P* < .05), especially *Dodonaea* (+59.5% ± 6.6) but also *Coprosma foliosa* Gray (+10.5% ± 4.2) and *Osteomeles anthyllidifolia* (Sm.) Lindl. (+8.3% ± 3.3). In the restoration area, native tree cover increased from 5.7% ± 3.0 to 18.5% ± 8.4 (*df* = 3, *P* = .32). Cover increases of *Dodonaea*, *Coprosma*, and *Osteomeles* represent both wild and planted individuals (see count plot data), whereas increases in plots of the two depleted native tree species *Pleomele auwahiensis* St. John (+1.2% ± 0.2) and *Xylosma hawaiiense* Seem. (+0.8% ± 0.6) represent planted individuals, surviving after 15 yr.

### *Restoration Area 1997–2012 (Cover Estimates)*

The most significant changes by guild (1997–2012) were decreases in total nonnative cover (–84.3%), especially nonnative graminoids (–73.1%) and nonnative herbs (–11.0%) and increases in total native cover (+57.6%), native shrubs (+59.7%), rock (+27.9%), and leaf litter (+51.5%) (Table 2, Figures 1 and 2). The most significant changes by species were decreases of kikuyu grass (–70.6%) and *Asclepias physocarpa* (Mey.) Schlechter (–11.1%) and increases of *Dodonaea* (+51.9%), *Osteomeles* (+6.2%), and *Coprosma* (+1.7%) (Table 3). Large-scale increases of native shrubs used as nurse plants, especially *Dodonaea*, are due to the growth and spread of planted individuals and subsequent natural recruitment. In cover estimates of *Dodonaea*, 66.2% cover increase was due to planted individuals, and 33.8% was from surviving original individuals and sapling recruitment. Cover of native tree species changed little (–3.1%)

TABLE 2

Mean Percentage Cover (SE) of Guilds within Restoration Site Based on Cover Estimates of 31 Randomly Located 100 m<sup>2</sup> Plots before and after 15 yr of Restoration

Cover Category	% Cover 1997	% Cover 2012
Nonnative grasses**	74.82 (3.90)	1.79 (0.42)
Native shrubs**	8.62 (2.58)	68.04 (3.23)
Nonnative herbs**	12.01 (3.09)	1.06 (0.45)
Native trees	12.17 (3.10)	10.43 (2.62)
Native grasses	0	1.07 (0.51)
Native vines	0.30 (0.06)	0.49 (0.19)
Native herbs	0	2.75 (2.25)
Native sedges	0.18 (0.06)	0.39 (0.18)
Nonnative sedges*	0.33 (0.10)	0.01 (0.003)
Native ferns	0.26 (0.12)	0.27 (0.21)
Nonnative tree	0.77 (0.25)	1.11 (0.55)

\* df = 30,  $P \leq .01$ , \*\*df = 30,  $P \leq .001$ .

according to cover estimates, perhaps reflecting the limitation of the technique of visually assigning cover estimates for tree species in structurally complex vegetation (Table 3).

#### Restoration Area 1997–2012 (Count Plots)

By 2012, seedlings of 14 of 22 native tree species and six of seven native shrub species had been observed in plots within the restoration area. The great majority (99%) of seedlings in plots belonged to six species: three native shrubs (*Dodonaea*, *Coprosma*, *Osteomeles*), two native trees [*Chamaesyce celsoides* (Bois.) Croizat & Degener var. *lorifolia* (Gray) Degener & Degener and *Nestegis sandwicensis* (Gray) Degener, Degener & Johnson], and one invasive tree (*Bocconia frutescens* L.) (Table




FIGURE 1. Mean percentage cover changes between 1997 and 2012 within restoration site based on cover estimates of 31 randomly located 100 m<sup>2</sup> plots before and after 15 yr of restoration. Bars represent standard error. All changes in percentage cover were significant to  $P \leq .001$ .



FIGURE 2. Auwahi restoration site, 2011.

TABLE 3

Mean Percentage Cover (SE) of Selected Species within Restoration Site Based on Cover Estimates of 31 Randomly Located 100 m<sup>2</sup> Plots before and after 15 yr of Restoration

Cover Category <sup>a,b</sup>	% Cover 1997	% Cover 2012
<b><i>Cenchrus clandestinus</i></b> <sup>***</sup>	70.90 (4.10)	0.26 (0.11)
<i>Dodonaea viscosa</i> <sup>***</sup>	3.60 (1.44)	55.55 (3.86)
<b><i>Asclepias physocarpa</i></b> <sup>**</sup>	11.18 (2.90)	0.12 (0.10)
<i>Osteomeles anthyllidifolia</i> <sup>*</sup>	2.64 (0.85)	8.86 (2.28)
<b><i>Coprosma foliosa</i></b> <sup>***</sup>	1.35 (0.62)	3.09 (0.66)
<i>Pleomele auwahiensis</i> <sup>2***</sup>	0.0003 (0.0003)	1.02 (0.24)
<i>Ocrosia haleakalae</i> <sup>1,4*</sup>	0	0.73 (0.35)
<i>Myrsine lessertiana</i>	1.67 (1.07)	1.00 (0.67)
<i>Chamaesyce celastroides</i> <sup>**</sup>	0.02 (0.01)	0.66 (0.24)
<i>Metrosideros polymorpha</i>	0.81 (0.57)	0.76 (0.43)
<i>Santalum ellipticum</i>	0.71 (0.43)	0.36 (0.25)
<i>Cheirodendron trigynum</i>	0.32 (0.32)	0
<i>Charpentiera obovata</i>	0.29 (0.21)	0
<i>Nothocestrum latifolium</i> <sup>1,4</sup>	0	0.26 (0.19)
<i>Polyscias oahuense</i>	0.32 (0.32)	0.10 (0.10)
<i>Xylosma hawaiiense</i>	0	0.19 (0.16)
<i>Santalum haleakalae lanaiensis</i> <sup>2,3</sup>	0	0.03 (0.03)
<i>Zantoxylum hawaiiense</i> <sup>1,3</sup>	0	0.02 (0.01)

<sup>a</sup> Nonnative species are in **boldface**.

<sup>b</sup> Endangerment status: <sup>1</sup>, Endangered IUCN Red List; <sup>2</sup>, Vulnerable IUCN Red List; <sup>3</sup>, Endangered USFWS; <sup>4</sup>, Candidate USFWS.

\* df = 30,  $P \leq .05$ , \*\*df = 30,  $P \leq .01$ , \*\*\*df = 30,  $P \leq .001$ .



TABLE 4

Mean (SE) Numbers of Naturally Occurring Tree Seedlings within 100 m<sup>2</sup> Count Plots inside Restoration Site (1997 and 2012, *n* = 84) and outside as Controls (2012 only, *n* = 68)

Taxa <sup>a,b</sup>	Control ( <i>n</i> = 68)	Restoration Site ( <i>n</i> = 84)	
	2012	1997	2012 <sup>c</sup>
<i>Dodonaea viscosa</i>	1.15 (0.30)	2.26 (0.31)	55.83 (6.28) <sup>*,**</sup>
<i>Coprosma foliosa</i>	0.91 (0.72)	0.74 (0.27)	30.18 (4.50) <sup>*,**</sup>
<i>Chamaecybe celastoides</i>	0	0.10 (0.06)	3.43 (1.10) <sup>*,**</sup>
<i>Osteomeles anthyllidifolia</i>	0.03 (0.02)	0.12 (0.04)	3.24 (0.68) <sup>*,**</sup>
<b><i>Bocconia frutescens</i></b>	0.04 (0.03)	0.23 (0.09)	1.82 (0.41) <sup>*,**</sup>
<i>Nestegis sandwicensis</i>	0	0.08 (0.05)	1.14 (0.29) <sup>*,**</sup>
<i>Pleomele auwahiensis</i> <sup>2</sup>	0	0	0.27 (0.13) <sup>*,**</sup>
<i>Myrsine lanaiense</i>	0	0.02 (0.02)	0.19 (0.14)
<i>M. Jessertiana</i>	0.01 (0.01)	0.036 (0.03)	0.17 (0.08)
<i>Wikstroemia monticola</i>	0	0	0.14 (0.05)
<i>Ochrosia haleakalae</i> <sup>1,4</sup>	0	0.01 (0.01)	0.10 (0.06)
<i>Santalum haleakalae</i> var. <i>lanaiensis</i> <sup>2,3</sup>	0	0	0.06 (0.06)
<i>Nothocestrum latifolium</i> <sup>1,4</sup>	0	0	0.036 (0.02)

<sup>a</sup> Nonnative species in **boldface**.

<sup>b</sup> Endangerment status: <sup>1</sup>, Endangered IUCN Red List; <sup>2</sup>, Vulnerable IUCN Red List; <sup>3</sup>, Endangered USFWS; <sup>4</sup>, Candidate USFWS.

<sup>c</sup> \*, Comparison with 2012 control mean (*t*-test, *df* = 150, *P* < .05); \*\*, comparison with 1997 restoration site mean (paired *t*-test, *df* = 83, *P* < .05).

4). By 2012, stem counts of native woody species had increased by an order of magnitude from  $12.4 \pm 1.4/100 \text{ m}^2$  to  $135.0 \pm 9.7$  (*df* = 83, *P* < .001). In plots, native species diversity increased from  $2.4 \pm 0.2/100 \text{ m}^2$  to  $6.6 \pm 0.2$  (*df* = 83, *P* < .001).

#### DISCUSSION

From 1997 to 2012, ungulate exclusion, weed control, and nurse planting have apparently been important factors in facilitating long-term suppression of formerly dominant invasive kikuyu grass (Table 3); reestablishment of native shrub understory (Table 2, Figure 2); seedling recruitment by some native tree, shrub, herb, grass, and vine species (Table 4); and apparently some degree of biotic resistance to reinvasion by light-demanding non-native flora, particularly fire-adapted grasses (D'Antonio and Vitousek 1992).

The absence of seed set and seed banks of kikuyu grass and resultant sparse recruitment of kikuyu grass seedlings after herbicide ap-

plications appears to have been an important factor in restoration efficacy at Auwahi. Treated kikuyu grass decomposed within ca. 1–2 yr, providing mulch and weed protection while nurse plants established (Medeiros et al. 2003, Medeiros and von Allmen 2006). Survival of outplanted nurse-plant seedlings, especially *Dodonaea*, was very high (>95% [A.C.M., unpubl. data]). For several years after outplanting, growth of nurse plants was rapid, perhaps due to heightened nutrient availability accompanying kikuyu grass decomposition. Within 2 yr, outplanted *Dodonaea* shrubs averaged 1 m<sup>3</sup> in cover and began to interlock canopies. Although no quantitative assessment was made, the thick mulch of decomposing kikuyu grass mats and shading provided by densely planting rapid-growing shrub nurse plants appeared to deter reinvasion by a regional suite of light-loving invasive plant species. Elsewhere, shrub nurse plants have had strongest benefits on target tree species including enhanced germination and seedling survival (Padilla and Pugnaire 2006, Gomez-Aparicio 2009).

Other dry forest restoration projects in Hawai'i have documented that just ungulate exclusion and invasive plant control have been insufficient to spur unassisted seedling recruitment of many dry forest species (Cabin et al. 2000, Cabin et al. 2002, Brooks et al. 2009). At Auwahi, the restoration strategy combining nonnative grass removal, ungulate exclusion, and reestablishment of native shrub understory through nurse planting (Table 2) appears to have recreated safe sites for enhanced native seedling establishment. As of 2012, unassisted seedling recruitment had been observed in the restoration site for 64% of native trees and 86% of native shrub species (Table 4). Following the rainy season, cohorts of hundreds of seedlings of *Dodonaea*, *Coprosma*, and *Chamaecybe* have been observed. In addition to the creation of safe sites, increases in seedling recruitment in some cases may be due to heightened levels of available seed as planted individuals begin to produce fruit.

By 2012, seven rare dry forest tree species, *Charpentiera obovata* Gaud., *Ochrosia haleakalae* St. John, *Pleomele auwahiensis*, *Nothocestrum latifolium* Gray, *Santalum haleakalae* Hillebr. var. *lanaiense* Rock (Harbaugh), *S. ellipticum* Gaud., and *Streblus pendulinus* (Endl.) Muell., had established seedlings and/or saplings within the restoration site. Though the number of observed seedlings was in some cases limited, the recruitment is important in that natural reproduction is largely lacking elsewhere in wild populations. Four of these species have IUCN Red List status; one (*S. haleakalae* var. *lanaiense*) is considered Endangered (USFWS). For example, in 2012, two wild *Nothocestrum latifolium* seedlings were discovered below perch trees distant from mature *Nothocestrum* individuals, apparently the result of seed dispersal by birds, likely the nonnative *Zosterops japonicus* (Japanese White-eye). Though modest, this recruitment represents the only currently known natural regeneration of the species. *Nothocestrum latifolium* is the sole native larval host plant for Blackburn's Sphinx Moth, *Manduca blackburni*, listed as Endangered by the USFWS (Rubinoff and San Jose 2010).

The conditions in the restoration area that promote recruitment of native species appear

also to facilitate establishment of certain nonnative woody species. Although most nonnative species declined dramatically, the invasive neotropical tree *Bocconia frutescens* increased in restoration area count plots. Despite control efforts, *Bocconia* recruitment in the restoration area was significantly higher than the same area in 1997 or adjacent unrestored areas in 2012 (Table 4). As such, weed control in the restoration area (ca. 48 person-hr/yr) is currently devoted predominantly to *Bocconia*. Without these efforts, *Bocconia* would likely increase and dominate portions of the restoration area. These results demonstrate the vulnerability of even relatively restored ecosystems to nonnative woody species such as *Bocconia* with the ability to recruit seedlings in shaded or semishaded sites.

Because success in rare plant reintroductions is mixed (Hobbs 2007), long-term survival and self-sustaining recruitment of reintroduced species in Auwahi is encouraging. Since 1997, nine highly depleted native species were reintroduced. Three survive but have yet to reach maturity: *Alectryon macrococcus* Radlk. var. *auwahiensis* Linney, *Xylosma hawaiiense*, and *Zanthoxylum hawaiiense* Hillebr.; two persist with limited recruitment: *Claoxylon sandwicense* Mull. Arg., *Pisonia brunoniana* Endl.; and one failed: *Sisyrinchium acre* Mann. All original outplantings of two short-lived Endangered (USFWS) species, *Bidens micrantha* Gaud. subsp. *kalealaba* Nagata & Ganders and *Vigna o-wahuensis* Vogel; and a highly depleted endemic grass, *Panicum tenuifolium* Hook. & Arnott, failed to persist. However, the three species became established in the restoration site through repeated cohorts of seedling recruitment, some at considerable distances from original planting sites.

One unfortunate but informative case of selective herbivory of an extremely rare species by a nonnative ungulate involved the vine *Vigna o-wahuensis*, a USFWS Endangered species with a total wild population of fewer than 100 individuals (Anon. 1994). In 2010, two juvenile feral pigs entered the restoration site, their smaller size allowing entry through perimeter fence mesh (since repaired). Before their removal, the pigs apparently searched

for and destroyed the entire outplanted population of approximately 100 established *Vigna* while minimally impacting other plant species. The pigs excavated the plants, consuming all parts including roots. By 2012, 24 newly emerged *Vigna* seedlings were recorded in count plots (all near original plantings).

Without native rodents, the Hawaiian biota largely lacks adaptations deterring high levels of predation on native plants, invertebrates, and birds, especially nesting populations (Drake and Hunt 2009). Rodents are a primary factor limiting seed production and perhaps seedling recruitment of Hawaiian plant species. Hawaiian dry forest trees with rodent-palatable seeds typically suffer near complete loss of seed crop and absence of seed bank (Chimera and Drake 2011), a fate shared with highly depleted species elsewhere in the Pacific (Meyer and Butaud 2009). Though predator-resistant fencing is costly and difficult, long-term rodent control would likely have positive and profound cascading effects on native invertebrate, bird, and plant populations released from predation (Innes et al. 2012).

The impact of nonnative rodents on seeds of certain native trees has been exacerbated by extinctions of native frugivorous birds, restricting “seed shadows” to beneath canopy areas (Foster 2009, Chimera and Drake 2010). Large-seeded (>7 mm) Hawaiian dry forest trees (*Alectryon*, *Nestegis*, *Pleomele*, and *Pouteria*), no longer dispersed by birds such as the extirpated Hawaiian Crow (*Corvus* spp.) capable of processing large seeds, often lose entire seed crops to rodent predation and characteristically lack seed banks (Culliney et al. 2012).

What will be the future trends at the Auwahi restoration site? The outcome clearly depends on climate change, invasive species, future land uses, management priorities, and perhaps other unforeseen factors. Without control of the invasive tree *Bocconia*, it appears likely that this species will continue to be dispersed into and invade the restoration site. More positively, the unassisted recruitment of native tree seedlings will increase the complexity and height of the emerging forest. Community composition will increasingly

reflect native species–native species competition instead of invasive species–native species competition. In some sense, a native version of a novel ecosystem has developed, as proposed by Hobbs et al. (2009, 2013). Mutualistic native–nonnative species interactions, such as pollination and seed dispersal, may further assist in restoring ecosystem functioning, as has been documented elsewhere in Hawai‘i (Cole et al. 1995, Foster and Robinson 2007). One example from the Auwahi restoration site is the apparent dispersal of small-seeded (<7 mm) native species such as *Coprosma*, *Leptecophylla tameiameia* (Cham. & Schldl.) Weller, *Osteomeles*, *Santalum ellipticum*, and *Wikstroemia monticola* Skotts. by nonnative birds, especially the small near-ubiquitous passerine *Zosterops japonicus* and gallinaceous Ring-necked Pheasant (*Phasianus colchicus*).

How transferable are lessons from Auwahi for restoration of other degraded forests? Several somewhat unique site factors may have contributed to efficacy of the restoration protocol at Auwahi. First, ranchlands and natural areas surrounding Auwahi harbor a relatively low diversity of nonnative species (Medeiros et al. 1986). This situation, combined with the absence of seed banks and relatively simple chemical control of kikuyu grass, creates an uncommonly manageable dominant invasive species. Secondly, *Dodonaea*, the nurse plant utilized here and a component of the original forest understory, was readily available, easy to propagate, grew rapidly, and had high outplanting survival.

Without appropriate management strategies, complete conversion of remaining Hawaiian dry forest will likely occur within the next century. The multicomponent restoration effort described here offers one strategy to conserve and restore tracts of dry forests in Hawai‘i and perhaps elsewhere. With climate change and rampant movement of nonnative species, native ecosystems are under siege worldwide, and restoration, already difficult, has become increasingly complicated (Harris et al. 2006). Future management efforts to mitigate climate change in areas such as watersheds may draw from challenges and lessons gleaned from restoration of degraded forests.

## ACKNOWLEDGMENTS

We thank and acknowledge Sumner, Pardee, and Betsy Erdman, ‘Ulupalakua Ranch owners, for their uncommon conservation ethic and support of science. We thank the Pacific Island Ecosystems Research Center and the Ecosystems and Invasives Programs of U.S. Geological Survey for essential support; Robert Cabin, Lloyd Loope, Richard Camp, Kevin Brinck, and two anonymous reviewers for manuscript comments; Andrea Buckman, Ka‘ai Bustamente, Keahi Bustamente, Diana Crow, Tracy Erwin, Stephanie Joe, Fernando Juan, Ainoa Kaiaokamelie, Paul Krushelnycky, Lindsey Manual, Luke McLean, Forest Starr, Kim Starr, and Brianna Welker for fieldwork assistance, and the tireless dedication and devotion of the hundreds of volunteers who gave up their Saturdays helping to plant and restore this unique dry forest community. Any use of trade, product, or firm names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

## Literature Cited

- Anon. 1994. Rules and Regulations, p. 56342, 59 Fed. Reg. 217 (18 November 1994).
- Athens, J. S. 2009. *Rattus exulans* and the catastrophic disappearance of Hawai‘i’s native lowland forest. *Biol. Invasions* 11:1489–1501.
- Blackmore, M., and P. M. Vitousek. 2000. Cattle grazing, forest loss, and fuel loading in a dry forest ecosystem at Pu‘u Wa‘a Wa‘a Ranch, Hawai‘i. *Biotropica* 32:625–632.
- Brooks, S., S. Cordell, and L. Perry. 2009. Broadcast seeding as a potential tool to reestablish native species in degraded dry forest ecosystems in Hawaii. *Ecol. Restor.* 27:300–305.
- Brueggemann, M. M. 1996. Hawaii’s dry forests. *Endang. Species Bull.* 11:26–27.
- Cabin, R. J., S. G. Weller, D. H. Lorence, S. Cordell, L. J. Hadway, R. Montgomery, D. Goo, and A. Urakami. 2002. Effects of light, alien grass, and native species additions on Hawaiian dry forest restoration. *Ecol. Appl.* 12:1595–1610.
- Cabin, R. J., S. G. Weller, D. H. Lorence, T. W. Flynn, A. K. Sakai, D. Sandquist, and L. Hadway. 2000. Effects of long-term ungulate exclusion and recent alien species control on the preservation and restoration of a Hawaiian tropical dry forest. *Conserv. Biol.* 14:439–453.
- Chimera, C. G., and D. R. Drake. 2010. Patterns of seed dispersal and dispersal failure in a Hawaiian dry forest having only introduced birds. *Biotropica* 42:493–502.
- . 2011. Could poor seed dispersal contribute to predation by introduced rodents in a Hawaiian dry forest? *Biol. Invasions* 13:1029–1042.
- Cole, F. R., L. L. Loope, A. C. Medeiros, J. A. Raikes, and C. S. Wood. 1995. Conservation implications of introduced game birds in high-elevation Hawaiian shrubland. *Conserv. Biol.* 9:306–313.
- Cordell, S., and D. R. Sandquist. 2008. The impact of an invasive African bunchgrass (*Pennisetum setaceum*) on water availability and productivity of canopy trees within a tropical dry forest in Hawaii. *Funct. Ecol.* 22:1008–1017.
- Culliney, L. W., S. L. Pejchar, R. Switzer, and V. Ruiz-Gutierrez. 2012. Seed dispersal by a captive corvid: The role of the ‘Alalā (*Corvus hawaiiensis*) in shaping Hawai‘i’s plant communities. *Ecol. Appl.* 22:1718–1732.
- Cuddihy, L. W., and C. P. Stone. 1990. Alteration of native Hawaiian vegetation. Cooperative National Park Resources Studies Unit, University of Hawai‘i at Mānoa, Honolulu.
- D’Antonio, C. M., and P. M. Vitousek. 1992. Biological invasions by exotic grasses, the grass/fire cycle, and global change. *Annu. Rev. Ecol. Syst.* 23:63–87.
- Drake, D. R., and T. L. Hunt. 2009. Invasive rodents on islands: Incorporating historical and contemporary ecology. *Biol. Invasions* 11:1483–1487.
- Foster, J. T. 2009. The history and impact of introduced birds. Pages 312–330 in T. K. Pratt, C. T. Atkinson, P. C. Banko, J. D. Jacobi, and B. L. Woodworth, eds. *Conservation biology of Hawaiian forest birds.*

- Yale University Press, New Haven, Connecticut.
- Foster, J. T., and S. K. Robinson. 2007. Introduced birds and the fate of Hawaiian rainforests. *Conserv. Biol.* 21:1248–1257.
- Fukumoto, G. K., and C. N. Lee. 2003. Kikuyu for forage. Cooperative Extension Service, College of Tropical Agriculture and Human Resources, University of Hawai'i, Livestock Management, 5.
- Giambelluca, T. W., Q. Chen, A. G. Frazier, J. P. Price, Y.-L. Chen, P.-S. Chu, J. Eischeid, and D. Delparte. 2011. The rainfall atlas of Hawai'i. <http://rainfall.geography.hawaii.edu>.
- Gillespie, T. W., and T. Jaffre. 2003. Tropical dry forests in New Caledonia. *Biodivers. Conserv.* 12:1687–1697.
- Gomez-Aparicio, L. 2009. The role of plant interactions in the restoration of degraded ecosystems: A meta-analysis across life-forms and ecosystems. *J. Ecol.* 97:1202–1214.
- Harris, J. A., R. J. Hobbs, E. Higgs, and J. Aronson. 2006. Ecological restoration and climate change. *Restor. Ecol.* 14:170–176.
- Hobbs, R. J. 2007. Managing plant populations in fragmented landscapes: Restoration or gardening? *Aust. J. Bot.* 55:371–374.
- Hobbs, R. J., E. Higgs, and C. Hall, eds. 2013. *Novel ecosystems: Intervening in the new ecological world order*. John Wiley & Sons, Chichester, United Kingdom.
- Hobbs, R. J., E. Higgs, and J. A. Harris. 2009. *Novel ecosystems: Implications for conservation and restoration*. *Trends Ecol. Evol.* 24:599–605.
- Innes, J., W. G. Lee, B. Burns, C. Campbell-Hunt, C. Watts, H. Phipps, and T. Stephens. 2012. Role of predator-proof fences in restoring New Zealand's biodiversity: A response to Scofield et al. (2011). *N. Z. J. Ecol.* 36:232–238.
- Janzen, D. 1988. Tropical dry forests. The most endangered major tropical ecosystem. Pages 130–137 in E. O. Wilson, ed. *Biodiversity*. National Academy of Sciences/Smithsonian Institution, Washington, D.C.
- Marais, J. P. 2001. Factors affecting the nutritive value of kikuyu grass (*Pennisetum clandestinum*): A review. *Trop. Grassl.* 35:65–84.
- Medeiros, A. C., C. F. Davenport, and C. G. Chimera. 1998. Auwahi: Ethnobotany of a Hawaiian dryland forest. Cooperative National Park Resources Studies Unit, University of Hawai'i at Mānoa, Department of Botany. *Pac. Coop. Stud. Unit Tech. Rep.* 117.
- Medeiros, A. C., T. L. Erwin, C. G. Chimera, and L. L. Loope. 2003. Vegetation trends at Auwahi dryland forest after five years of restoration (Hawaii). *Ecol. Restor.* 21:207–208.
- Medeiros, A. C., L. L. Loope, and R. A. Holt. 1986. Status of native flowering plant species on the south slope of Haleakalā, East Maui, Hawaii. Cooperative National Park Resources Studies Unit, University of Hawai'i at Mānoa, Department of Botany. *Tech. Rep.* 59. Honolulu.
- Medeiros, A. C., and E. I. von Allmen. 2006. Restoration of native Hawaiian dryland forest at Auwahi, Maui. U.S. Geological Survey Fact Sheet FS 2006–3035.
- Meyer, J.-Y., and J.-F. Butaud. 2009. The impacts of rats on the endangered native flora of French Polynesia (Pacific islands): Drivers of plant extinction or coup de grace species? *Biol. Invasions* 11:1569–1585.
- Miles, L., A. C. Newton, R. S. DeFries, C. Ravilious, I. May, S. Blyth, V. Kapos, and J. E. Gordon. 2006. A global overview of the conservation status of tropical dry forests. *J. Biogeogr.* 33:491–505.
- Mueller-Dombois, D., and H. Ellenberg. 2003. *Aims and methods of vegetation ecology*. The Blackburn Press, Caldwell, New Jersey.
- Olson, D. M., and E. Dinerstein. 2002. The global 200: Priority ecoregions for global conservation. *Ann. Mo. Bot. Gard.* 89:199–224.
- Padilla, F. M., and F. I. Pugnaire. 2006. The role of nurse-plants in the restoration of degraded environments. *Front. Ecol. Environ.* 4:196–202.
- Perkins, K. S., J. R. Nimmo, and A. C. Medeiros. 2012. Effects of native forest restoration on soil hydraulic properties, Auwahi,

- Maui, Hawaiian Islands. *Geophys. Res. Lett.* doi:10.1029/2012GL051120.
- Rock, J. F. 1913. The indigenous trees of the Hawaiian Islands. Publ. privately, Honolulu. (Reprinted with addendum in 1974, Charles E. Tuttle Co., Rutland, Vermont).
- Rubinoff, D., B. S. Holland, A. Shibata, R. H. Messing, and M. G. Wright. 2010. Rapid invasion despite lack of genetic variation in the *Erythrina* gall wasp (*Quadrastichus erythrinae* Kim). *Pac. Sci.* 64:23–31.
- Rubinoff, D., and M. San Jose. 2010. Life history and host range of Hawaii's Endangered Blackburn's Sphinx Moth (*Manduca blackburni* Butler). *Proc. Hawaii. Entomol. Soc.* 42:53–59.
- Scholl, M. A., T. W. Giambelluca, S. B. Gingerich, M. A. Nullet, and L. L. Loope. 2007. Cloud water in windward and leeward mountain forests: The stable isotope signature of orographic cloud water. *Water Resour. Res.* 43: W12411. doi:10.1029/2007WR006011.
- Weber, E. 2003. Invasive plant species of the world: A reference guide to environmental weeds. CABI Publishing, Wallingford, United Kingdom.
- Wilmshurst, J. M., T. L. Hunt, C. P. Lipo, and A. J. Anderson. 2011. High-precision radiocarbon dating shows recent and rapid initial human colonization of East Polynesia. *Proc. Natl. Acad. Sci. U.S.A.* 108:1815–1820.
- Zar, J. H. 1999. *Biostatistical analysis*. 4th ed. Prentice Hall, Saddle River, New Jersey.

