

somatic alteration of the remaining allele in a tumour from a patient with a germline *hPMS2* mutation supports the idea that inactivation of both alleles of a mismatch repair gene is required for tumour formation. □

Received 21 June; accepted 22 July 1994.

- Lynch, H. T. et al. *Gastroenterology* **104**, 1535–1549 (1993).
- Peinado, M. A., Malkhosyan, S., Velazquez, A. & Perucho, M. *Proc. natn. Acad. Sci. U.S.A.* **89**, 10065–10069 (1992).
- Ionov, Y., Peinado, A., Malkhosyan, S., Shibata, D. & Perucho, M. *Nature* **363**, 558–561 (1993).
- Thibodeau, S. N., Bren, G. & Schaid, D. *Science* **260**, 816–819 (1993).
- Aaltonen, L. A. et al. *Science* **260**, 812–816 (1993).
- Strand, M., Prolla, T. A., Liskay, R. M. & Petes, T. D. *Nature* **365**, 274–276 (1993).
- Parsons, R. P. et al. *Cell* **75**, 1227–1236 (1993).
- Umar, A. et al. *J. biol. chem.* **269**, 14367–14370 (1994).
- Peltomaki, P. et al. *Science* **260**, 810–812 (1993).
- Lindblom, A., Tannergard, P., Werelius, B. & Nordenskjold, M. *Nature Genet.* **5**, 279–282 (1993).
- Nystrom-Lahti, M. et al. *Proc. natn. Acad. Sci. U.S.A.* **91**, 6054–6058 (1994).
- Fishel, R. et al. *Cell* **75**, 1027–1038 (1993).
- Leach, F. S. et al. *Cell* **75**, 1215–1225 (1993).
- Paiombo, F., Hughes, M. & Jiricny, J. *Nature* **367**, 417–418 (1994).
- Bronner, C. E. et al. *Nature* **368**, 258–261 (1994).
- Papadopoulos, N. et al. *Science* **263**, 1625–1629 (1994).
- Adams, M. D. et al. *Science* **252**, 1651–1656 (1991).
- Frohman, M. A., Dush, M. K. & Martin, G. R. *Proc. natn. Acad. Sci. U.S.A.* **85**, 8998–9002 (1988).
- Kozak, M. *Cell* **44**, 283–292 (1986).
- Prolla, T. A., Christie, D. M. & Liskay, R. M. *Molec. cell. Biol.* **14**, 407–415 (1994).
- Liu, B. et al. *Cancer Res.* (in the press).
- Powell, S. M. et al. *New Engl. J. Med.* **329**, 1982–1987 (1993).
- Van Der Luitj, R. et al. *Genomics* **20**, 1–4 (1994). (Initials?)
- Dietz, H. C., et al. *Science* **259**, 680–683 (1993).
- Prolla, T. A. et al. *Science* (in the press).
- Kramer, W., Kramer, B., Williamson, M. S. & Fogel, S. J. *J. Bact.* **171**, 5339–5346 (1989).
- Schuler, G. D., Altschul, S. F. & Lipman, D. J. *Proteins Struct. Funct. Genet.* **9**, 180–190 (1991).
- Johnson, C. V. et al. *Meth. cell. Biol.* **35**, 73–99 (1991).
- Sidransky, D. et al. *Science* **252**, 706–709 (1991).

ACKNOWLEDGEMENTS. We would like to thank R. Wang, C. Fischer, V. Madike, and the sequencing staff from HGS for technical support. This work was supported by grants from the Clayton Fund, the Academy of Finland, the Finnish Cancer Societies, the Sigrid Juselius Foundation, the Folkhalsan Institute of Genetics, the Cooperative Human Tissue Network (NCI) and the NIH. N.C.N. and N.P. contributed equally to this work.

Crystal structure of an isoleucine-zipper trimer

Pehr B. Harbury*†, Peter S. Kim† & Tom Alber‡

* Department of Biological Chemistry and Molecular Pharmacology, Harvard Medical School, Boston, Massachusetts 02115, USA
 † Howard Hughes Medical Institute, Whitehead Institute, Department of Biology, Massachusetts Institute of Technology, 9 Cambridge Center, Cambridge, Massachusetts 02142, USA
 ‡ Department of Molecular and Cell Biology, University of California, Berkeley, California 94720, USA

SUBUNIT oligomerization in many proteins is mediated by short coiled-coil motifs^{1,2}. These motifs share a characteristic seven-amino-acid repeat containing hydrophobic residues at the first (a) and fourth (d) positions. Despite this common pattern, different sequences form two-, three- and four-stranded helical ropes. We have investigated the basis for oligomer choice by characterizing variants³ of the GCN4 leucine-zipper dimerization domain that adopt trimeric or tetrameric structures in response to mutations at the a and d positions. We now report the high-resolution X-ray crystal structure of an isoleucine-containing mutant that folds into a parallel three-stranded, α -helical coiled coil. In contrast to the dimer and tetramer structures^{3,4}, the interior packing of the trimer can accommodate β -branched residues in the most preferred rotamer at both hydrophobic positions. Compatibility of the shape of the core amino acids with the distinct packing spaces in the two-, three- and four-stranded conformations appears to determine the oligomerization state of the GCN4 leucine-zipper variants.

The GCN4-pII peptide, which differs from the wild-type GCN4 leucine zipper by isoleucine substitutions at four a and four d positions (Fig. 1a; Table 1), forms a parallel trimer in solution³. A similar isoleucine repeat occurs in the trimeric σ 1 protein of reovirus serotype 1 (refs 5, 6) in haemagglutinins from several strains of human parainfluenza virus^{7,8} and in trout axonemal dynein⁹. The trimeric coiled coils from influenza haemagglutinin^{10,11} and the yeast heat-shock factor¹² (HSF) resemble the GCN4-pII sequence at the d positions of the heptad repeat, which contain predominantly β -branched residues (IIIL in the five heptads of HSF).

The X-ray crystal structure of GCN4-pII at 1.8 Å resolution (Fig. 1) shows that three α -helical peptide monomers wrap in a gradual left-handed superhelix. The superhelix describes a cylinder that is ~24 Å wide and ~48 Å long. The isoleucine residues point into the core of the trimer, and cross-sectional layers containing isoleucine at the a positions alternate with layers containing isoleucine at the d positions (Fig. 1d). In seven of eight layers, the dihedral angles χ_1 and χ_2 of the isoleucine side chains are approximately $-60,180$, corresponding to the most abundant rotamer¹³. The isoleucine residues in the C-terminal a layer assume dihedral angles near $+60,180$. Residues at positions e and g pack against the isoleucines at d and a, respectively, to complete the hydrophobic core. Interhelical salt bridges form with high frequency between charged side chains at the g position of one heptad and the e position of a succeeding heptad (for

TABLE 1 Helix-helix interactions*

	Dimer	Trimer	Tetramer	Globular proteins ¹⁷
GCN4 peptide variant†	p1‡ and pL	pI	pL	
Residues at four a positions	V(N)	I	L	
Residues at four d positions	L	I	I	
Superhelix parameter				
Supercoil radius, R_0 (Å)	4.9	6.7	7.6	
Residues per supercoil turn, ω_0	100	118	139	
Supercoil pitch (Å)	148	175	205	
Radius of curvature (Å)	118	124	149	
Superhelix crossing angle (χ)	23.4°	26.8°	26.0°	
Position a orientation angle, ϕ	21.6°	20.4°	19.8°	
α-Helix parameter¹⁷				
Residues per α -helix turn, n	3.62	3.60	3.59	3.64 ± 0.18
Rise/residue, d (Å)	1.51	1.53	1.52	1.51 ± 0.12
α -Helix radius ($C\alpha$), R_1 (Å)	2.28	2.24	2.26	—
Pairwise helix-crossing angle, Ω	23.4°	23.2°	18.3°	19° ± 24°
Pairwise interhelix distance, D (Å)	9.8	11.5	10.6	10.2 ± 2.0
Interhelical salt bridges				
g to e	3 of 6	8 of 9	5 of 12	
g to b and c to e	0 of 8	1 of 12	9 of 16	

* Superhelical characteristics were obtained by fitting the $C\alpha$ backbones to a supercoil parametrization suggested by Crick²² (Fig. 1a). The radius (R_0), frequency (ω_0) and pitch of the superhelix, and the radius (R_1) and phase (ϕ) of the α -helix were treated as variables. The frequency of the α -helix was fixed at 4π radians per 7 amino acids to preserve the periodicity of the heptad structure. α -helix parameters were calculated as defined in ref. 17. The properties of α -helices in globular proteins were taken from that reference and are expressed as the mean value ± 2 standard deviations. The values of D and Ω for globular proteins are averages for the 3–4 ridges-into-grooves packing class¹⁷. Consistent with earlier analyses^{3,22,23}, the superhelical radius and pitch, and the radius of helix curvature increase with the number of strands in the coiled coil.

† The leucine zipper variants were derived from the wild-type GCN4 coiled-coil peptide, GCN4-p1 (ref. 24), by collective mutation to the indicated residues at four a positions (dashed box in Fig. 1a) and four d positions (dashed oval in Fig. 1a)³.

‡ Geometric properties of the dimer conformation were obtained from the structure of GCN4-p1 (ref. 4) because the crystallographic analysis of GCN4-pL has not been completed. The GCN4-p1 sequence contains four leucines at the varied d positions, and three valines and one asparagine at the varied a positions. Interestingly, replacement of the asparagine residue (Asn 16) by valine generates a peptide, GCN4-pVL, that forms a mixture of dimers and trimers³. This result indicates that the buried polar side chain of Asn 16 directs dimer formation.







FIG. 1 Crystal structure of the GCN4-pII trimer. **a**, Helical wheel projection of residues 1 (Arg) to 32 (Glu) of the GCN4-pII sequence. View is from the N terminus, and residues in the first two helical turns are boxed or circled. Heptad positions are labelled **a-g**. The R_0 , ω_0 , α , ϕ and R_1 variables from a coiled-coil parameterization suggested by Crick^{3,22} and the D and Ω variables defined in ref. 17 are illustrated. **b**, A portion of the 6.0–1.8 Å resolution $2F_o - F_c$ electron-density map superimposed on the final model. A side view of residues Ser C14 to Asn C21 is shown. **c**, Axial view of the GCN4-pII trimer. The van der Waals surfaces are coloured purple for residues at **a** positions and green for residues at **d** positions. **d**, Side view of the trimer. The van der Waals surfaces of the N-terminal methionine layer are yellow.


METHODS. The GCN4-pII peptide, Ac-R MKQIEDK IEEILSK IYHIENE IAR IKKIGER (where mutant **a** residues are underlined and mutant **d** residues double-underlined; Ac, acetyl), was crystallized by vapour diffusion from 100 mM sodium citrate, pH 4.8, 600 mM NaBr and 5% PEG 400. The crystals had $P2_1$ symmetry ($a = 35.32$ Å, $b = 39.67$ Å, $c = 36.97$ Å, $\beta = 98.38^\circ$) with a trimer in the asymmetric unit. Data to 1.8 Å resolution were collected using an R -Axis image plate detector (7,052 unique reflections, $R_{\text{merge}} = 0.045$). An idealized model was built²⁵ using an empirical method for calculation of coiled-coil structures based on an algebraic parameterization²². The model was used in a combined rotation–translation search (with 6–2.9 Å resolution data $> 2\sigma$) along a non-crystallographic 3-fold symmetry axis (located with GLRF²⁶) using the program XPLOR²⁷. The coordinates corresponding to the peak solution (initial $R = 52\%$) were annealed and rebuilt into simulated-annealing omit maps²⁸. Water molecules were added and the coordinates and individual B values were refined using TNT²⁹ and FRODO³⁰. The final model, which includes 96 amino acids of the trimer and 48 water molecules, has a crystallographic R -factor of 0.175 (6–1.8 Å data) with root-mean-square deviations from ideal bond lengths and bond angles of 0.013 Å and 2.4° respectively. All main-chain dihedral angles fall within allowed regions of the Ramachandran plot, and most side chains assume well populated rotamer conformations. The coordinates of the search model and the refined structure differ by a root-mean-square deviation of 0.46 Å for main-chain and core side-chain atoms. These results demonstrate the feasibility of using a predicted structure to initiate crystallographic analysis. The coordinates have been deposited in the Brookhaven Protein Data Bank.



example, Glu B22–Lys C27; Table 1 and Fig. 1a). This observation supports proposals that heterotrimer formation could be directed by electrostatic attraction¹⁴ or repulsion^{15,16} between these residues.

Although the pitch, radius and residues per turn of the superhelices in the trimer (GCN4-pII), dimer (GCN4-pI) and

tetramer (GCN4-pLI) differ greatly, the individual α -helices are virtually identical on a local scale (Table 1). Moreover, pairs of adjacent helices in the three structures have similar separations (D) and crossing angles (Ω). This conserved pairwise arrangement of helices corresponds to the optimal geometry for one class of helix association in globular proteins¹⁷ (Table 1).



The GCN4-pII trimer shows 'knobs-into-holes' packing, in which the side chains (knobs) at positions **a** and **d** fit into the spaces (holes) between four residues on a neighbouring helix¹⁸. In both the **a** and **d** layers, the $Ca-C\beta$ bond of the knob side chain makes an acute angle with the $Ca-C\alpha$ vector defining the base of the apposed hole (Fig. 2*b, e*). This shared geometry results in a similar placement of atoms around the side chains at positions **a** and **d** (Fig. 3*b*). However, the acute angles in the two types of layers differ by $\sim 25^\circ$ (Fig. 3*b*). In addition, the third helix of the trimer sits to the C-terminal side of the hole in the **a** layer (anti-clockwise from hole in Fig. 2*b*) but to the N-terminal side of the hole in the **d** layer (clockwise from hole in Fig. 2*e*). Thus, for example, a threonine knob residue with a $(-)\chi_1$ dihedral angle would direct its hydroxyl group into the hydrophobic interface at an **a** position, but towards solvent at a **d** position.

The interior packing of the trimer differs from that seen in the dimeric and tetrameric leucine-zipper variants. In the GCN4-pI dimer, the $Ca-C\beta$ bond of the knob side chain at position **a** points out of the interface (parallel packing; Fig. 2*a*), and the $Ca-C\beta$ bond of the knob side chain at position **d** points directly at the adjacent helix (perpendicular packing; Fig. 2*d*). This pattern is reversed in the GCN4-pLI tetramer (Figs 2*c*, and 3*a, c*), which shows perpendicular packing in the **a** layers and parallel packing in the **d** layers. Thus the packing geometries in the dimer, trimer and tetramer are distinct from one another at both hydrophobic positions (Fig. 2).


FIG. 3 Superposition³¹ of parallel, perpendicular and acute knobs-into-holes packing. *a*, Parallel packing in the dimer **a** layer (green) and the tetramer **d** layer (blue). *b*, Acute packing in the trimer **a** layer (green) and the trimer **d** layer (blue). *c*, Perpendicular packing in the tetramer **a** layer (green) and the dimer **d** layer (blue).

FIG. 2 Packing differences in parallel coiled coils^{3,18}. Helix cross-sectional layers centred on the **a** positions (containing the residues at positions **f, g, a** and **b**) and on the **d** positions (containing the residues at positions **c, d, e** and **f**) are depicted³¹. Knobs formed by the side chains of one helix fit into holes formed by the spaces between side chains on a neighbouring helix. Looking from the N terminus down the superhelix axis, each knob at an **a** position (red) packs into a hole formed between the **g** and **a** residues (yellow) and two **d** residues in adjacent layers (not shown) of the anti-clockwise-related monomer. Similarly, each knob at a **d** position (red) packs into a hole formed between the **d** and **e** residues (yellow) and two **a** residues in adjacent layers (not shown) of the clockwise related monomer. *a*, Position **a** of the dimer: the $Ca-C\beta$ bond of each valine knob (red) is oriented parallel to the $Ca-C\alpha$ vector at the base of the hole into which it projects (yellow). *b*, Position **a** of the trimer; the $Ca-C\beta$ bond of each isoleucine knob (red) forms an acute angle ($\sim 50^\circ$) with the $Ca-C\alpha$ vector at the base of the recipient hole (yellow). *c*, Position **a** of the tetramer; the $Ca-C\beta$ bond of each leucine knob (red) is perpendicular to the $Ca-C\alpha$ vector at the base of the apposed hole (yellow). *d*, Perpendicular packing at position **d** of the dimer. *e*, Acute packing at position **d** of the trimer. *f*, Parallel packing at position **d** of the tetramer.

The three modes of knobs-into-holes packing exhibit different preferences for specific amino acids at the knob position. In dimeric coiled-coil sequences, the **a** positions (parallel packing) are enriched for the β -branched residues valine and isoleucine, whereas the **d** positions (perpendicular packing) are sharply depleted of β -branched residues and enriched for leucine^{19,20}. In trimeric coiled-coil sequences, however, leucine and the β -branched amino acids are approximately equally represented and distributed evenly between the **a** and **d** positions²¹ (consistent with the geometric similarity of the packing at **a** and **d** in the trimer; Fig. 3*b*). Thus parallel geometry favours β -branched residues, perpendicular geometry disfavors β -branched residues and favours leucine, and acute geometry shows little preference.

The oligomerization states of the GCN4 leucine-zipper mutants are apparently dictated by matching the core residues, determined by sequence, with appropriate packing geometry, determined by structure. The dimer of GCN4-pIL and the tetramer of GCN4-pLI, for example, place the leucine residues in perpendicular geometry and the isoleucine residues in parallel geometry. By contrast, a dimer or tetramer of GCN4-pII would mismatch four of the eight buried isoleucines with perpendicular geometry. The trimer conformation, however, places all the isoleucines in acute geometry, thereby accommodating β -branched amino acids at both the **a** and **d** positions.

The packing properties of the β -branched residues isoleucine and valine are not identical. For example, the peptide



GCN4-pVL forms both dimers and trimers and the peptide GCN4-pLV is trimeric³. Comparison of the oligomerization properties of GCN4-pVL and GCN4-pIL (dimer), or GCN4-pLV and GCN4-pLI (tetramer), suggests that isoleucine exhibits a stronger preference than valine for parallel over acute geometry. Thus, the GCN4 variant coiled coils provide a striking example of how side-chain shape and packing can dramatically affect the global fold of a protein. □

Received 27 April; accepted 8 July 1994.

- Cohen, C. & Parry, D. A. D. *Proteins* **7**, 1–15 (1990).
- Cohen, C. & Parry, D. A. D. *Science* **263**, 488–489 (1994).
- Harbury, P. B., Zhang, T., Kim, P. S. & Alber, T. *Science* **262**, 1401–1407 (1993).
- O'Shea, E. K., Klemm, J. D., Kim, P. S. & Alber, T. *Science* **254**, 539–544 (1991).
- Duncan, R., Horne, D., Cashdollar, L. W., Joklik, W. K. & Lee, P. W. K. *Virology* **174**, 399–409 (1990).
- Strong, J. E., Leone, G., Duncan, R., Sharma, R. K. & Lee, P. W. K. *Virology* **184**, 23–32 (1991).
- Elango, N., Coligan, J. E., Jambou, R. C. & Venkatesan, S. *J. Virol.* **57**, 481–489 (1986).
- Coelingh, K. L. V. W., Winter, C. C. & Murphy, B. R. *Virology* **162**, 137–143 (1988).
- Garber, A. T., Retief, J. D. & Dixon, G. H. *EMBO J.* **6**, 1727–1734 (1989).
- Wilson, I. A., Skehel, J. J. & Wiley, D. C. *Nature* **289**, 366–373 (1981).
- Carr, C. M. & Kim, P. S. *Cell* **73**, 823–832 (1993).
- Sorger, P. K. & Nelson, H. C. *Cell* **59**, 807–813 (1989).
- Ponder, J. W. & Richards, F. M. *J. molec. Biol.* **193**, 775–791 (1987).
- Beck, K., Dixon, T. W., Engel, J. & Parry, D. A. *J. molec. Biol.* **231**, 311–323 (1993).
- Coelingh, K. L. V. W., Winter, C. C. & Murphy, B. R. *Virology* **162**, 137–143 (1988).
- O'Shea, E. K., Lumb, K. J. & Kim, P. S. *Curr. Biol.* **3**, 658–667 (1993).
- Chothia, C., Levitt, M. & Richardson, D. J. *molec. Biol.* **145**, 215–250 (1981).
- Crick, F. H. C. *Acta crystallogr.* **6**, 689–697 (1953).
- Hu, J. C. & Sauer, R. T. in *Progress in Nucleic Acids and Molecular Biology* (eds Eckstein, F. & Lilley, D. M. J.) 82–101 (Springer, Berlin, 1992).
- Conway, J. F. & Parry, D. A. D. *Int. J. biol. Macromol.* **12**, 328–334 (1990).
- Conway, J. F. & Parry, D. A. D. *Int. J. biol. Macromol.* **13**, 14–16 (1991).
- Crick, F. H. C. *Acta crystallogr.* **6**, 685–689 (1953).
- Seo, J. & Cohen, C. *Proteins* **5**, 223–234 (1993).
- O'Shea, E. K., Rutkowski, R. & Kim, P. S. *Science* **243**, 538–542 (1989).
- Harbury, P. H. thesis, Harvard Univ. (1994).
- Tong, L. & Rossmann, M. G. *Acta crystallogr.* **A46**, 783–792 (1990).
- Brunger, A. T., Kuriyan, J. & Karplus, M. *Science* **235**, 458–460 (1987).
- Brunger, A. T. *J. molec. Biol.* **203**, 803–816 (1988).
- Tronrud, D. E., TenEyck, L. F. & Matthews, B. W. *Acta crystallogr.* **A43**, 489–501 (1987).
- Jones, T. A. *J. appl. Crystallogr.* **11**, 268–272 (1978).
- Kraulis, P. J. *J. appl. Cryst.* **24**, 946–950 (1991).

ACKNOWLEDGEMENTS. We thank M. Milhollen for peptide synthesis. This work was supported by grants from the National Institutes of Health (to P.S.K. and T.A.) and the Lucille P. Markey Charitable Trust (to T.A.).

CORRECTION

Flattening of the sea-floor depth–age curve as a response to asthenospheric flow

Jason Phipps Morgan & Walter H. F. Smith

Nature **359**, 524–527 (1992)

WE are grateful to Carol and Seth Stein¹ for alerting us to errors in our discussion of the results shown in Fig. 2b of our Letter. In the calculations shown in this Figure, the asthenospheric viscosity beneath the South Atlantic lithosphere was 4.2×10^{19} Pa s and the African plate was assumed to be stationary, instead of migrating eastward at 10 mm yr^{-1} as stated in the Letter. In addition, the thermal subsidence rate and ridge-crest depth, which were not mentioned, were $280 \text{ m Myr}^{-1/2}$ and $2,200 \text{ m}$ respectively. In contrast, the values used for the Pacific plate were, as stated, $300 \text{ m Myr}^{-1/2}$ and $2,644 \text{ m}$. Hence the fit to the Pacific and South Atlantic plates results from assuming subsidence rates differing by 7% and viscosities differing by a factor of 21. □

1. Stein, C. & Stein, S. *Geophys. Res. Lett.* **21**, 709–712 (1994).

PLEASE follow these guidelines so that your manuscript may be handled expeditiously.

Nature is an international journal covering all the sciences. Contributors should therefore bear in mind those readers who work in other fields and those for whom English is a second language, and write clearly and simply, avoiding unnecessary technical terminology. Space in the journal is limited, making competition for publication severe. Brevity is highly valued. One printed page of *Nature*, without display items, contains about 1,300 words.

Manuscripts are selected for publication according to editorial assessment of their suitability and reports from independent referees. They can be sent to London or Washington and should be addressed to the Editor. Manuscripts may be dealt with in either office, depending on the subject matter, and will where necessary be sent between offices by overnight courier. High priority cannot be given to pre-submission enquiries; in urgent cases they can be made in the form of a one-page fax. All manuscripts are acknowledged on receipt but fewer than half are sent for review. Those that are not reviewed are returned as rapidly as possible so that they may be submitted elsewhere without delay. Contributors may suggest reviewers; limited requests for the exclusion of specific reviewers are usually heeded. Manuscripts are usually sent to two or three reviewers, chosen for their expertise rather than their geographical location. Manuscripts accepted for publication are processed from the London office.

Nature requests authors to deposit sequence and crystallographic data in the databases that exist for this purpose.

Once a manuscript is accepted for publication, contributors will receive proofs in about 4 weeks. *Nature's* staff will edit manuscripts with a view to brevity and clarity, so contributors should check proofs carefully. Manuscripts are generally published 2–3 weeks after receipt of corrected proofs. *Nature* does not exact page charges. Contributors receive a reprint order form with their proofs; reprint orders are processed after the manuscript is published and payment received.

Categories of paper

Review Articles survey recent developments in a field. Most are commissioned but suggestions are welcome in the form of a one-page synopsis addressed to the Reviews Coordinator. Length is negotiable in advance.

Progress articles review particularly topical developments for a nonspecialist readership. They do not exceed 4 pages in length. Suggestions may be made to the Reviews Coordinator in the form of a brief synopsis.

Articles are research reports whose conclusions are of general interest and which are sufficiently rounded to be a substantial advance in understanding. They should not have more than 3,000 words of text (not including figure legends) or more than six display items and should not occupy more than five pages of *Nature*.

Articles start with a heading of 50–80 words written to advertise their content in general terms, to which editors will pay particular attention. The heading does not usually contain numbers, abbreviations or measurements. The introduction to the study is contained in the first two or three paragraphs of the article, which also briefly summarize its results and implications. Articles have fewer than 50 references and may contain a few short subheadings.

Letters to Nature are short reports of outstanding novel findings whose implications are general and important enough to be of interest to those outside the field. Letters should have 1,000 or fewer words of text and four or fewer display items. The first paragraph describes, in not more than 150 words and without the use of abbreviations, the background, rationale and chief conclusions of the study for the particular benefit of non-specialist readers. Letters do not have subheadings and contain fewer than 30 references.

Commentary articles deal with issues in, or arising from, research that are also of interest to readers outside research.

News and Views articles inform nonspecialist readers about new scientific advances, sometimes in the form of a conference report. Most are commissioned but proposals can be made in advance to the News and Views Editor.

Scientific Correspondence is for discussion of topical scientific matters, including those published in *Nature*, and for miscellaneous contributions. Priority is given to letters of fewer than 500 words.

Preparation of manuscripts

All manuscripts should be typed, double-spaced, on one side of the paper only. An original and four copies are required, each accompanied by artwork. If photographs are included, five sets of originals are required, for line drawings, one set of originals and four good-quality photocopies are acceptable. Reference lists, figure legends and tables should all be on separate sheets, all of which should be double-spaced and numbered. Three copies of relevant manuscripts in press or submitted for publication elsewhere should be included with submitted manuscripts, clearly marked as such. Five copies of revised and resubmitted manuscripts, labelled with their manuscript numbers, are required, together with five copies of a letter detailing the changes made.

Titles are brief and simple. Active verbs, numerical values, abbreviations and punctuation are to be avoided. Titles should contain one or two key words for indexing purposes.

Artwork should be marked individually and clearly with the author's name and, when known, the manuscript number. Ideally, no figure should be larger than 28 by 22 cm. Figures with several parts are to be avoided and are permitted only if the parts are closely related, either experimentally or logically. Unlettered originals of photographs should be provided. Suggestions for cover illustrations, with captions and labelled with the manuscript number, are welcome. Original artwork is returned when a manuscript cannot be published.

Protein/nucleotide sequences should ideally be in the three-letter and not the single-letter code for amino acids. One column width of *Nature* can accommodate 20 amino acids or 60 base pairs. **Colour artwork.** A charge of £500 per page is made as a contribution towards the cost of reproducing colour figures. Inability to pay these costs will not prevent publication of essential colour figures if the circumstances are explained. Proofs of colour artwork may be sent to contributors under separate cover from their galley proofs.

Figure legends should not exceed 300 words and ideally should be shorter. The figure is described first, then, briefly, the method. Reference to a method published elsewhere is preferable to a full description. Methods are not described in the text.

References are numbered sequentially as they appear in the text, followed by those in tables and finally by those in figure legends. Only papers published or in the press are numbered and included in the reference list. All other forms of reference should be cited in the text as a personal communication, manuscript submitted or in preparation. Text is not included in reference lists. References are abbreviated according to the *World List of Scientific Periodicals* (Butterworths, London, 1963–65). The first and last page numbers are included; reference to books includes publisher, place and date.

Abbreviations, symbols, units and Greek letters should be identified the first time they are used. Acronyms should be avoided whenever possible and, if used, defined. Footnotes are not used in the text.

Acknowledgements are brief and appear after the reference list; grant and contribution numbers are not allowed.

Supplementary Information is material relevant to Articles or Letters which cannot, for lack of space, be published in full, but which is available from *Nature* on request.

Submission. Manuscripts can be sent to the Editor at 4 Little Essex Street, London WC2R 3LF, UK or at 1234 National Press Building, Washington, DC 20045, USA. A telephone and fax number should be included. Manuscripts or proofs sent by air courier to London should be declared as manuscripts and 'value \$5' to prevent the imposition of import duty and value-added tax.