

Preventive Care Benefits for Children

All Marketplace health plans and many other plans must cover the following list of preventive services without charging you a copayment or coinsurance. This is true even if you haven't met your yearly deductible.

IMPORTANT

These services are free only when delivered by a doctor or other provider in your plan's network.

Coverage for children's preventive health services:

1. Alcohol and drug use assessments for adolescents
2. Autism screening for children at 18 and 24 months
3. Behavioral assessments for children ages: 0 to 11 months, 1 to 4 years, 5 to 10 years, 11 to 14 years, 15 to 17 years
4. Blood pressure screening for children ages: 0 to 11 months, 1 to 4 years, 5 to 10 years, 11 to 14 years, 15 to 17 years
5. Cervical dysplasia screening for sexually active females
6. Depression screening for adolescents
7. Developmental screening for children under age 3
8. Dyslipidemia screening for children at higher risk of lipid disorders ages: 1 to 4 years, 5 to 10 years, 11 to 14 years, 15 to 17 years
9. Fluoride chemoprevention supplements for children without fluoride in their water source
10. Gonorrhea preventive medication for the eyes of all newborns
11. Hearing screening for all newborns
12. Height, weight and body mass index (BMI) measurements for children ages: 0 to 11 months, 1 to 4 years, 5 to 10 years, 11 to 14 years, 15 to 17 years
13. Hematocrit or hemoglobin screening for all children
14. Hemoglobinopathies or sickle cell screening for newborns
15. Hepatitis B screening This link takes you to a website not operated by the federal government. The site may have different privacy and security policies. for adolescents at high risk, including adolescents from countries with 2% or more Hepatitis B prevalence, and U.S.-born adolescents not vaccinated as infants and with at least one parent born in a region with 8% or more Hepatitis B prevalence: 11 - 17 years.
16. HIV screening for adolescents at higher risk
17. Hypothyroidism screening for newborns
18. Immunization vaccines for children from birth to age 18 — doses, recommended ages, and recommended populations vary:

| | |
|---|-------------------------------|
| Diphtheria, Tetanus, Pertussis (Whooping Cough) | Haemophilus influenzae type b |
| Hepatitis A | Hepatitis B |
| Human Papillomavirus (PVU) | Inactivated Poliovirus |
| Influenza (flu shot) | Measles |
| Meningococcal | Pneumococcal |
| Rotavirus | Varicella (Chickenpox) |
19. Iron supplements for children ages 6 to 12 months at risk for anemia
20. Lead screening for children at risk of exposure
21. Medical history for all children throughout development ages: 0 to 11 months, 1 to 4 years, 5 to 10 years, 11 to 14 years, 15 to 17 years
22. Obesity screening and counseling
23. Oral health risk assessment for young children ages: 0 to 11 months, 1 to 4 years, 5 to 10 years
24. Phenylketonuria (PKU) screening for newborns
25. Sexually transmitted infection (STI) prevention counseling and screening for adolescents at higher risk
26. Tuberculin testing for children at higher risk of tuberculosis ages: 0 to 11 months, 1 to 4 years, 5 to 10 years, 11 to 14 years, 15 to 17 years
27. Vision screening for all children