

Champion positive health outcomes and behaviors through innovative programs and community engagement

Respiratory Disease Surveillance Report Jefferson County, Missouri

Week 10: March 6, 2022 – March 12, 2022

OVERVIEW: The Jefferson County Health Department (JCHD) monitors influenza and other respiratory pathogens, including COVID-19, throughout Jefferson County, Missouri. The Jefferson County Health Department conducts influenza surveillance using passive surveillance and syndromic surveillance. Although influenza illness can occur year-round, the seasonal influenza reporting begins on the Morbidity and Mortality Weekly Report (MMWR¹) Week 40 of a given year and continues through MMWR Week 20 of the following calendar year. COVID-19 surveillance began in March 2020 and has been ongoing. COVID-19 data reflects cases that are entered into Missouri's communicable disease database. Vaccination data for COVID-19 is reported to the Jefferson County Health Department weekly by the Missouri Department of Health and Senior Services (MO DHSS). *All data and information are conditional and may change as more reports are received*.

-

consist of 52 weeks.

¹ Data is reported in epidemiologic weeks established by the CDC's Morbidity and Mortality Weekly Report (MMWR). The MMWR week starts on a Sunday and ends on Saturday. Values for MMWR week range from 1 to 53, although most years

INFLUENZA SURVEILLANCE

Passive Surveillance

Table 1. Number of Laboratory-positive² Influenza Cases by Week, Jefferson County, Missouri

CDC Reporting					
Week	Week Start Date	Α	В	Unknown	Total
40	10/3/2021	5	2	0	7
41	10/10/2021	0	0	0	0
42	10/17/2021	3	0	0	3
43	10/24/2021	2	1	0	3
44	10/31/2021	7	2	0	9
45	11/7/2021	2	4	0	6
46	11/14/2021	3	1	0	4
47	11/21/2021	3	1	0	4
48	11/28/2021	33	7	0	40
49	12/5/2021	73	6	0	79
50	12/12/2021	217	19	0	236
51	12/19/2021	215	6	0	221
52	12/26/2021	167	6	0	173
1	1/2/2022	66	7	0	73
2	1/9/2022	41	4	0	45
3	1/16/2022	46	2	0	48
4	1/23/2022	25	2	0	27
5	1/30/2022	11	2	0	13
6	2/6/2022	10	2	0	12
7	2/13/2022	21	5	0	26
8	2/20/2022	19	4	0	23
9	2/27/2022	26	3	0	29
10	3/6/2022	15	1	0	16

² Laboratory-positive influenza includes the following test methods: rapid influenza diagnostic tests (antigen), reverse transcriptase polymerase chain reaction (RTPCR) and other molecular assays, immunofluorescence antibody staining (Direct (DFA) or Indirect (IFA)), or viral culture.

Table 2. Number of Laboratory-positive Influenza Cases by Age Group, Jefferson County, Missouri

For Cases Reported 10/3/2021- 3/12/2022

Age Group	Type A	Type B	Unknown Type	Total	% TOTAL
00 to <02	14	2	0	16	1.5%
02 to 04	70	5	0	75	6.8%
05 to 14	357	18	0	375	34.2%
15 to 24	228	15	0	243	22.2%
25 to 49	197	29	0	226	20.6%
50 to 64	75	11	0	86	7.8%
65+	69	7	0	76	6.9%
TOTAL	1010	87	0	1097	100.0%

Figure 1. Percentage of Influenza Cases by Type, Jefferson County, Missouri

For Cases Reported 10/3/2021- 3/12/2022





Figure 2. Number of Laboratory-positive Influenza Cases by Week, Influenza Seasons 2018-19 to 2021-22, Jefferson County, Missouri



^{*}There were 53 weeks in 2020. The estimates for the other three years' values for the weeks 53 are the averages of weeks 52 and weeks 1.

Syndromic Surveillance

As determined by ESSENCE, Influenza-Like-Illness (ILI³) accounted for 0% to 0.47% of daily visits to hospital emergency departments (EDs) for Jefferson County residents during Week 10. The number of visits to local hospital EDs for ILI ranged from zero to one per day during Week 10.

ILI as a Percentage of Daily ED Visits, Jefferson County, MO 6.0 5.5 5.0 4.5 4.0 Percent 3.5 3.0 2.5 2.0 1.5 1.0 0.5 0.0 Data: Warning Data: Alert Data: Normal

Figure 3. Syndromic Surveillance for Influenza-Like-Illness (ILI) Jefferson County, Missouri

Jefferson County Influenza Outbreaks

No influenza outbreaks have been reported in Jefferson County as of Week 10.

Jefferson County Influenza Deaths⁴

Two influenza-associated deaths have been reported in Jefferson County as of Week 10.

³ ILI is defined by ESSENCE as Emergency Department chief complaints for Influenza or (FeverPlus and (Cough or SoreThroat) and not NonILIFevers).

⁴ All influenza-associated deaths became reportable in Missouri in 2016.

COVID-19 SURVEILLANCE

From 3/5/22 to 3/12/22, the average number of new COVID-19 cases diagnosed among Jefferson County residents decreased by 45.00 percent from 14.29 to 7.86 cases per day. The current rate of weekly COVID-19 diagnoses (24.44 cases per 100,000 residents per week) is moderate.

Jefferson County, MO Covid-19 Case Count and 7-Day Rolling **Average** 800 700 600 Number of Cases 500 400 300 200 100 9/2020 11/2020 3/2021 5/2021 7/2021 9/2021 3/2020 5/2020 7/2020 11/2021 1/2022 Lab Collection Date Daily New Cases 7-day Avg

Figure 4. Confirmed and Probable Covid-19 Cases Over Time and 7-Day Average, Jefferson County, Missouri


During Week 10 a total of eleven Covid-19 associated deaths were reported in Jefferson County. A cumulative total of 511 Covid-19 associated deaths have been reported in Jefferson County as of Week 10.

Table 3. Covid-19 Fatalities and Fatality Rates by Age Group,
Jefferson County, Missouri

Age Range:	Fatalities	Rate Per 100
0-9	0	0.0
10-19	0	0.0
20-29	2	0.0
30-39	8	0.2
40-49	20	0.4
50-59	54	1.1
60-69	108	2.7
70-79	141	6.7
80+	178	15.5

As of March 11, 2021, the percentage of Jefferson County residents who have initiated vaccination for COVID-19 is 54.45%. The percentage of residents who have completed the full series of vaccination is 50.41%. The percentage of residents who have received a third dose or booster is 20.00%.

Figure 5. Covid-19 Doses Administered to Jefferson County Residents Over Time and 7-Day Average, Jefferson County, Missouri



ADDITIONAL DATA SOURCES AND INFORMATION

Jefferson County Health Department's Covid-19 Data Dashboard https://www.jeffcohealth.org/covid19-data

Centers for Disease Control and Prevention, National Influenza Surveillance: https://www.cdc.gov/flu/weekly/

Missouri Department of Health and Senior Services, Statewide Influenza Surveillance: https://health.mo.gov/living/healthcondiseases/communicable/influenza/reports.php

World Health Organization, International Influenza Surveillance: http://www.who.int/influenza/surveillance_monitoring/en/

The National Respiratory and Enteric Virus Surveillance System (NREVSS): https://www.cdc.gov/surveillance/nrevss/

Passive Surveillance

Influenza is a reportable condition in the state of Missouri. JCHD conducts passive influenza surveillance by collecting and aggregating data on all laboratory-positive influenza tests in Jefferson County residents. Passive surveillance provides information on the true burden of influenza illness in Jefferson County but is limited by variations in testing and reporting practices. If diagnostic tests are not conducted on patients presenting to health care providers with influenza-like-illness, or if test results (e.g., of rapid influenza tests) are not reported to JCHD, those persons will not be included in the passive surveillance data. Thus, passive surveillance is likely an underestimate of the true burden of influenza illness in Jefferson County. Passive surveillance helps the health department to keep track of the incidence of influenza cases in the community and facilitates early detection of potential outbreaks.

Syndromic Surveillance

JCHD uses the ESSENCE system to conduct syndromic surveillance for Influenza-Like-Illness (ILI) at Missouri hospital emergency departments (EDs). ESSENCE captures data on all ED visits in persons with chief complaints (rather than final diagnoses or positive laboratory tests) of ILI. ILI chief complaints are those which include the word "influenza" or those that the ESSENCE system parses to [fever and (cough or sore throat)]. The syndromic surveillance data presented above include all ED visits for ILI from those with a Jefferson County residential address. Data from private physicians, clinics, or urgent care centers are not included in ESSENCE. Syndromic surveillance contributes to our understanding of the burden of influenza experienced by the health care system, as well as demonstrating the relative impact over time of influenza with respect to other illnesses.