

**Delegates:**  
**McDonald, Elizabeth**

**Editor:**  
**Parvizi, Javad**

## **F- 21 - FAILURE PREDICTORS**

### **F- 21 - What are the predictors of treatment failure in patients who have undergone two-stage exchange for infected total ankle arthroplasty?**

#### **Response/Recommendation:**

Predictors for treatment failure in patients undergoing two-stage exchange for infected TAA include compromised soft tissues (e.g. sinus tract, exposed hardware, etc.); significant bone involvement/osteomyelitis; and insufficient timing of antibiotic course before reimplantation.

**Strength of Recommendation:** Limited

#### **Rationale:**

The optimal management of patients with infected TAA is not well known due to limited number of studies.<sup>1-5</sup> While comparisons and deductions can be made from the knee and hip PJI literature on two-stage exchange, management of infected TAA can differ from hip and knee arthroplasty because of the precarious soft tissue coverage around ankle, oftentimes history of multiple preceding operations, and arthroplasty design updates coupled with limited surgical experience.<sup>3</sup> Two-stage exchange is a well-accepted surgical management approach for PJI.

There is limited detail in the TAA literature on two-stage exchange failure. A study by Patton et al reported on 12 cases of two-stage revision for infected TAA but offered no details of the cases that failed.<sup>3</sup> another study by Kessler et al reported on 34 patients undergoing surgical management for infected TAA. Of the patients treated for infected TAA, 10% (1/10) of two-stage exchanges resulted in failure.<sup>6</sup> This two-stage failure is not described in detail.<sup>6</sup> However, in the described cohort, the presence of compromised soft-tissue significantly increased the rate of failure after revision.<sup>6</sup>

Another problem with the soft tissues is the presence of a sinus tract around the ankle. Not only do sinus tracts often have indurated soft tissue around the ankle, but also a sinus tract has the potential to limit pre-operative cultures and organism identification, which in itself may predispose patient to a future failure.<sup>7-11</sup> Further, certain comorbidities such as metastatic disease, renal and/or liver dysfunction, and advanced cardiac disease are indicated to influence rate of PJI,<sup>6,12</sup> but these comorbidities may not necessarily be tied to treatment failure after two-stage exchange arthroplasty.

In North America patients undergoing two-stage exchange arthroplasty for treatment of PJI, are often subjected to six weeks of antibiotic course. Based on data from the hip and knee PJI, inadequate administration of antibiotics has been linked to the presence of positive cultures during reimplantation that in turn increases the risk of failure after reimplantation.<sup>7,13</sup> While inadequate antibiotic therapy has been linked with subsequent failure, the exact duration of

antibiotic treatment, the benefit of IV-to-PO antibiotics, and the timing of IV-to-PO switch have not been determined. Recent PJI literature suggests that a short IV antibiotic period lasting at least 5-7 days followed by pathogen-specific PO therapy may be a viable option for treatment of patients with PJI.<sup>14,15</sup>

## References:

1. Reuver JM, Dayerizadeh N, Burger B, Elmans L, Hoelen M, Tulp N. Total ankle replacement outcome in low volume centers: short-term followup. *Foot Ankle Int.* 2010;31(12):1064-1068. doi:10.3113/FAI.2010.1064
2. Gougoulas N, Khanna A, Maffulli N. How successful are current ankle replacements?: a systematic review of the literature. *Clin Orthop.* 2010;468(1):199-208. doi:10.1007/s11999-009-0987-3
3. Patton D, Kiewiet N, Brage M. Infected total ankle arthroplasty: risk factors and treatment options. *Foot Ankle Int.* 2015;36(6):626-634. doi:10.1177/1071100714568869
4. Lee K-B, Cho S-G, Hur C-I, Yoon T-R. Perioperative complications of HINTEGRA total ankle replacement: our initial 50 cases. *Foot Ankle Int.* 2008;29(10):978-984. doi:10.3113/FAI.2008.0978
5. Young JL, May MM, Haddad SL. Infected total ankle arthroplasty following routine dental procedure. *Foot Ankle Int.* 2009;30(3):252-257. doi:10.3113/FAI.2009.0252
6. Kessler B, Sendi P, Graber P, et al. Risk factors for periprosthetic ankle joint infection: a case-control study. *J Bone Joint Surg Am.* 2012;94(20):1871-1876. doi:10.2106/JBJS.K.00593
7. Osmon DR, Berbari EF, Berendt AR, et al. Executive summary: diagnosis and management of prosthetic joint infection: clinical practice guidelines by the Infectious Diseases Society of America. *Clin Infect Dis Off Publ Infect Dis Soc Am.* 2013;56(1):1-10. doi:10.1093/cid/cis966
8. Jackson WO, Schmalzried TP. Limited role of direct exchange arthroplasty in the treatment of infected total hip replacements. *Clin Orthop.* 2000;(381):101-105.
9. Buechel FF, Femino FP, D'Alessio J. Primary exchange revision arthroplasty for infected total knee replacement: a long-term study. *Am J Orthop Belle Mead NJ.* 2004;33(4):190-198; discussion 198.
10. Winkler H, Stoiber A, Kaudela K, Winter F, Menschik F. One stage uncemented revision of infected total hip replacement using cancellous allograft bone impregnated with antibiotics. *J Bone Joint Surg Br.* 2008;90(12):1580-1584. doi:10.1302/0301-620X.90B12.20742
11. Del Pozo JL, Patel R. Clinical practice. Infection associated with prosthetic joints. *N Engl J Med.* 2009;361(8):787-794. doi:10.1056/NEJMcp0905029
12. Kessler B, Knupp M, Graber P, et al. The treatment and outcome of peri-prosthetic infection of the ankle: a single cohort-centre experience of 34 cases. *Bone Jt J.* 2014;96-B(6):772-777. doi:10.1302/0301-620X.96B6.33298
13. Zimmerli W, Trampuz A, Ochsner PE. Prosthetic-Joint Infections. *N Engl J Med.* 2004;351(16):1645-1654. doi:10.1056/NEJMra040181
14. Scarborough M, Li HK, Rombach I, et al. Oral versus intravenous antibiotics for the treatment of bone and joint infection (oviva): a multicentre randomised controlled trial. *Orthop Proc.* 2017;99-B(SUPP\_22):42-42. doi:10.1302/1358-992X.2017.22.042

15. Frank JM, Kayupov E, Moric M, et al. The Mark Coventry, MD, Award: Oral Antibiotics Reduce Reinfection After Two-Stage Exchange: A Multicenter, Randomized Controlled Trial. *Clin Orthop*. 2017;475(1):56-61. doi:10.1007/s11999-016-4890-4