

Inflation, quantum fields, and CMB anisotropies¹

Iván Agulló², José Navarro-Salas³

Departamento de Física Teórica and IFIC, Universidad de Valencia-CSIC, Facultad de Física, Burjassot-46100, Valencia, Spain.

Gonzalo J. Olmo⁴

*Instituto de Estructura de la Materia, CSIC, Serrano 121, 28006 Madrid, Spain and
Perimeter Institute for Theoretical Physics, Waterloo, Ontario, N2L 2Y5 Canada*

Leonard Parker⁵

*Physics Department, University of Wisconsin-Milwaukee, P.O. Box 413, Milwaukee, WI
53201, USA*

March 13, 2009

Abstract

Inflationary cosmology has proved to be the most successful at predicting the properties of the anisotropies observed in the cosmic microwave background (CMB). In this essay we show that quantum field renormalization significantly influences the generation of primordial perturbations and hence the expected measurable imprint of cosmological inflation on the CMB. However, the new predictions remain in agreement with observation, and in fact favor the simplest forms of inflation. In the near future, observations of the influence of gravitational waves from the early universe on the CMB will test our new predictions.

¹Essay written for the Gravity Research Foundation 2009 Awards for Essays on Gravitation

²ivan.agullo@uv.es

³jnavarro@ific.uv.es

⁴olmo@iem.cfmac.csic.es

⁵leonard@uwm.edu

One of the most exciting ideas of contemporary physics is to explain the origin of the observed structures in our universe as a result of quantum fluctuations in the early expanding universe. As first shown in the sixties [1], the amplification of quantum field fluctuations is an unavoidable consequence in a strongly time-dependent gravitational field [2, 3]. Fundamental physical implications were implemented some years later to culminate, in the seventies, with the prediction of the evaporation of black holes with a black-body spectrum [4] and, in the eighties, when the inflationary model of the universe was introduced [5], predicting that small density perturbations are likely to be generated in the very early universe with a nearly scale-free spectrum [6]. In the nineties, the detection of temperature fluctuations in the cosmic microwave background (CMB) by the COBE satellite [7] appeared to be consistent with the inflationary cosmology predictions. In the present decade, the predictions of inflation have been confirmed in the specific pattern of anisotropies imprinted in the full sky map of the CMB, as reported, for instance, by the WMAP mission [8]. Moreover, an inflationary-type expansion also predicts the creation of primordial gravitational waves [9], whose effects still remain undetectable. Forthcoming experiments, such as the PLANCK satellite [10], may measure effects of relic gravitational waves and offer new trends for gravitational physics in the next decade. Therefore, it is particularly important to scrutinize the quantitative predictions of quantum field theory in an inflationary background. This is the aim of this essay.

As remarked above, a strongly time-dependent gravitational field necessarily amplifies vacuum fluctuations. This happens, typically, in a rapidly expanding universe and also in a gravitational collapse. The event horizon of a black hole acts as a magnifying glass that exponentially stretches very

short wavelengths to macroscopic scales and generates a thermal flux of outgoing particles. Similarly, during exponential inflation, $ds^2 = -dt^2 + e^{2Ht}d\vec{x}^2$, a typical physical length, with comoving wavenumber k , increases exponentially $k^{-1}e^{Ht}$ and reaches the Hubble radius, $H^{-1} = \text{constant}$, at some time t_k ($ke^{-Ht_k} = H$). These quantum fluctuations produce scale-free density perturbations and relic gravitational waves via a quantum-to-classical transition at the time of Hubble horizon exit t_k . The cosmic expansion farther stretches these scale-free primordial perturbations to astronomical scales.

Let us focus on the production of relic gravitational waves by considering fluctuating tensorial modes $h_{ij}(\vec{x}, t)$ in an exponentially expanding, spatially flat universe: $g_{ij} = a^2(t)(\delta_{ij} + h_{ij})$, with $a(t) = e^{Ht}$. The perturbation field h_{ij} can be decomposed into two polarization states described by a couple of massless scalar fields $h_{+,\times}(\vec{x}, t)$, both obeying the wave equation $\ddot{h} + 3H\dot{h} - a^{-2}\nabla^2 h = 0$ (see, for instance, [11]; we omit the subindex $+$ or \times). On scales smaller than the Hubble radius the spatial gradient term dominates the damping term and leads to the conventional flat-space oscillatory behavior of modes. However, on scales larger than the Hubble radius the damping term $3H\dot{h}$ dominates. If one considers plane wave modes of comoving wavevector \vec{k} obeying the adiabatic condition [1, 2, 3] and de Sitter invariance one finds

$$h_{\vec{k}}(\vec{x}, t) = \sqrt{\frac{16\pi G}{2(2\pi)^3 k^3}} e^{i\vec{k}\vec{x}} (H - ike^{-Ht}) e^{i(kH^{-1}e^{-Ht})}. \quad (1)$$

These modes oscillate until the physical wave length reaches the Hubble horizon length. A few Hubble times after horizon exit the modes amplitude freezes out to the constant value $|h_{\vec{k}}|^2 = \frac{GH^2}{\pi^2 k^3}$. Because of the loss of phase information, the modes of the perturbations soon take on clas-

sical properties [12]. The freezing amplitude is usually codified through the quantity $\Delta_h^2(k) \equiv 4\pi k^3 |h_{\vec{k}}|^2$. Taking into account the two polarizations, one easily gets the standard scale free tensorial power spectrum [11] $P_t(k) \equiv 4\Delta_h^2(k) = \frac{8}{M_P^2} \left(\frac{H}{2\pi}\right)^2$, where $M_P = 1/\sqrt{8\pi G}$. It is easy to see that $\Delta_h^2(k)$ gives the formal contribution, per $d \ln k$, to the variance of the gravity wave fields $h_{+,\times}$

$$\langle h^2 \rangle = \int_0^\infty \frac{dk}{k} \Delta_h^2(k) . \quad (2)$$

Due to the large k behavior of the modes the above integral is divergent. It is a common view [11] to bypass this point by regarding $h(\vec{x}, t)$ as a classical random field. One then introduces a window function $W(kR)$ in the integral to smooth out the field on a certain scale R and to remove the Fourier modes with $k^{-1} < R$. One can also consider unimportant the value of $\langle h^2 \rangle$ and regard the (finite) two-point function $\langle h(x_1)h(x_2) \rangle$, uniquely defined by $\Delta_h^2(k)$, as the basic object. However, $\langle h^2 \rangle$ represents the variance of the Gaussian probability distribution associated to $h(\vec{x}, t)$, which means that at any point $h(\vec{x}, t)$ may fluctuate by the amount $\pm \sqrt{\langle h^2(\vec{x}, t) \rangle}$ defining this way a classical perturbation. It is our view to regard the variance as the basic physical object and treat h as a proper quantum field. Renormalization is then the natural solution to keep the variance finite and well-defined. Since the physically relevant quantity (power spectrum) is expressed in momentum space, the natural renormalization scheme to apply is the so-called adiabatic subtraction [13], as it renormalizes the theory in momentum space. Adiabatic renormalization [14, 2, 3] removes the divergences present in the formal

expression (2) by subtracting counterterms mode by mode in the integral (2)

$$\langle h^2 \rangle_{ren} = \int_0^\infty \frac{dk}{k} \left[4\pi k^3 |h_{\vec{k}}|^2 - \frac{16\pi G k^3}{4\pi^2 a^3} \left(\frac{1}{w_k} + \frac{\dot{a}^2}{2a^2 w_k^3} + \frac{\ddot{a}}{2a w_k^3} \right) \right], \quad (3)$$

with $w_k = k/a(t)$. The subtraction of the first term ($16\pi G k^3/4\pi^2 a^3 w_k$) cancels the typical flat space vacuum fluctuations. The additional terms, proportional to \dot{a}^2 and \ddot{a} , are necessary to properly perform the renormalization in an expanding universe.

For the idealized case of a strictly constant H , the subtractions exactly cancel out the vacuum amplitude [13], at any time during inflation, producing a vanishing result for the variance. Therefore, the physical tensorial power spectrum, the integrand of (3), is zero. Note that this surprising result does not contradict the fact that quantum fluctuations in de Sitter space produce a Hawking-type radiation with temperature $T_H = H/2\pi$ [15]. This temperature stems from the comparison of the modes (1) with those defining the vacuum of a static observer with metric $ds^2 = -(1 - H^2 r^2) dt^2 + (1 - H^2 r^2)^{-1} dr^2 + r^2 d\Omega^2$. The different time/phase behavior of both sets of modes (captured by non-vanishing Bogolubov coefficients) produces the Hawking temperature. However, their amplitudes are exactly the same [16].

Does it mean that inflation does not produce gravitational waves? No. For more realistic inflationary models, $H \equiv \sqrt{\frac{8\pi G}{3} V(\phi_0)}$ slowly decreases as ϕ_0 (the classical homogeneous part of the inflaton field) rolls down the potential towards a minimum. Tensorial perturbations are then expected, on dimensional grounds, to be produced with amplitude proportional to \dot{H} , instead of H^2 . The form of the modes is now $h_{\vec{k}}(t, \vec{x}) = (-16\pi G \tau \pi / 4 (2\pi)^3 a^2)^{1/2} \times H_\nu^{(1)}(-k\tau) e^{i\vec{k}\vec{x}}$, where the index of the Bessel function is $\nu = \sqrt{9/4 + 3\epsilon}$ and

ϵ is the slow-roll parameter $\epsilon \equiv -\dot{H}/H^2 = (M_P^2/2)(V'/V)^2$. The conformal time $\tau \equiv \int dt/a(t)$ is given here by $\tau = -(1 + \epsilon)/aH$. The loss of phase information in the modes still occurs at a few Hubble times after horizon exit, converting the fluctuations to classical perturbations. Therefore, it is natural to evaluate the new integrand of (3) (i.e, the tensorial power spectrum) a few Hubble times after the time t_k . Since the results will not be far different from those at t_k , we use the time t_k to characterize the results. The new tensorial power spectrum turns out to be then

$$P_t(k) = \frac{8\alpha}{M_P^2} \left(\frac{H(t_k)}{2\pi} \right)^2 \epsilon(t_k) \equiv -\frac{8\alpha}{M_P^2} \dot{H}(t_k) , \quad (4)$$

where $\alpha \approx 0.904$ is a numerical coefficient. As expected, it is just the deviation from an exactly constant H , parameterized by ϵ , which generates a non-zero tensorial power spectrum.

The above result would then imply that the tensor-to-scalar ratio $r = P_t(k)/P_{\mathcal{R}}(k)$ may be well below the standard predictions of single-field inflationary models. However, this is not necessarily the case since the scalar power spectrum $P_{\mathcal{R}}(k)$, which constitutes the seeds for structure formation, is also affected by renormalization. A detailed calculation, sketched in [17], leads to

$$P_{\mathcal{R}} = \frac{1}{2M_P^2\epsilon(t_k)} \left(\frac{H(t_k)}{2\pi} \right)^2 (\alpha\epsilon(t_k) + 3\beta\eta(t_k)) , \quad (5)$$

where $\beta \approx 0.448$ is numerical coefficient and $\eta \equiv M_P^2(V''/V)$ is the second slow-roll parameter. Note that this contrasts with the standard prediction: $P_{\mathcal{R}} = \frac{1}{2M_P^2\epsilon(t_k)} \left(\frac{H(t_k)}{2\pi} \right)^2$ [11]. Since the scalar amplitude is also modulated by

the slow-roll parameters the ratio r is given by

$$r = \frac{16\epsilon^2(t_k)}{\alpha\epsilon(t_k) + 3\beta\eta(t_k)}, \quad (6)$$

which contrasts with the standard result $r = 16\epsilon(t_k)$. To translate this difference to a closer empirical level we have to introduce the scalar and tensorial spectral indices $n_s \equiv 1 + d \ln P_{\mathcal{R}}/d \ln k$, $n_t \equiv d \ln P_t/d \ln k$, and the running tensorial index $n'_t \equiv dn_t/d \ln k$. The standard expression for the relation between the ratio $r \equiv P_t/P_{\mathcal{R}}$ and spectral indices (consistency condition) is: $r = -8n_t$. It is expected to be verified by any single-field slow-roll inflationary model, irrespective of the particular form of the potential. However, if we invoke renormalization we get a more involved consistency condition $r = r(n_t, n_s, n'_t)$. For instance, in the simplest case of $n'_t \approx 0$, and approximating $\alpha \approx 2\beta$, the new consistency condition becomes

$$r = 1 - n_s + \frac{96}{25}n_t + \frac{11}{5}\sqrt{(1 - n_s)^2 + \frac{96}{25}n_t^2}. \quad (7)$$

Note that this expression allows for a null tensorial tilt $n_t \approx 0$ while being compatible with a non-zero ratio $r \approx \frac{16}{5}(1 - n_s)$.

We can compare the new predictions with the standard ones on the basis of the five year WMAP results. We find [17], see Figure 1, that the new predictions agree with observation and improve the likelihood that the simplest potential energy functions (quadratic and quartic, respectively) are responsible for driving the early inflationary expansion of the universe. The influence of relic gravitational waves on the CMB will soon come within the range of planned satellite measurements, and this will be a definitive test of the new predictions.




Figure 1: Plot of r versus n_s . The contours show the 68% and 95% CL derived from WMAP5 (in combination with BAO+SN) [8]. We consider two representative inflation models: $V(\phi) = m^2\phi^2$ (solid line), $V(\phi) = \lambda\phi^4$ (dashed line). The symbols show the prediction from each of this models in terms of the number N of e-folds of inflation for the monomial potentials. The top part corresponds to the prediction of our formulae, while the bottom one corresponds to the standard prediction.

References

- [1] Parker L., *Phys.Rev.Lett.* **21** 562 (1968); *Phys. Rev.* **183**, 1057(1969).
- [2] Parker L. and Toms D.J., *Quantum field theory in curved spacetime: quantized fields and gravity*, Cambridge University Press, (in press).
- [3] Birrel N.D. and Davies P.C.W., *Quantum fields in curved space*, Cambridge University Press, (1982).
- [4] Hawking S.W., *Comm. Math. Phys.* **43**, 199 (1975). Parker L., *Phys. Rev.* **D12**, 1519 (1975). Wald R.M., *Commun. Math. Phys.* **82**, 548 (1975).
- [5] Guth A., *Phys. Rev.* **D23**, 347 (1981). Starobinsky A.A., *Phys. Lett.* **B91**, 99 (1980). Linde, A.D., *Phys. Lett.* **B108**, 389 (1982); *Phys. Lett.* **B129**, 177 (1983). Albrecht A. and Steinhardt P. J., *Phys. Rev. Lett.* **48**,1220 (1982). Sato, K., *Mon. Not. Roy. Astron. Soc.* **195**, 467 (1981).
- [6] Mukhanov V.F. and Chibisov G.V., *JETP Letters* **33**, 532(1981). Hawking, S. W., *Phys. Lett.* **B115**, 295 (1982). Guth A. and Pi, S.-Y., *Phys. Rev. Lett.* **49**, 1110 (1982). Starobinsky, A. A., *Phys. Lett.* **B117**, 175 (1982). Bardeen, J.M., Steinhardt, P.J. and Turner, M.S., *Phys. Rev.***D28**, 679 (1983).
- [7] Smoot, G., *et al. Astrophys. J. Lett.* **396**, L1-L4 (1992).
- [8] Komatsu, E. et al, arXiv:0803.0547.
- [9] Grishchuk, L.P., *Sov. Phys. JETP* **40**, 409 (1975); Starobinsky, A.A., *JETP Lett.* **30**, 682 (1979).

- [10] Efstathiou G., Lawrence C. and Tauber J. (coordinators), ESA-SCI(2005)1.
- [11] Liddle A. R. and Lyth D.H., *Cosmological inflation and large-scale structure*, Cambridge University Press, (2000).
- [12] Kiefer C., Lohmar I., Polarski D. and Starobinsky A. A., *Class. Quant. Grav.* **24**, 1699 (2007)
- [13] Parker L., hep-th/0702216
- [14] Parker L. and Fulling S.A., *Phys. Rev. D* **9** 341 (1974).
- [15] Gibbons G. and Hawking S.W., *Phys. Rev. D* **15**, 2738 (1977).
- [16] Agulló I., Navarro-Salas J., Olmo G.J. and Parker L., *Phys. Rev. Lett.* **101**, 171301 (2008).
- [17] Agulló I., Navarro-Salas J., Olmo G.J. and Parker L., arXiv:0901.0439.