

OREGON HEALTH AUTHORITY

ANATOMY BASED SEXUALLY TRANSMITTED INFECTION SCREENING RECOMMENDATIONS

	Sexually Active Individuals with a Cervix				Sexually Active Individuals with a Penis		
	Aged <25	Aged ≥25	Pregnant <ul style="list-style-type: none"> • Aged <25 • Aged ≥25 if at increased risk¹ or high-prevalence setting² 	Living with HIV	Only has Partners with a Cervix	Has Partners with a Penis ³	Living with HIV
Chlamydia Gonorrhea* Retest 3 months after treatment	At least annually	Screen at least annually if increased risk ¹ or high prevalence setting ²	At 1st prenatal visit Perform test-of-cure 3-4 weeks after treatment CT only: Retest during 3rd trimester	At least annually Repeat as often as every 3 months if at increased risk ¹	Screen at least annually if increased risk ¹ or high prevalence setting ²	At least annually Repeat as often as every 3 months if at increased risk ¹	At least annually Repeat as often as every 3 months if at increased risk ¹
Syphilis	At least one once between ages 15-44, repeat according to risk ¹	At least one lifetime test between ages 15-44, repeat according to risk ¹	At 1st prenatal visit, at 28-32 weeks, and at delivery ⁴	At least annually Repeat as often as every 3 months if at increased risk ¹	Screen at least annually if increased risk ¹ or high prevalence setting ²	At least annually Repeat as often as every 3 months if at increased risk ¹	At least annually Repeat as often as every 3 months if at increased risk ¹
Hepatitis C	At least once between ages 18-79, repeat according to risk ⁵	At least once between ages 18-79, repeat according to risk ⁵	Screen according to risk ⁵	At least once, repeat according to risk ⁵	At least once between ages 18-79, repeat according to risk ⁵	At least once between ages 18-79, repeat according to risk ⁵	At least once, repeat according to risk ⁵
HIV	At least once between ages 13-64, repeat according to individual risk ¹	At least once between ages 13-64, repeat according to individual risk ¹	At 1st prenatal visit Repeat in 3rd trimester if increased individual risk ¹ or high prevalence setting ²		At least once between ages 13-64, repeat according to individual risk ¹	At least annually Repeat every 3-6 months as indicated by individual risk ¹	

* Screen all sites of contact reported by patient:

- ◇ CT: urethra, rectum
- ◇ GC: urethra, rectum, pharynx

¹ Individual risk factors include: previous STI, recreational drug use (particularly methamphetamine and other stimulants), multiple sex partners, condomless intercourse.

² High-prevalence settings include but are not limited to: correctional facilities, adolescent clinics, STD clinics.

³ Individuals in this category either 1) exclusively have partners with a penis or 2) have partners with a penis and partners with a cervix

⁴ Three screenings recommended in pregnancy due to ongoing syphilis epidemic and increase in congenital syphilis cases in OR.

⁵ Risk factors include: history of current or past injection drug use, intranasal drug use, or incarceration; have an unregulated tattoo; received long-term hemodialysis; receipt of a blood transfusion before 1992.

Screening transgender patients: Assess risks for STIs and HIV based on current anatomy and sexual behaviors.

