

Consent for Restylane® Injections



Indications: Restylane® is a sterile gel consisting of stabilized hyaluronic acid. Medicis, the manufacturer, states that it is biodegradable, and can be safely and completely metabolized by the body. Restylane® injections are given to correct facial wrinkles and/or lip augmentation. Restylane® has been approved by the FDA (Food and Drug Administration) for correction of facial wrinkles in the nasolabial area (nose-lips) and the folds between the cheeks and the nose/upper lip (“no-label” use). I understand that the safety and effectiveness of treating facial areas (other than the nasolabial folds) has not been studied; however, Restylane® has been used to enhance the appearance of lips in over 60 other countries. This “off-label” aspect of the treatment has been explained to me.

Alternatives: There are alternatives to Restylane® injections including no treatment, collagen for lip or other facial soft tissue augmentation, cosmetics, Botulinum Toxin A, laser skin resurfacing, chemical peels, or plastic surgery for wrinkle reduction.

Results: I understand that the actual degree of improvement cannot be predicted or guaranteed. Furthermore, I understand that the effect will gradually wear off and additional treatments may be necessary to maintain the desired effect. **Please allow a full two weeks for injections to settle and to see results.**

Side effects and complications include but are not limited to:

- *Potential allergic reaction* as with any product, allergies can develop during or after injection.
- *Injection site reaction* a lumpy or “thick” feeling at or just under the skin, and/or bruising, redness, itching, pain, tenderness, or slight swelling.
- *Injections into the Lip area* could trigger a recurrence of facial cold sores (herpes simplex infections for patients with a history of prior cold sores).

Treatment Areas Requested:

Marionette Lines Oral Commissures Nasolabial Folds Cheeks Tear Troughs Lips

PRECAUTIONS: Due to potential for an allergic reaction, Restylane® is not recommended for patients with severe allergies or a history of anaphylaxis. The risk of bruising or bleeding may be increased by **medications with anticoagulant effects, such as aspirin and non-steroidal anti-inflammatory drugs (e.g. Ibuprofen, Aleve, Motrin, Celebrex), high doses of Vitamin E and certain herbs (Ginkgo Biloba, St. John’s Wart).** **Please do not perform Restylane® injections if pregnant or breastfeeding.**

I understand the need for a topical application of anesthetic gel and/or injections for a nerve block to reduce the discomfort of the procedure. I understand the above, and have had the risks, benefits, and alternatives explained to me, and have had the opportunity to ask questions. No guarantees about results have been made. To the best of my knowledge, I am not pregnant and I am not breastfeeding. I give my informed consent for Restylane® injections today as well as future treatments as needed. I hereby authorize Renew laser and Skin, Dr. Nina Desai and/or Dr. Dhara Patel to perform injections as discussed during consultation.

Patient Signature _____ Date _____