


# LAMI™

Lamium maculatum



**LAMI™**  
**Dark Purple**



- Fast growing, multicolored leaves with large purple flowers
- Trailing ground cover plant for shady to partial shade locations
- Mature height and width is 6 inches x 12 inches (15cm x 30cm)

For additional technical growing information:

LAMI™ [CLICK HERE](#)



# LAMI™

Lamium maculatum



## PROPAGATION TIPS

- Stick on priority – Number 2 out of 4 categories



Flowering time  
Spring / Summer  
Flowering



Width  
(12" to 14")  
30 to 36 cm



Height  
(4" to 6")  
10 to 15cm

Average Time	Temperature	Hormone	Fertilization	Fungicide
5 weeks	Weeks 1 - 2 70°-72° F (21° - 22°C)	Optional	Weeks 1-2 50 ppm N	Spray fungicide to control Botrytis and bacteria day of sticking
	Weeks 3 - 5 65° - 68° (18° - 20° C)		Weeks 3-5 100 to 125 ppm N	Day of sticking & Week 2



**Irrigation specification** - Remove from mist as soon as possible. Scout specifically for Botrytis



Daminozide at 1500 – 2000 ppm week 3 and/or 5 as required  
Pinch at transplant

Rooting	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 1	5.8 / 6.2	0.80	72 / 74 F (22 / 23 C)	50 ppm N in mist			Day of sticking	
Week 2	5.8 to 6.2	.80 to .90	72 / 74 F (22 / 23 C)	50 ppm N in mist			Second fungicide app	
Week 3	5.8 to 6.2	.90 to 1.0	Cool to 70 F (21 C)	Feed 100 ppm to 125 ppm N		Daminozide as needed		
Week 4	5.8 to 6.2	.90 to 1.0	68 F (20 C)	feed 100 to 125 ppm N				
Week 5	5.8 to 6.2	.90 to 1.0	65 to 68 F (18 to 20 C)	feed 100 to 125 ppm N	None required	Daminozide as needed		Pinch at transplant

For additional technical growing information:

LAMI™ [CLICK HERE](#)



 **DANZIGER**

www.danzigeronline.com |   

# LAMI™

Lamium maculatum



## FINISHING TIPS

- Pinch at transplant or soon after transplant
- Moderate moist soils will avoid marginal browning of leaves from drying
- Avoid bright direct light
- Cool temperature finishing for best quality

Average Time (from liners)	Temperature	Pinch/Daylength Modification	Fertilization	Plant Growth Regulator
5 to 7 wks (12 cm) 5" pot	Average Day 68° to 72°F (20° to 22°C) early in crop	Pinch – Required	150 - 200 ppm N	To control stretch if finishing warm
6 to 8 wks (15 cm) 6 ½"	65° - 68° (18° - 20° C) finishing temperatures		Soil EC 0.9 to 1.0 pH 5.8 to 6.2	

Pests	Aphids	ACETAMIPRID, FLONICAMID, IMIDACLOPRID, DICHLORVOS
	Snail & Slugs	Ferric Phosphate Pellets or Sprays
Diseases	Botrytis	CYPRODINIL+FLUDIOXONIL, IPRODIONE, POLYOXIN
	Pythium	PROPAMOCARB
	Phytophthora	PROPAMOCARB



- PGR - Spray Daminozide at 1500 to 2500 ppm early on finished plants
- Maintain good airflow and allow plants to dry before nightfall
- Scout for Aphid and Slugs/Snail on regular basis

Finishing	pH	EC	Temp	Feed	Light	PGR	Fungicide	Comments
Week 6 Transplant	5.8 to 6.2	0.9 to 1.0	65 to 68 F (18 to 20 C)	feed 150 to 200 ppm		Daminozide as needed	Spray fungicide after transplant	
Week 6 to 10	5.8 to 6.2	0.9 to 1.0	65 to 68 F (18 to 20 C)	feed 150 to 200 ppm				

For additional technical growing information:

LAMI™ [CLICK HERE](#)

