

Cost-Savings of School-Based Health Centers

Research and evaluations have demonstrated that school-based health centers represent cost-effective investments of public resources:

- A study by Johns Hopkins University found that school-based health centers **reduced inappropriate emergency room use**, increased use of primary care, and resulted in **fewer hospitalizations** among regular users.¹
- A study of Medicaid-enrolled children served by a SBHC in Atlanta, Georgia found **significantly lower inpatient, nonemergency department transportation, drug, and emergency department Medicaid expenses** as compared to children without a SBHC. In 1996 the total yearly expense per individual for the SBHC was \$898.98, as compared to \$2360.46 for individuals without a SBHC.²
- The number of **hospitalizations and emergency department visits decreased for children with SBHCs** in Cincinnati schools (2.4-fold and 33.5% respectively) – with an estimated savings of nearly \$1,000 per child.³
- Students in New York City schools with SBHCs were **less likely to have been hospitalized for asthma** at least once in the past year (10.5%) compared to those in schools without SBHCs (17.1%).⁴
- In South Carolina, prevention-oriented health care provided in a SBHC **decreased emergency department visit rates** by 41% to 57% -- 18% greater than the decrease in students who did not use the SBHC.⁵
- Adolescents with access to SBHCs in Denver had **38% - 55% fewer after-hours care (emergent or urgent) visits** than those without school-based health center access.⁶
- Students in Ohio who used an SBHC reported more positive self-perceptions of their health, which correlated with **lower Medicaid costs**. SBHC patients cost Medicaid an average of \$30.40 less than comparable, non-SBHC patients.⁷
- A study that explored the cost-benefit of a nationwide SBHC program to manage childhood asthma estimated total savings for opportunity costs of work loss and premature death at \$23.13 billion.⁸

1. Santelli J, Kouzis A, Newcomer S. School-Based Health Centers and Adolescent Use of Primary Care and Hospital Care. *Journal of Adolescent Health*. 1996;19:267-275.

2. Adams EK, Johnson V. An Elementary School-Based Health Clinic: Can it Reduce Medicaid Costs? *Pediatrics*. Apr 2000;105(4 Pt 1):780-788.

3. Guo JJ, Jang R, Keller KN, McCracken AL, Pan W, Cluxton RJ. Impact of School-Based Health Centers on Children with Asthma. *Journal of Adolescent Health*. Oct 2005;37(4):266-274.

4. Webber MP, Carpinello KE, Oruwariye T, Lo Y, Burton WB, Appel DK. Burden of Asthma in Inner-City Elementary Schoolchildren: Do School-Based Health Centers Make a Difference? *Archives of pediatrics & adolescent medicine*. Feb 2003;157(2):125-129.

5. Key JD, Washington EC, Hulseley TC. Reduced Emergency Department Utilization Associated with School-Based Clinic Enrollment. *Journal of Adolescent Health*. Apr 2002;30(4):273-278.

6. Kaplan D, Calonge B, Guernsey B, Hanrahan M. Managed Care and School-based Health Centers: Use of Health Services. *Archives of Pediatric and Adolescent Medicine*. 1998;152:25-33.

7. Wade TJ, Guo JJ. Linking Improvements in Health-Related Quality of Life to Reductions in Medicaid Costs Among Students Who Use School-Based Health Centers. *American journal of public health*. Sep 2010;100(9):1611-1616.

8. Tai T, Bame SI. Cost-Benefit Analysis of Childhood Asthma Management Through School-Based Clinic Programs. *J Community Health*. Apr 2011;36(2):253-260.