




University of Massachusetts Dartmouth
 School for Marine Science and Technology
 836 South Rodney French Boulevard
 New Bedford, MA 02844



Final Report

2019 Drop Camera Survey of Benthic Communities and Substrate in Vineyard Wind Lease Area OCS-A 0522 Study Area



Submitted to:
 Vineyard Wind LLC
 700 Pleasant Street
 New Bedford, MA 02740

January 2020

SMAST-CE-REP-2020-075

Final Report

2019 Drop Camera Survey of Benthic Communities and Substrate in Vineyard Wind Lease Area OCS-A 0522 Study Area

Principal Investigators: N. David Bethoney, PhD. & Kevin D. E. Stokesbury, Ph.D.

Report Co-Author: Caitlyn Riley

Address: School for Marine Science and Technology,
University of Massachusetts Dartmouth,
836 S. Rodney French Blvd.
New Bedford, MA 02744

Phone Number: (508) 910-6373, (508) 910-6374

E-mail: nbethoney@umassd.edu, kstokesbury@umassd.edu

Date: January 17, 2020

You may cite this report as:

Bethoney, N.D., Riley, C., & Stokesbury, K.D.E. (2020). 2019 Drop Camera Survey of Benthic Communities and Substrate in Vineyard Wind Lease Area OCS-A 0522 Study Area. University of Massachusetts Dartmouth - SMAST, New Bedford, MA. SMAST-CE-REP-2020-075. 40 pp.

SMAST-CE-REP-2020-075

Project Summary: The University of Massachusetts Dartmouth School for Marine Science and Technology (SMAST) conducted drop camera surveys to examine the benthic community and substrate in Vineyard Wind's Outer Continental Shelf (OCS) Lease Area OCS-A 0522 (522 Study Area). The primary goal of this project was to collect preliminary data to help determine the sampling intensity needed to collect enough baseline data for environmental assessment of windfarm development impact. Our objectives were to provide:

- 1) distribution and density estimates of dominant benthic megafauna and,
- 2) classify substrate across the survey domain.

We utilized a centric systematic sampling design to sample 22 stations in the 522 Study Area. Stations were placed 5.6 km apart following a grid design. At each station a sampling pyramid was deployed, and a high-resolution camera was used to take four quadrats (2.3 m² images) samples. The area was surveyed in July and October 2019 using a commercial scallop vessel to deploy the sampling pyramid. Eighteen different benthic animal groups were observed in the 522 Study Area. The most common groups were similar to the dominant groups in other OCS Study Areas, but sea stars, moon snails, and moon snail eggs cases were observed at higher frequencies. Decreases in animal occurrence in October, except skates, were observed but wide confidence intervals are associated with most estimates. Sand was the substrate type at the majority of stations, but gravel was present at 5 stations during July. In October, only sand was observed.

Table of Contents

List of Tables	5
List of Figures	6
Introduction	8
Goal and Objectives	11
Methods	12
Results and Discussion	14
References	39
Appendix I	42

List of Tables

Table 1. The most frequently observed benthic animal groups, in order of most to least quadrats present, during the 2019 SMAST drop camera survey of OCS 522	15
---	----

List of Figures

Figure 1. Spatial extent of SMAST drop camera surveys in the northern hemisphere.....	9
Figure 2. Example of digital still image taken by the SMAST drop camera survey in complex habitat of the Rhode Island Wind Energy Lease Area on Cox’s Ledge during a survey in 2013	10
Figure 3. Drop camera survey station grids and Wind Energy Lease Areas	12
Figure 4. SMAST drop camera survey pyramid with cameras and lights used for data collection	13
Figure 5. Density of common benthic animals and associated 95% confidence intervals in the July (J) and October (O) 2019 drop camera survey of the OCS 522 Study Area.....	16
Figure 6. The average number of quadrats benthic animals were present in at each station during the July (J) and October (O) 2019 drop camera survey of the OCS 522 Study Area.....	17
Figure 7. The distribution of sea stars in the 2019 July drop camera survey of OCS 522 Study Area.....	18
Figure 8. The distribution of sea stars in the 2019 October drop camera survey of OCS 522 Study Area	19
Figure 9. The distribution of crabs in the 2019 July drop camera survey of OCS 522 Study Area	20
Figure 10. The distribution of crabs in the 2019 October drop camera survey of OCS 522 Study Area.....	21
Figure 11. The distribution of hermit crabs in the 2019 July drop camera survey of OCS 522 Study Area	22
Figure 12. The distribution of hermit crabs in the 2019 October drop camera survey of OCS 522 Study Area	23
Figure 13. The distribution of red hake in the 2019 July drop camera survey of OCS 522 Study Area.....	24
Figure 14. The distribution of red hake in the 2019 October drop camera survey of OCS 522 Study Area	25
Figure 15. The distribution of skates in the 2019 July drop camera survey of OCS 522 Study Area.....	26
Figure 16. The distribution of skates in the 2019 July drop camera survey of OCS 522 Study Area.....	27
Figure 17. The distribution of moonsnails in the 2019 July drop camera survey of OCS 522 Study Area	28
Figure 18. The distribution of moonsnail egg cases in the 2019 July drop camera survey of OCS 522 Study Area	29
Figure 19. The distribution of silver hake in the 2019 July drop camera survey of OCS 522 Study Area	30

Figure 20. The distribution of sand dollars in the 2019 July drop camera survey of OCS 522 Study Area	31
Figure 21. The distribution of sand dollars in the 2019 October drop camera survey of OCS 522 Study Area	32
Figure 22. The distribution of anemones in the 2019 October drop camera survey of OCS 522 Study Area	33
Figure 23. The distribution of bryozoans or hydrozoans in the 2019 October drop camera survey of OCS 522 Study Area	34
Figure 24. The distribution of holes (burrowing animals) in the 2019 July drop camera survey of OCS 522 Study Area.....	35
Figure 25. The distribution of holes (burrowing animals) in the 2019 July drop camera survey of OCS 522 Study Area.....	36
Figure 26. The distribution of substrate types in the October 2019 drop camera survey of OCS 522 Study Area	37
Figure 27. The distribution of substrate types in the October 2019 drop camera survey of OCS 522 Study Area	38

Introduction

In 2019, Vineyard Wind LLC leased a 516 km² area for renewable energy development on the Atlantic OCS (OCS-A) named Lease Area OCS-A 0522, located south of Nantucket, Massachusetts off the south coast of Massachusetts. Vineyard Wind is conducting fisheries surveys within Lease Area OCS-A 0522 (the “522 Study Area,”) which is the focus of this report. Vineyard Wind is also conducting fisheries studies within the northern portion of Lease Area OCS-A 0501 (the “501 North Impact Area”), considered the development area, and within the southern portion of Lease Area OCS-A 0501 (the “501 South Study Area”); these studies are reported separately.

The University of Massachusetts Dartmouth School for Marine Science and Technology (SMAST) has developed a minimally invasive, image-based drop camera survey that allows for practical data collection of the epibenthic community without causing a disturbance to the seafloor. The SMAST drop camera survey can be used to better understand benthic macrofaunal community characteristics, substrate habitats, and the spatial and temporal scales of potential impacts on these communities and habitats. The survey techniques were developed collaboratively with scallop (*Placopecten magellanicus*) fishermen and apply quadrat sampling methods based on diving studies (Stokesbury and Himmelman 1993,1995). Initial surveys in the early 2000s focused on estimating the density of scallops within closed portions of the U.S. Georges Bank fishery and the survey approach has since expanded to cover most of the scallop resource in U.S. and Canadian waters ($\approx 100,000$ km², Figure 1). Information from the survey has been incorporated into the scallop stock assessment through the Stock Assessment Workshop process and reliably provided to the New England Fisheries Management Council to aid in annual scallop harvest allocation (NEFSC 2010, 2018).




Figure 1. Spatial extent of SMAST drop camera surveys in the northern hemisphere. All stations surveyed since 1999 are displayed.

Data from the drop camera survey has contributed in numerous ways to understanding the ecology of non-scallop species (Marino et al. 2009, MacDonald et al. 2010, Bethoney et al 2017, Ascii et al. 2018, Rosellon-Druker and Stokesbury 2019) and the characterization of benthic habitat (Stokesbury and Harris 2006, Harris and Stokesbury 2010, NEFMC 2011, Harris et al. 2012). This work contributed to several ecosystem-based management activities such as the New England Fisheries Management Council Swept Area Seabed Impact model (NEFMC 2011). Drop camera surveys have also been used to define habitat characteristics and spatial distribution of benthic marine invertebrates in potential wind energy areas off the coasts of Maryland and southern New England (Guida et al 2017). Ecologically and economically important species that would be difficult to sample with a net or dredge, such as longfin squid (*Doryteuthis pealeii*) egg clusters or habitat forming filamentous fauna (bryozoans or hydrozoans), can be counted using the drop camera survey (Figure 2).




Figure 2. Example of digital still image taken by the SMAST drop camera survey in complex habitat of the Rhode Island Wind Energy Lease Area on Cox’s Ledge during a survey in 2013. A longfin squid (*Doryteuthis pealeii*) egg cluster is present (top, middle).

The data collected by the drop camera survey can be used in an impact assessment to determine whether a change to the environment occurred due to a specific stressor, such as wind development, and to what extent the components are affected (Smith 2006). The Before-After Control-Impact (BACI) study is an experiment designed for assessing anthropogenic impacts on natural habitats and is particularly useful in large-scale anthropogenic disturbances or environmental management (Green 1979; Underwood 1991; Kerr et al. 2019). To account for naturally fluctuating characteristics, a designated area outside of the impact area, but containing similar environments and communities, is chosen to be the control site (Eberhardt 1976). The approach is strengthened with an asymmetrical design that uses multiple control sites at different distances from the impact site, incorporating the concepts of Beyond BACI (Underwood 1993) and Before After Gradient (Ellis and Scheider 1997). The standardized, systematic approach of the drop camera survey allows each survey the potential to become a dataset integrated into this design with the ultimate goal of comparing epibenthic faunal variance between impact and control sites over time. Based on the drop camera survey’s history and this analytical approach, drop camera surveys within and near areas slated for offshore wind energy development will aid in building a regional, standardized baseline dataset needed to address

development impacts on epibenthic communities and habitats. From this study the data collected can be used as a preliminary estimate enabling the calculation of a power analysis detailing the number of samples required to detect a significant change with a specific level of precision. This will enable an accurately designed BACI experiment prior to development of the area.

Goal and Objectives

The primary goal of this project was to collect preliminary data on the benthic community and substrate in the southern portion of Vineyard Wind's OCS Lease Area OCS-A 0522 (522 Study Area). This data could be used to help determine the sampling intensity needed to collect enough baseline data for environmental assessment of windfarm development impact in the 522 Study Area. To do this we used information from drop camera surveys of the 522 Study Area during two different time periods to (Figure 3):

- 1) Map the distribution and estimate the density of dominant benthic megafauna
- 2) Classify substrate

These two objectives documented the primary epibenthic animals and habitats within the 522 Study Area and help identify the sampling intensity needed for future analysis of variance. They also document seasonal changes in distribution and density.




Figure 3. 2019 Drop camera survey station grids and Wind Energy Lease Areas.

Methods

We utilized a centric, systematic sampling design to sample survey stations in the 522 Study Area. Stations were placed 5.6 km apart following a grid design (Figure 3). At each station a sampling pyramid was deployed, and a high-resolution camera was used take four quadrat (2.3 m² images) samples (Figure 4). This provided the same sampling resolution as the 2012 and 2013 drop camera surveys of Massachusetts Wind Energy Areas. The information from these surveys was used to determine the 1.5 km station distance applied to the 501 North Impact and Control areas (Figure 3).




Figure 4. SMAST drop camera survey pyramid with cameras and lights used for data collection. The camera used for the small view was turned to the side to provide a view parallel to the seafloor for some stations.

At each station, we deployed the drop camera pyramid (Figure 4) affixed with cameras and lights to the seafloor from a commercial fishing vessel (Stokesbury 2002, Stokesbury et al. 2004, Bethoney and Stokesbury 2018). A mobile studio including monitors, computers for image capturing and data entry, and survey navigation (software integrated with the differential global positioning system) was assembled in the vessel's wheelhouse. The two downward facing cameras mounted on the sampling pyramid provided 2.3 m² and 2.5 m² quadrat images of the sea floor for all stations. Additionally, a third camera providing a 0.6 m² view or view parallel to the seafloor was also deployed. Images from all cameras and video footage from the 2.5 m² quadrat view of the first quadrat was saved and then the pyramid was raised, so the seafloor could no longer be seen. The vessel was allowed to drift approximately 50 m and the pyramid was lowered to the seafloor again to obtain a second quadrat; this was repeated four times, so that each station had four images from each camera. Onboard the survey vessel, scallop counts, station location, and depth were recorded and saved through a specialized field application for entry into a SQL Server Relational Database Management System.

After the survey, the high resolution digital still images were used as the primary data source (Figure 2). Other images and video collected were used as aids. Within each quadrat, macrobenthos were counted or noted as present, and the substrate was identified (Stokesbury 2002, Stokesbury et al. 2004, Bethoney and Stokesbury 2018). Fifty taxa of macrobenthos are identified if present in the sample, counted or noted as present or absent (Appendix I). For animals noted as present, the percent of a quadrat they were present within was calculated by portioning the quadrat into equal sized cells and recording presence or absence for each cell. In addition, longfin squid egg clusters (*Doryteuthis pealeii*), which are not typically enumerated, were counted. Sediments were visually identified following the Wentworth particle grade scale from images, where the sediment particle size categories (in grain diameters) are based on a doubling or halving of the fixed reference point of 1 mm; sand = 0.0625 to 2.0 mm, gravel = 2.0 to 256.0 mm and boulders > 256.0 mm (Lincoln et al. 1992). Gravel was divided into two categories, granule/pebble = 2.0 to 64.0 mm and cobble = 64.0 to 256.0 mm (Lincoln et al. 1992). The presence of each sediment category was noted for each image. Maps and analysis focused on classifying stations by the largest sediment particle size observed in a digital still image from that station (Harris and Stokesbury 2010). Shell debris was also identified. After the images were digitized, a quality assurance check was performed on each image to ensure accuracy of counted and identified species and sediments.

Mean densities and standard errors of animals counted were calculated using equations for a two-stage sampling design where the mean of the total sample is (Cochran 1977):

$$\bar{x} = \sum_{i=1}^n \left(\frac{\bar{x}_i}{n} \right)$$

where n is the number of stations and \bar{x}_i is the mean of the 4 quadrats at station i . The SE of this 2-stage mean was calculated as:

$$S.E.(\bar{x}) = \sqrt{\frac{1}{n}(s^2)}$$

where:

$$s^2 = \sum_{i=1}^n (\bar{x}_i - \bar{x})^2 / (n - 1)$$

According to Cochran (1977) and Krebs (1989) this simplified version of the 2-stage variance is appropriate when the ratio of sample area to survey area (n/N) is small. In this case, thousands of square meters (n) are sampled compared with millions of square meters (N) in the study area. A similar multi-stage approach was used to calculate mean presence values. Mean density or quadrats present per station within the 522 Study Area were mapped (Figures 7-27). The analysis was limited to the 12 most common benthic animal groups in the 522 Study Area, to focus results on the groups detected at high enough rates for future analysis of variance (Bethoney et al. 2017). Densities for animal group were compared by graphing mean estimates with their associated 95% confidence intervals (Sokal and Rohlf 2012).

Results and Discussion

The two drop camera surveys of the 522 Study Area were conducted on July 18th and October 19th, 2019. All images and video collected were shared with Vineyard Wind. All 22 stations in the area were surveyed in July and October. Whelks. Eighteen different benthic animal groups were observed in the 522 Study Area. The most common groups were similar to the dominant groups in the 501N Impact and Control Areas, but sea stars, moon snails, and moon snail eggs cases replaced waved whelks, sponges, and skate egg cases (Table 1). In addition to these animals several others were observed at lower frequencies. These animals included scallops (1 in October) and flat fishes (2 in July and 1 in October). Decreases in animal occurrence in October, except skates, was observed but wide confidence intervals are associated with most estimates (Figures 5 & 6). Confidence intervals were extremely large for sea stars as they were found at one station in July and October at very high densities (Figures 7 & 8). Crabs also had a large confidence interval compared to the other animals, but high-density stations were spread through the survey area (Figures 9 & 10)). All other animals appear to be common at depths shallower than 50 meters, except burrowing animals (Figures 11-25). Sand was the largest substrate type at the majority of stations, but gravel was present at 5 stations during July (Figure 26). In October, only sand was observed (Figure 27).

Table 1. The 12 most common benthic animal groups, in order of most to least quadrats present, during the 2019 SMAST drop camera survey of the 522 Study Area. Groups left blank in the “Counts” column are tracked as present or absent.

Animal Group	Quadrats Present	Counts
Holes (burrowing animals)	55	
Crabs (cancer spp.)	34	75
Hermit Crabs	18	22
Sand Dollars	17	
Red hake	13	21
Silver hake	13	13
Sea Stars	10	141
Anemones	6	
Skates	6	6
Moonsnails	5	6
Moonsnail Egg Cases	5	5
Bryozoans/Hydrozoans	4	




Figure 5. Density of common benthic animals found in the July (J) and October (O) 2019 drop camera surveys of the OCS 522 Study Area. Error bars are 95% confidence intervals.




Figure 6. The average number of quadrats benthic animals were present in at each station during the July (J) and October (O) 2019 drop camera surveys of the OCS 522 Study Area. Holes represent burrowing animals and Bry./Hyd. indicates bryozoans and hydrozoans. Four quadrats (2.3m² images) were observed at each station.




Figure 7. The distribution of sea stars in the 2019 July drop camera survey of OCS 522 Study Area. Density categories equally divide the data above zero based on observations in July and October.




Figure 8. The distribution of sea stars in the 2019 October drop camera survey of OCS 522 Study Area. Density categories equally divide the data above zero based on observations in July and October.




Figure 9. The distribution of crabs in the 2019 July drop camera survey of OCS 522 Study Area. Density categories equally divide the data above zero based on observations in July and October.




Figure 10. The distribution of crabs in the 2019 October drop camera survey of OCS 522 Study Area. Density categories equally divide the data above zero based on observations in July and October.




Figure 11. The distribution of hermit crabs in the 2019 July drop camera survey of OCS 522 Study Area. Density categories equally divide the data above zero based on observations in July and October.




Figure 12. The distribution of hermit crabs in the 2019 October drop camera survey of OCS 522 Study Area. Density categories equally divide the data above zero based on observations in July and October.




Figure 13. The distribution of red hake in the 2019 July drop camera survey of OCS 522 Study Area. Density categories equally divide the data above zero based on observations in July and October.




Figure 14. The distribution of red hake in the 2019 October drop camera survey of OCS 522 Study Area. Density categories equally divide the data above zero based on observations in July and October.




Figure 15. The distribution of skates in the 2019 July drop camera survey of OCS 522 Study Area. Density categories represent zero, one, or two skates observed at a station.




Figure 16. The distribution of skates in the 2019 July drop camera survey of OCS 522 Study Area. Density categories represent zero, one, or two skates observed at a station.




Figure 17. The distribution of moonsnails in the 2019 July drop camera survey of OCS 522 Study Area. No moonsnails were observed in an October survey of the same area. Density categories represent zero, one, two or three moonsnails observed at a station.




Figure 18. The distribution of moon snail egg cases in the 2019 July drop camera survey of OCS 522 Study Area. No moon snail egg cases were observed in an October survey of the same area. Density categories represent zero, one or two moon snail egg cases observed at a station.




Figure 19. The distribution of silver hake in the 2019 July drop camera survey of OCS 522 Study Area. No silver hake were observed in an October survey of the same area. Density categories represent zero, one, two or three silver hake observed at a station.




Figure 20. The distribution of sand dollars in the 2019 July drop camera survey of OCS 522 Study Area. Four quadrats (2.3m^2 images) were observed at each station.




Figure 21. The distribution of sand dollars in the 2019 October drop camera survey of OCS 522 Study Area. Four quadrats (2.3m^2 images) were observed at each station.




Figure 22. The distribution of anemones in the 2019 October drop camera survey of OCS 522 Study Area. No anemones were observed in an October survey of the same area. Four quadrats (2.3m² images) were observed at each station.




Figure 23. The distribution of bryozoans or hydrozoans in the 2019 October drop camera survey of OCS 522 Study Area. No bryozoans or hydrozoans were observed in an October survey of the same area. Four quadrats (2.3m^2 images) were observed at each station.




Figure 24. The distribution of holes (burrowing animals) in the 2019 July drop camera survey of OCS 522 Study Area. Four quadrats (2.3m^2 images) were observed at each station.




Figure 25. The distribution of holes (burrowing animals) in the 2019 July drop camera survey of OCS 522 Study Area. Four quadrats (2.3m^2 images) were observed at each station.




Figure 26. The distribution of substrate types in the October 2019 drop camera survey of OCS 522 Study Area. Four quadrats (2.3m^2 images) were observed at each station.




Figure 27. The distribution of substrate types in the October 2019 drop camera survey of OCS 522 Study Area. Four quadrats (2.3m² images) were observed at each station.

References

- Asci SC, Langton RW, Stokesbury KDE (2018) Estimating similarity in benthic communities over decades and in areas open and closed to fishing in the central Gulf of Maine. *Mar. Ecol. Prog. Ser.* 595: 15-26.
- Bethoney ND, Zhao L, Chen C, Stokesbury KDE (2017) Identification of persistent benthic assemblages in areas with different temperature variability patterns through broad-scale mapping. *PLoS ONE* 12 (5), e0177333. <https://doi.org/10.1371/journal.pone.0177333>
- Bethoney ND, Stokesbury K (2018) Methods for Image-based Surveys of Benthic Macroinvertebrates and Their Habitat Exemplified by the Drop Camera Survey for the Atlantic Sea Scallop. *J. Vis. Exp.* (137), 57493. doi:10.3791/57493
- Cochran WG (1977) *Sampling Techniques* 3rd Edition. John Wiley & Sons, New York
- Eberhardt LL (1976) Quantitative ecology and impact assessment. *J. Environ. Manage.* 4, 27–70.
- Ellis JI, Schneider DC (1997) Evaluation of a gradient sampling design for environmental impact assessment. *Environ. Monit. Assess.* 48, 157 – 172.
- Green RH (1979) *Sampling Design and Statistical Methods for Environmental Biologists*. Wiley, Chichester, UK.
- Guida V, Drohan A, Welch H, McHenry J, Johnson D, Kentner V, Brink J, Timmons D, Estela-Gomez E (2017) *Habitat Mapping and Assessment of Northeast Wind Energy Areas*. Sterling, VA: US Department of the Interior, BOEM. OCS Study BOEM 2017-088. 312 p.
- Harris BP, Stokesbury KDE (2010) The spatial structure of local surficial sediment characteristics on Georges Bank, USA. *Cont. Shelf. Res.* 30, 1840-1853
- Harris BP, Cowles GW, Stokesbury KDE (2012) Surficial sediment stability on Georges Bank in the Great South Channel and on eastern Nantucket Shoals. *Cont. Shelf. Res.* 49, 65-72
- Kerr LA, Kritzer JP, Cadrin SX (2019) Strengths and limitations of before–after–control–impact analysis for testing the effects of marine protected areas on managed populations, *ICES J. Mar. Sci.*, fsz014, <https://doi.org/10.1093/icesjms/fsz014>.
- Krebs CJ (1989) *Ecological methodology*. Harper & Row, New York
- Lincoln RJ, Boxshall GA, Clark PF (1992) *A dictionary of ecology, evolution, and systematics*. Cambridge University Press, Cambridge.

- MacDonald AM, Adams CF, Stokesbury KDE (2010) Abundance estimates of skates (*Rajidae*) on the continental shelf of the northeastern USA using a video survey. *Trans. Am. Fish. Soc.* 139, 1415-1420
- Marino II MC, Juanes F, Stokesbury KDE (2009) Spatio-temporal variations of sea star *Asterias* spp. distributions between sea scallop *Placopecten magellanicus* beds on Georges Bank. *Mar. Ecol. Prog. Ser.* 382, 59–68
- NEFSC (Northeast Fisheries Science Center) (2010) Stock assessment for Atlantic sea scallops in 2014. In: 59th Northeast Regional Stock Assessment Workshop (59th SAW) Assessment Report. U.S. Dept. of Commerce, NEFSC Ref. Doc. 14-09
- NEFSC (2018) Sea scallop assessment summary for 2018. In: 65th Northeast Regional Stock Assessment Workshop (65th SAW) Assessment Summary Report. U.S. Dept of Commer, Northeast Fish Sci Cent Ref Doc 18-08, p 13-19
- Rosellon-Druker J, Stokesbury KDE (2019) Characterization and quantification of echinoderms (Echinodermata) on Georges Bank and the potential role of marine protected areas on these populations. *Invert. Biol.* 132(2): <https://doi.org/10.1111/ivb.12243>
- Sokal RR, Rohlf FJ (2012) Hypothesis testing and interval estimation. Pages 119–176 in M. Rolfes, editor. *Biometry*, 4th edition. Freeman, New York, New York.
- Smith E (2006) BACI Design. *Encyc. of Environmetrics*. Vol. 1, pp141-148. <https://doi.org/10.1002/9781118445112.stat07659>
- Stokesbury KDE, Himmelman JH (1993) Spatial distribution of the giant scallop *Placopecten magellanicus* in unharvested beds in the Baie des Chaleurs, Québec. *Mar. Ecol. Prog. Ser.* 96, 159-168
- Stokesbury KDE, Himmelman JH (1995) Examination of orientation of the giant scallop, *Placopecten magellanicus*, in natural habitats. *Can. J. Zool.* 73, 1945-1950.
- Stokesbury KDE (2002) Estimation of Sea Scallop Abundance in Closed Areas of Georges Bank, USA. *Trans. Am. Fish. Soc.* 131:1081–1092.
- Stokesbury KDE, Harris BP (2006) Impact of limited short-term sea scallop fishery on epibenthic community of Georges Bank closed areas. *Mar. Ecol. Prog. Ser.* Vol 307: 85-100.
- Stokesbury KDE, Harris BP, Marino MC, Nogueira JI (2004) Estimation of sea scallop abundance using a video survey in off-shore US waters. *J. Shellfish Res.* Vol 23, No. 1, 33-40.

Underwood AJ (1991) Beyond BACI: experimental designs for detecting human and environmental on temporal variations in natural populations. *Aust. J. Mar. Freshwater Res.* 42: 569–587.

Underwood AJ (1993) The mechanics of spatially replicated sampling programmes to detect environmental impacts in a variable world. *Australian J. Ecol.* 18: 99–116.

Appendix I: Georges Bank species are grouped into taxonomic categories (Stokesbury and Harris, 2006).

Category	Scientific name	Common name	
Scallop	<i>Placopecten magellanicus</i>	Sea scallop	
Starfishes	<i>Solaster endeca</i>	Purple sunstar	
	<i>Crossaster papposus</i>	Spiny sunstar	
	<i>Leptasterias Polaris</i>	Polar sea star	
	<i>Asterias spp.</i>	Sea stars	
	<i>Henricia spp</i>	Blood star	
Sand dollars	<i>Echinarachnius parma</i>	Sand dollar	
Bryozoans/hydrozoans	<i>Flustra foliacea</i>	Bryozoans	
	<i>Callopora aurita</i>	Bryozoans	
	<i>Electra monostachys</i>	Bryozoans	
	<i>Cribrilina punctate</i>	Bryozoans	
	<i>Eucretea loricate</i>	Bryozoans	
	<i>Tricellaria ternate</i>	Bryozoans	
	<i>Eudendrium capillare</i>	Hydrozoans	
	<i>Sertularia cupressina</i>	Sea cypress hydroid	
	<i>Sertularia argentea</i>	Squirrel's tail hydroid	
	<i>Diphasia fallax</i>	Hydrozoans	
	<i>Filograna implexa</i>	Lacy tube worm	
	Sponges	<i>Suberites ficus</i>	Fig sponge
		<i>Haliclona oculata</i>	Finger sponge
		<i>Halichondria panacea</i>	Crumb of bread sponge
		<i>Cliona celata Grant</i>	Boring sponge
		<i>Polymastia robusta</i>	Encrusting sponge
<i>Isodictya palmate</i>		Palmate sponge	
<i>Microciona prolifera</i>		Red beard sponge	
<i>Homarus americanus</i>		American lobster	
Crabs		<i>Cancer irroratus Say</i>	Atlantic rock crab
		<i>Cancer borealis Stimpson</i>	Jonah crab
Hermit crabs		<i>Diogenidae</i>	Left-handed hermit crabs
	<i>Paguridae</i>	Right-handed hermit crabs	
Eel pout	<i>Parapaguridae</i>	Deep water hermit crabs	
Flounder	<i>Zoarces americanus</i>	Ocean pout	
	<i>Paralichthys dentatus</i>	Summer flounder	
	<i>Paralichthys oblongus</i>	Fourspot flounder	
	<i>Scophthalmus aquosus</i>	Windowpane flounder	
	<i>Pseudopleuronectes americanus</i>	Winter flounder	
	<i>Limanda ferruginea</i>	Yellowtail flounder	
	<i>Glyptocephalus cynoglossus</i>	Witch flounder	
	<i>Trinectes maculatus</i>	Hogchoaker	
	Haddock	<i>Melanogrammus aeglefinus</i>	Haddock
		<i>Merluccius bilinearis</i>	Silver hake
Hake	<i>Urophycis spp.</i>	Red and white hake	
	<i>Myoxocephalus octodecemspinosus</i>	Longhorn sculpin	
Sculpins	<i>Prionotus carolinus</i>	Northern sea robin	
Skates	<i>Leucoraja erinacea</i>	Little skate	
	<i>Leucoraja ocellata</i>	Winter skate	
	<i>Dipturus laevis</i>	Barndoor skate	
Other fish	<i>Myxine glutinosa</i>	Atlantic hagfish	
	<i>Scyliorhinus rotifer</i>	Chain dogfish	
	<i>Squalus acanthias</i>	Spiny dogfish	
	<i>Anguilla rostrate</i>	American eel	
	<i>Conger oceanicus</i>	Conger eel	
	<i>Clupea harengus</i>	Atlantic herring	
	<i>Brosme brosme</i>	Cusk	
	<i>Gadus morhua</i>	Atlantic cod	
	<i>Lophius americanus</i>	Goosefish	
	<i>Ammodytes dubius</i>	Northern sand lance	
	<i>Scomber scombrus</i>	Atlantic mackerel	
	<i>Sebastes fasciatus</i>	Acadian refish	
	<i>Anarhichas lupus</i>	Atlantic wolffish	
	Shell debris	<i>Buccinum undatum</i>	Waved whelk
		<i>Euspira heros</i>	Northern moonshell
		<i>Mercenaria mercenaria</i>	Northern quahog
<i>Modiolus modiolus</i>		Northern horse mussel	
<i>Ensis directus</i>		Atlantic jackknife	
	<i>Placopecten magellanicus</i>	Sea scallops	