

# FACT SHEET

## Maternal Mental Health (MMH)

### KEY POINTS

- > Maternal mental health (MMH) conditions are the most common complications of pregnancy and childbirth, affecting 1 in 5 women (800,000 women each year in the United States).<sup>1-3</sup>
- > MMH conditions include depression, anxiety disorders, obsessive-compulsive disorder, post-traumatic stress disorder, bipolar illness (which may include psychotic symptoms), and substance use disorders.<sup>1-3</sup>
- > The “baby blues” are a normal period of transition affecting up to 85% of new mothers in the first 2-3 weeks after baby is born. Baby blues typically include emotional sensitivity, weepiness, and/or feeling overwhelmed. Baby blues resolve without treatment.<sup>4</sup>
- > MMH conditions are caused by a combination of changes in biology, psychology, and environment.<sup>4</sup>
- > Women at increased risk of MMH conditions are those who have a personal or family history of mental illness; lack social support, especially from their partner; experienced a traumatic birth or previous trauma in their lives; or have a baby in the neonatal intensive care unit.<sup>1,4</sup>
- > Suicide and overdose are the leading causes of death in the first year postpartum, with 100% of these deaths deemed preventable.<sup>5-7</sup>
- > All parents — including fathers, partners, and adoptive parents — can experience changes in mood when there is a new baby in the household.<sup>8,9</sup>

**1/5**



women will experience MMH conditions during pregnancy or first year following pregnancy<sup>1-3</sup>



of women who experience MMH symptoms go untreated<sup>10</sup>

#### Cost of not treating MMH conditions

is \$32,000 per mother-infant pair (adding up to \$14 billion nationally)<sup>11</sup>



### SIGNS & SYMPTOMS

DEPRESSION	ANXIETY
<ul style="list-style-type: none"> <li>• Feeling hopeless, helpless, or worthless</li> <li>• Lacking motivation, concentration, or energy</li> <li>• Loss of interest or pleasure in activities</li> <li>• Feelings of anger, guilt, irritability, rage, or regret</li> </ul>	<ul style="list-style-type: none"> <li>• Feeling easily stressed, worried, or overwhelmed</li> <li>• Being hypervigilant with baby</li> <li>• Having scary, intrusive, or racing thoughts</li> <li>• Feeling keyed up, on edge, restless, or panicked</li> </ul>

#### Women experiencing MMH conditions might say...

- Having a baby was a mistake.
- I'm not bonding with my baby.
- I'm afraid to be alone with my baby.
- I'm exhausted, but I can't sleep, even when my baby sleeps.
- I'm such a bad mother; my baby would be better off without me.

### WOMEN AT INCREASED RISK

**Women living in poverty and women of color are MORE likely to experience MMH conditions and LESS likely to get help due to:**<sup>12, 13</sup>

- Lack of access to healthcare, including culturally appropriate mental health care
- Cultural and racial biases in the healthcare system
- More barriers to care, such as lack of transportation or childcare
- Fear that child protective services or immigration agencies will become involved



### TERMINOLOGY

**Perinatal:** ~2-year timeframe from conception to baby's first birthday

**Antenatal or Prenatal:** During pregnancy

**Postpartum or Postnatal:** First year following pregnancy

The following terms are used interchangeably to describe the mental health conditions women experience during pregnancy and the first year following pregnancy:

- Postpartum depression (PPD) has long been used as an umbrella term encompassing mood changes following childbirth
- Antenatal / prenatal / perinatal / postnatal depression and anxiety
- Perinatal mood disorders (PMDs) or perinatal mood and anxiety disorders (PMADs)
- Maternal mental health (MMH) challenges / complications / conditions / disorders / illnesses / issues

# CONSEQUENCES OF UNTREATED MMH CONDITIONS

Untreated MMH conditions can have long-term negative impact on mother, baby, and entire family.

MOTHER	CHILD
<p><b>Women with untreated MMH conditions are more likely to:</b><sup>4, 14-17</sup></p> <ul style="list-style-type: none"> <li>• Not manage their own health</li> <li>• Have poor nutrition</li> <li>• Use substances such as alcohol, tobacco, or drugs</li> <li>• Experience physical, emotional, or sexual abuse</li> <li>• Be less responsive to baby's cues</li> <li>• Have fewer positive interactions with baby</li> <li>• Experience breastfeeding challenges</li> <li>• Question their competence as mothers</li> </ul>	<p><b>Children born to mothers with untreated MMH conditions are at higher risk for:</b><sup>12, 16-19</sup></p> <ul style="list-style-type: none"> <li>• Low birth weight or small head size</li> <li>• Pre-term birth</li> <li>• Longer stay in the NICU</li> <li>• Excessive crying</li> <li>• Impaired parent-child interactions</li> <li>• Behavioral, cognitive, or emotional delays</li> </ul> <p>Untreated mental health issues in the home may result in an Adverse Childhood Experience, which can impact the long-term health of the child.<sup>20</sup></p>




**Parents who are depressed or anxious are more likely to:**<sup>15,16</sup>

- > Make more trips to the emergency department or doctor's office
- > Find it particularly challenging to manage their child's chronic health conditions
- > Not follow guidance for safe infant sleep and car seat usage

**Most maternal mental health conditions are temporary & treatable.**

## STEPS TO WELLNESS<sup>17</sup>


LOWER COST  
LOWER BARRIER



### Editorial Team

A multidisciplinary editorial team provided input for this Fact Sheet representing the fields of obstetrics, pediatrics, nursing, psychiatry, psychology, and public health. Team members from MMHLA are Adrienne Griffen, MPP; Pooja Lakshmin, MD; Kelly Sheppard, PhD; and Terri Wright, PhD, MPH. Additional editorial team members include Nancy Byatt, DO, MBA; Wendy Davis, PhD; Sue Kendig, JD, WHNP; Tiffany Moore Simas, MD, MPH; and Debra Waldron, MD, MPH.

**Of women who experience anxiety or depression during pregnancy or first year of baby's life<sup>21</sup>**



### Citations

- 1 ACOG Committee Opinion 757 (2018).
- 2 Gavin (2005). *Obstetrics & Gynecology*, 106, 1071-83.
- 3 Fawcett (2019). *Journal of Clinical Psychiatry* (80)4.
- 4 National Institute of Mental Health (2013). *Postpartum Depression Facts* NIH13-8000.
- 5 Davis (2019). *Pregnancy-related deaths: Data from 14 U.S. Maternal Mortality Review Committees, 2008-2017*.
- 6 Metz (2016). *Obstetrics & Gynecology*, 128(6): 1233-1240.
- 7 Goldman-Mellor (2019). *American Journal of Obstetrics & Gynecology*, 221(489): e1-9.
- 8 Da Costa (2019). *Journal of Women's Health*, 21(9): 830-836.
- 9 Foli (2016). *Journal of Affective Disorders*, 200: 293-302.
- 10 Byatt (2015). *Obstetrics & Gynecology*, 126(5): 1048-1058.
- 11 Luca (2020). *American Journal of Public Health*, 110(6): 888-896.
- 12 Grote (2010). *Archives of General Psychiatry*, 67(10): 1012-1024.
- 13 Taylor (2019). *Eliminating racial disparities in maternal and infant mortality*. Center for American Progress.
- 14 Zhou (2019). *Journal of Women's Health*, 28(8): 1068-1076.
- 15 Field (2010). *Infant Behavioral Health*, 33(1): 1-14.
- 16 Sriraman (2017). *Pediatrics in Review*, 38(12): 541-551.
- 17 Fitelson (2011). *International Journal of Women's Health*, 3: 1-14.
- 18 Cherry (2016). *International Journal of Women's Health*, 8: 233-242.
- 19 Stein (2014). *The Lancet*, 384: 1800-1819.
- 20 Felitti (1998). *American Journal of Preventative Medicine*, 14(4): 245-258.
- 21 Wisner (2013). *JAMA Psychiatry*, 7(5): 490-498.

**Download this and other fact sheets at [mmhla.org/factsheets](http://mmhla.org/factsheets)**