

TREATMENT

PA Get Help Now
1-800-662-HELP (4357)

**DBHIDS Behavioral Health
Special Initiative**
If you do not have insurance
215-546-1200

**Community Behavioral
Health (CBH)**
If you have insurance
888-545-2600

**Philadelphia 24HR Suicide
Crisis & Intervention Line**
215-685-6440

Find more resources at
[SubstanceUsePhilly.com/
treatment.](https://www.substanceusephilly.com/treatment)

MOBILE SITES

 **PREVENTION
POINT**

2913-15 Kensington Ave.
Philadelphia, PA 19134
ppponline.org

Services available

- OD prevention education
- Naloxone (Narcan),
fentanyl test strips, and
xylazine test strips
- HCV and HIV screening
- Medical Triage

RESOURCES & SERVICES

**Get Free Narcan & Fentanyl
Test Strips at**
[SubstanceUsePhilly.com/
get-supplies.](https://www.SubstanceUsePhilly.com/get-supplies)

Using Alone?

*Anonymously connect with a
peer who can contact a friend
for you or call 911 in case you
become unresponsive:*

- Never Use Alone Hotline
877-696-1996
- SafeSpot Hotline
800-972-0590



**KNOW
WHAT'S IN
YOUR DRUGS**

HOW TO USE FENTANYL TEST STRIPS

Test Strip Directions

1. Mix fresh water with a small amount of the drug (or residue) in the empty bag/cooker you just prepped.
2. Dip the wavy-line end of the test strip in water for 15 seconds (don't pass the solid blue line).
3. Remove the fentanyl test strip and lay it on a flat surface. Results are ready in 1 minute.

Testing Pills

If you take pills orally:

- Scrape a small amount off the pill or cut the pill in half.
- Scrape from the middle for best results.
- Follow test strip directions.

If you snort:

- Crush pills and add water.
- You can also use a few grains from the pill, but testing residue is safer.

Test Results



One line = **Positive**
for fentanyl.



Two lines = **Negative**
for fentanyl.

If the second line (closest to the white end) is faint, it is negative for fentanyl.

Test is invalid if:

1. the test has no lines OR



2. the red line closest to the solid blue end is missing.



If using, try a small amount first and go slow.

Fentanyl may still be present even if the test is negative.



Always carry naloxone!

