

TREATMENT

PA Get Help Now
1-800-662-HELP (4357)

**DBHIDS Behavioral Health
Special Initiative**
If you do not have insurance
215-546-1200

**Community Behavioral
Health (CBH)**
If you have insurance
888-545-2600

**Philadelphia 24HR Suicide
Crisis & Intervention Line**
215-686-4420

Find more resources at
SubstanceUsePhilly.com/
treatment.

MOBILE SITES

 **PREVENTION
POINT**

2913-15 Kensington Ave.
Philadelphia, PA 19134
ppponline.org

Services available

- OD prevention education
- Naloxone (Narcan),
fentanyl test strips,
xylazine test strips, and
medetomidine test strips
- HCV and HIV screening
- Medical Triage

RESOURCES & SERVICES

Get Free Narcan & Test Strips
at [SubstanceUsePhilly.com/
get-supplies](https://SubstanceUsePhilly.com/get-supplies).

Using Alone?

*Anonymously connect with a
peer who can contact a friend
for you or call 911 in case you
become unresponsive:*

- Never Use Alone Hotline
877-696-1996
- SafeSpot Hotline
800-972-0590



**KNOW
WHAT'S IN
YOUR DRUGS**

HOW TO USE FENTANYL TEST STRIPS

Test Strip Directions

1. Mix 5 cc (1 tsp) of clean water with a microscop (10mg) of the drug or residue in the empty cooker or bag the drug was packaged in.
2. Dip the wavy-line end of the test strip in water for 15 seconds (don't pass the solid blue line).
3. Remove the fentanyl test strip and lay it on a flat surface. Results are ready in 1 minute.

Testing

How to test pills:

Crush the pill into powder or cut the pill in half and scrape from the middle. Use about 10mg for the test.

How to test crystals:

Crush the shards as much as possible. Do not use any shards. Use about 10mg for the test.

NOTE:

Some drugs can cause false positives. If you get a positive, add a shot glass amount of water and retest with a new test strip.

Test Results



One line = **Positive**
for fentanyl.



Two lines = **Negative**
for fentanyl.

If the second line (closest to the white end) is faint, it is negative for fentanyl.



Test is invalid if:

1. the test has no lines OR



2. the red line closest to the solid blue end is missing.



If using, try a small amount first and go slow.



Fentanyl may still be present even if the test is negative.

True to size microscop (10mg)



Always carry naloxone!

